



BIRDS
OF
SOUTH SUDAN

Gerhard Nikolaus

BRITISH MUSEUM
(NATURAL HISTORY)
25 OCT 1993
EXCHANGED
TRING LIBRARY

Birds of South Sudan

Richard Nikolaus

Academy of Science Supplement Number 3

BIRDS OF SOUTH SUDAN

Gerhard Nikolaus_k

Scopus Special Supplement Number 3

BRN 294182
AN424478


BRITISH MUSEUM
(NATURAL HISTORY)
25 OCT 1993
EXCHANGED
TRING LIBRARY

© Gerhard Nikolaus, 1989

Bosenbüttel 4, D-2859 Spieka, Federal Republic of Germany

Published by the Ornithological Sub-Committee
of the East Africa Natural History Society and
available from D.A. Turner, P.O. Box 48019,
Nairobi, Kenya.

Printed in Kenya by
English Press, Enterprise Road
P.O. Box 30127 Nairobi.



Map of the South Sudan showing main vegetational areas

CONTENTS

Introduction	1
Topography and Climate	2
Habitats	2
Status	3
Order and Nomenclature	5
Acknowledgements	5
Species Recorded	7
Gazetteer	99
References	101
Index of Scientific Names	105
Index of English Names	116

INTRODUCTION

Ornithologically, the Sudan was one of the better known parts of Africa early this century, for the mystery of the Nile had attracted natural historians for a number of decades previously. However, most early work was based in the north, and the southern areas became known mainly as a result of collecting by Emin Pasha in the 1880s and the work of F.O. Cave and J.D. MacDonald in the 1930s and 1940s. Field work in this region has always been hampered by a long rainy season, with difficult roads and supply, disease and a variety of unpleasant biting insects. Ten years ago there were still large gaps in knowledge concerning the range of races, breeding season, moult cycles and Palaearctic migration through the area.

In 1972 the Southern Region came under an independent administration based at Juba and covering the three old provinces of Equatoria, Bahr-el-Ghazal and Upper Nile. This region, which came to be called the 'South Sudan' became much more accessible at this time and offered new scope for discovery to ornithologists now equipped with a greater background understanding of African avifaunas and migration systems, and armed with the technique of mist-netting. Several observers were resident for short spells over the following decade, and the author and his wife spent the four years 1976-1980 based at Imatong, but travelled widely throughout the region whilst undertaking a comprehensive ornithological survey. Several species were added to the South Sudan list, one new subspecies (of the Spotted Ground Thrush *Turdus fischeri*) was discovered, and a better understanding of migration through the region was established.

This work briefly reviews the distribution and occurrence of 790 species which can be accepted for the South Sudan list. In addition another 61 species are discussed, either because they have been claimed in the past, but are no longer acceptable, or because they occur so close to the region's border that their future addition to the list seems highly likely. The South Sudan can boast more species than the other countries in the Sudan belt of Africa. This is due to the existence within the region of a variety of habitats, for it embraces the edge of the Ethiopian highlands, part of the Turkana desert and representatives of both West and East African rain forests. To continue the uniform treatment of ornithological literature in eastern Africa this work has been modelled on *Birds of East Africa* (Britton 1980).

Throughout their study the author and his wife were sponsored by the Alexander Koenig Museum in Bonn and the Naturkunde Museum in Stuttgart. They also received generous support from the Sudan Wildlife Administration, and it is hoped that this work will be of use in the future management and protection of a valuable resource, the beautiful birds of the South Sudan.

TOPOGRAPHY AND CLIMATE

South Sudan lies in northeastern Africa between approximately 4°N and 12°N, and between 24°E and 36°E and has an area of 684 000 km². It consists mainly of the large Nile basin known as the Sudd, but also includes part of the Ethiopian highlands, the Turkana desert, the Albertine Rift Valley, the Zaire rain forest and the West African tree savanna. The three mountains, Imatong, Dongotona and Didinga rise along the Uganda border to 3180 m, but otherwise the altitude of the mainly flat country lies between 400 and 800 m, rising towards the southern border. The White Nile and Sobat rivers bring water from Uganda and Ethiopia respectively. South Sudan is rich in cattle and game and has various National Parks and Game Reserves. Much of the country is sparsely populated open grassland or tree savanna, but fairly intensive agriculture exists along the southern border.

Most of the country has a mean annual rainfall of 800-1500 mm, but more than 2000 mm is experienced on the southern mountains and less than 400 mm near Lake Turkana. There is generally a single rainy season, lasting from April/May to September. However, in the extreme southeast, in the Turkana basin, there are two short rainy seasons, in April/May and November/December.

HABITATS

WETLANDS comprise a dominant habitat, especially during and after the rainy season. The largest area is the 'Sudd', one of Africa's most important wetlands, fed by the Nile and its major tributaries, the Bahr el Arab, Bahr el Ghazal, Lol, Sue, Jur and Sobat. The Sudd has a large inundation zone with seasonally tall grassland, a floating vegetation of reeds, grass, papyrus and water hyacinth, and a variable area of open water. The complex is seriously endangered by the construction of the Jongolei canal. Elsewhere seasonal wetlands result from the flooding of grassland.

GRASSLANDS are found in four types of major importance: The Sudd inundation zone is largely covered by tall grassland during and immediately after the rains, but this is regularly burnt off towards the end of the calendar year by the cattle-owning people. The soil in the whole Nile basin is black 'cotton' clay. Semi arid areas exist in the extreme southeast and around Renk in the north, and here there are patches of open short grassland within *Acacia* bushland. Patches of short grass also exist within the tree savanna in the southwest, in ironstone areas with poor soil. Finally, luxuriant short grass grows above 1800 m on Imatong, Dongotona and Didinga, mainly on the northern sides of the mountains. Only in recent years has this been regularly burnt in connection with the extension of agricultural activity.

BUSH - AND TREE SAVANNA: The *Acacia* bushland in the semi-arid north and southeast sometimes forms small *Acacia* forests along larger seasonal river-banks (*Khors*). More importantly, however, almost half of the South Sudan, the area west and south of the Sudd, and the Ethiopian highland, is covered by broad-leaved tree savanna or open bushland with seasonal tall grass. Nearer to the south and southwest borders this

often forms better developed woodland and local forest patches.

LOWLAND FORESTS: The only true lowland forests exist around the Imatong Mountains. The three areas concerned, the Talanga, Lotti and Laboni forests, are endangered by forestry and agricultural developments. A number of larger forest patches also exist west of the Nile. These are mainly gallery forests, but some, the so-called 'bowl forests' of Cave and MacDonald cover larger pans. The main examples are along the Kajiko River on the Aloma Plateau, the Aza Forest near Maridi, along the Zaire border near Yambio, and at Bengengai. These western forests are also threatened by development.

HIGHLAND FOREST is confined to the upper slopes of the three mountains along the Uganda border. The forests on Didinga and Dongotona are very small, but Imatong is heavily forested at all altitudes. There are three major types: *Albizia* - *Croton* - *Macaranga* from 1600 to 2400 m, *Podocarpus* up to 3000 m and a small patch of *Ericacea* on Mt. Kinyetti above 3000 m which, however, has been practically destroyed in recent years by regular fires. Most of the Imatong forest is still primary, and forestry activities, including the planting of pine and cypress, are limited to the northern edge around Gilo in the Upper Kinyetti river basin. Here, secondary growth, cultivations and plantations are the major habitats.

STATUS

1. **MIGRATORY STATUS.** The following abbreviations are given in the heading of each species: PMW = overwintering Palaearctic migrant; PMP = Palaearctic passage migrant; AM = intra-African migrant (Sudan birds have a distinct seasonal migration within the Afrotropical Region, usually across the equator); LM - local migrant (although distinct migrations are not well established the species is only present seasonally, at least in some localities); R= resident (present throughout the year and presumed to breed); V = vagrant (probably extralimital, known from very few records). A question mark indicates that the migratory status given is speculative.

2. **BREEDING STATUS.** Except in the case of Palaearctic visitors, each species is shown as either BR = breeding recorded (followed by breeding months based on observed or estimated egg-laying time) or NBR - no breeding records. When a breeding month is given in brackets, breeding is only presumed (e.g. birds in breeding condition) but not confirmed.

3. **ABUNDANCE.** The following terms are used to denote decreasing numerical categories; abundant (many seen each day in the appropriate habitat); common (a few most days); fairly common (a few per month); uncommon (a few each year) and rare. These are somewhat subjective impressions, but provide useful indications and comparisons.

DISTRIBUTION

During the period 1976 - 1980 the author visited all parts of the Region, and special effort was made to survey every major locality and habitat at least once during both the wet and dry season. Areas most often visited were Imatong, Juba, Maridi and Wau. Distributional data for the whole of the Sudan has been plotted in Atlas form using a 1° X 1° square grid (Nikolaus 1987). In the present work distributional descriptions on the South Sudan are given in greater detail, forming the bulk of the species accounts. Sufficient reference to place names and topographical features is given for a species' known range in the Region to be easily interpreted. Apart from the southern mountains and major rivers there are few striking topographical features, and distribution patterns tend to be uncomplicated. Species often range over large areas of similar habitat, so that relatively few localities need be given. For example: 'southwest of Aweil - Wau - Juba' would imply occurrence throughout the southwestern tree savanna area. The species accounts need to be read in conjunction with Maps 1 and 2, and with the gazetteer. Compass points and 'central' are usually abbreviated to N, S, C etc. when used to delimit an area which is part of a larger named area, but the words are given in full when used in other contexts. Mountain(s) is abbreviated to Mt(s); the words river and lake are given in full.

SOURCES OF DATA

This work is based largely on observations made by the author and his wife during continuous fieldwork between 1976 and 1980 (Nikolaus 1979a, b; 1982 a, b), during which distributional data were routinely plotted on a 1° square grid, and intensive netting and ringing produced information on rarer species, on migratory status (Nikolaus & Backhurst 1982) and on the distribution of different races (Nikolaus in prep.). To assist in racial determination about 1000 specimens were collected, and these are now housed in the Naturkunde Museum Stuttgart and the Alexander Koenig Museum Bonn (v.d.Elzen and Konig 1983). Also included here are recent data kindly supplied by the Chicago Field Museum (Imatong/Bengengai, February/March 1976; Traylor & Archer 1982 and *in litt.*); the Copenhagen Museum (Imatong, November/December 1980; F.P. Jensen *in litt.*); and recent field observations from P.J. Dare (November/December 1960), J.S. Fairclough (Mbili/Wau 1981, 1982); J.C. & S Hillman (Bengengai 1980, 1981); F.R. Lambert (Nile and Imatong March/April 1983; and H. Lege (November 1981/January 1982). Most of the 'old' literature reviewed by Cave & MacDonald has been included, as also have data from the collection at the Khartoum Natural History Museum (Bowen 1926, 1931) and from the British Museum (Nat. Hist.) at Tring, from the African Atlases of Hall & Moreau (1970) and Snow (1978), and from field notes of F.O. Cave (Wau 1955; files in the British Museum).

In addition to the author's own records, breeding data are from Mackworth-Praed & Grant (1952, 1955), which included most of the 'old' English literature, Koenig (1911, 1926), M.A. Traylor (*in litt.*) and F.O. Cave (files in the British Museum).

Considerable efforts have been made to trace original old sources, although only major publications are included in the references. Because of the space available for the species accounts, references are only given to support records or statements of special interest, but a more detailed bibliography is available from the author.

ORDER AND NOMENCLATURE

The order and nomenclature followed is generally that used by Britton (1980). Thus, nomenclature is essentially based on White (1960, 1961, 1962a,b, 1963, 1965), though Vaurie (1959,1965) is usually followed for Palaearctic species. Order of families is that of Wetmore (1960) as modified by Benson et al. (1970), and of sub-families within families that of Thompson (1964). Genera and species are listed alphabetically.

FORMAT OF SPECIES IN THE LIST

Each species is listed under a family heading (and Sub-family heading where appropriate) with information given in the following sequence :

- a. Sequential number on the Sudan list (this number is used in the index, not the page number). Comments are included on species not yet recorded, but expected to occur in the Sudan, and on species formerly listed but no longer considered acceptable; these species are included in the systematic list but are not numbered.
- b. Scientific name for the species.
- c. English name (generally those used in Britton 1980).
- d. For ease of cross-reference, the species number in Mackworth-Praed & Grant (1952, 1955) is shown on the right hand side.
- e. The suggested status is given (PMW, PMP, AM, LM, R, V).
- f. Breeding status is given for all Afrotropical species (BR, NBR), together with months of breeding.
- g. A concise resume of known distribution, status, abundance and times of occurrence is given, with information on each sub-species occurring in the region.

ACKNOWLEDGEMENTS

Even though few people were able to supply new field data, there have been a great number helping in many ways. My wife Minna assisted me in the field in the South Sudan for four years. Hans Rupp has to be specially mentioned because we completed many field trips together until his unfortunate death in 1979 during the course of his work on

the small mammals of South Sudan. For valuable field data and other comments I would like to thank A.L. Archer, Mrs Barker, J.S. Fairclough, C. & S. Hillman, F.R. Lambert, H. Lege, F.P. Jensen and M.A. Traylor, and especially P.L. Britton and P.J. Dare who gave valuable comments on an earlier draft of this manuscript. For assistance in the identifications of skins and notes on literature I am grateful to the late C.W. Benson, P.R. Colston, G.R. Cunningham-van Someren, Dr. R. v.d.Elzen, Dr. C.H. Fry, J. Fryxell, I.C.J. Galbraith, A. Guillet, P. Hogg, Dr. C. König, Dr. M. Louette, Dr. J. Nikolai, R. Schlenker and Dr. H. Wolters; also to Dr. E. Schüz who helped in various other ways as well. Very essential was the assistance given by various individuals in Khartoum and Juba, especially Mr. Abu Sineina, Dr. El Rayah, H. Minga and C. Acire. John Blower recommended me to the Wildlife Department, whose support proved basic to the whole work. Last but not least D.J. Pearson and G.C. Backhurst commented on the manuscript and finally edited it. Many thanks to them and to any others not mentioned who helped in one way or another.

Gerhard Nikolaus, Bosenbüttel 4, 2859 Spieka, F.R. Germany.

SPECIES RECORDED

Family STRUTHIONIDAE Ostrich

1. *Struthio camelus* Ostrich 1
R LM BR (1, 3, 10)
The race *camelus* is uncommon in the tree savanna and open grassland north of a line from Juba to Wau, decreasing in numbers though locally still common. A 1980 game count in Boma Park gave 2150 for the dry season and 1300 for the wet season.

Family PODICIPEDIDAE Grebes

2. *Tachybaptis ruficollis* Little Grebe 4
R? LM AM? NBR
The race *capensis* is uncommon on large waters. Probably breeding on the White Nile and Lake No. All records are from October - April. Some are possibly resident. Birds in April in Aweil were on passage.

Family PELECANIDAE Pelicans

3. *Pelecanus onocrotalus* White Pelican 31
AM PMW? NBR
The few records are all in January, from the Nile, Sobat and Lol Rivers. The origin of these birds is unknown.
4. *Pelecanus rufescens* Pink-backed Pelican
R LM BR (1, 2)
Fairly common throughout on all suitable waters; more common in winter.

Family PHALACROCARIDAE Cormorants

5. *Phalacrocorax africanus* Long-tailed Cormorant 27
R? LM NBR
The nominate race is common in the dry season on any waters but moves north with the rains. Hundreds roost at Lake No.
- Phalacrocorax carbo* Cormorant 25
The only record, from Wau, needs confirmation. Probably a rare visitor to large rivers.

Family ANHINGIDAE Darters

6. *Anhinga rufa* Darter 28
R LM BR (8, 9)
The nominate race is a common resident on rivers and larger waters throughout.

Family ARDEIDAE Herons, Bitterns, Egrets

Most species vacate the large Sudd area in the rainy season, probably moving to the northern savanna or Eurasia for breeding. The year-round occurrence of several species around Aweil suggests that some breed locally.

Sub-family Botaurinae

Botaurus stellaris Bittern

52

The Palaearctic nominate race probably reaches South Sudan. It is known as a rare winter visitor to the North, while there is a single record from Uelle in NE Zaire.

7. *Ixobrychus minutus* Little Bittern

50

PMP R? LM

NBR

Uncommon in swamps and reedbeds. The exact status of the two races is not known. There are specimens of *payesii* from Maridi and Tambura, while several examples of the Palaearctic nominate race were caught in October at the Baro River in neighbouring Ethiopia.

8. *Ixobrychus sturmii* Dwarf Bittern

51

R? LM?

NBR

Uncommon throughout. There are dated records from Aweil (September), Wau (March), Boma (February) and Torit (November). No information about seasonal movements.

Sub-family Ardeinae

9. *Ardea cinerea* Grey Heron

33

PMW R?

NBR

The nominate race is a fairly common Palaearctic visitor from October to March to marshy places. A few oversummer, but breeding has not yet been reported.

10. *Ardea goliath* Goliath Heron

35

R? LM

BR (11, 12)

Common in the dry season on larger rivers with deep water, where it breeds, moving north during the rains.

11. *Ardea melanocephala* Black-headed Heron

34

R LM

NBR

Common throughout in a variety of open habitats, especially as a dry season visitor. It breeds in the North Sudan from June to August.

12. *Ardea purpurea* Purple Heron

36

PMW R?

BR (5)

The nominate race arrives in September as a locally common Palaearctic visitor along the Nile in the Sudd. Few are seen in summer. Once reported breeding near Shambe.

13. *Ardeola ralloides* Squacco Heron

41

PMW LM?

NBR

The nominate race is a common Palaearctic visitor to marshes and swamps, mainly from September to February; less common in March-April. A few probably oversummer.

14. *Ardeola ibis* Cattle Egret

41

R? AM

NBR

The nominate race is a very common bird throughout, often accompanying cattle or game. Most birds leave in summer, presumably for the North Sudan for breeding. It is likely that they also breed in the northern part of South Sudan.

15. *Butorides striatus* Green-backed Heron 45
R LM? NBR
The race *atricapillus* is an uncommon but widespread resident of rivers and swamps with dense fringing vegetation. Seasonal movements need confirmation.
16. *Egretta alba* Great White Egret 37
PMW R? LM NBR
The race *melanorhynchos* is fairly common, mainly associated with larger rivers and lakes. Nearly all dated records are for the period September to April. They occur throughout the year.
17. *Egretta ardesiaca* Black Heron 39
R? NBR
There are only five records, from Aweng, north of the River Lol, Aweil, Sobat and Lado. It frequents creeks and rivers.
18. *Egretta garzetta* Little Egret 40
PMW LM NBR
The nominate race is common in wetland habitats. All dated records are for the period August to April.
19. *Egretta intermedia* Yellow-billed Egret 38
R? LM NBR
The race *brachyrhyncha* is a non-breeding visitor to swamps and marshes. Mainly absent from May to August. It is less common east of the Nile.
20. *Gorsachius leuconotus* White-backed Night Heron 49
R? AM? NBR
Rare, and possibly only a non-breeding visitor. Once recorded from Maridi in February. This species of thick bush vegetation with trees along rivers is common in NE Zaire, and there are records from the Baro River in Ethiopia.
21. *Nycticorax nycticorax* Night Heron 48
PMW R LM BR (5, 6)
The nominate race is fairly common throughout around wetlands. Breeding is reported from Aweil and Shambe. Concentrations in very large roosts at Lake No and the Sobat River in January/February are most probably Palaearctic visitors.

Family BALAENICIPITIDAE Shoebill

22. *Balaeniceps rex* Shoebill 54
R BR (1, 9 - 12)
A resident of the Sudd, much endangered by destruction of papyrus and heavy cattle grazing. Recent counts gave about 6000 birds with the biggest population in the River Lol area and the Bahr-el-Zeral Game Reserve.

Family SCOPIDAE Hamerkop

23. *Scopus umbretta* Hamerkop 53
R LM NBR
The nominate race is a common resident throughout wherever there is water, preferably stagnant pools. Recorded up to 3000 m in the Imatong Mountains.

Family CICONIIDAE Storks

Sub-family Ciconiinae

24. *Anastomus lamelligerus* Open-billed Stork 59
AM R? BR (6, 7)
The nominate race is a very common dry season visitor throughout, avoiding only large tracts of dense forest. Exceptional numbers are seen after bush-fires in the Sudd. Birds in the Nimule region August to October probably originate from nearby breeding colonies. Breeding possibly occurs around Aweil.
25. *Ciconia abdimii* Abdim's Stork 58
AM LM? BR (4,5)
A breeding visitor in the rains. The main breeding area is in North Sudan, but a few pairs breed in Juba, Bor, Rumbek and the northern parts of the area. Large numbers are seen on passage in April, fewer on southward passage in September and October. Some birds frequent open grasslands in the Sudd and the southeast through the dry season.
26. *Ciconia ciconia* White Stork 55
PMW PMP
The nominate race is an uncommon Palaearctic visitor to swamps and grasslands, mainly from November to February. There is a marked southward passage in October and November.
27. *Ciconia episcopus* Woolly-necked Stork 57
LM AM BR (1, 2, 11, 12)
The race *microscelis* is a breeding visitor to open country in South Sudan in the dry season. Numbers are always small, sometimes frequenting burnt grassland. There are no records from forest areas along the Zaire border.
28. *Ciconia nigra* Black Stork 56
PMW
This Palaearctic migrant is an uncommon winter visitor. All records are of single birds, from Tonj (January), Juba (March), the Imatong Mountains (December, April), Nimule (January) and Bengengai.
29. *Ephippiorhynchus senegalensis* Saddle-billed Stork 60
R BR (1, 9 - 12)
Fairly common throughout, nearly always in pairs, at larger swamps with permanent water. So far, breeding is only confirmed from the Upper Nile region, but recently fledged birds seen in February and April suggest more widespread breeding.
30. *Leptoptilos crumeniferus* Marabou 61
R LM BR (1, 10 - 12)
A widespread and common breeding visitor. Some of the twenty or more documented breeding colonies are small; but the largest, at Thiet, has at least 100 pairs. Concentrations are a feature of bush-fires and drying waterpools late in the dry season (February to March). Few remain during the rains.

Sub-family Mycteriniinae

31. *Mycteria ibis* Yellow-billed Stork 62
 R LM BR (10 - 12)
 Uncommon, frequenting marshes and rivers with permanent water.
 Two breeding colonies are known at Ayod and Jebel Kasangor.
 Breeding is also suspected at Aweil. Most or all vacate South
 Sudan during the rains (May - September) and migrate north.

Family THRESKIORNITHIDAE Ibises, Spoonbills

Sub-family Threskiornithinae

32. *Bostrychia hagedash* Hadada 65
 R. NBR
 The race *brevirostris* is a common resident throughout. It favours
 wooded localities near swamps and rivers.
33. *Plegadis falcinellus* Glossy Ibis 68
 PMW PMP? AM? NBR
 The nominate race is a rather uncommon winter visitor in small
 numbers to large swamps, recorded from Malakal, Lake No, Aweil,
 Kongor and Juba. Up to 500 arrive in Aweil and Kosti in August.
 Numbers decrease markedly in early September. These, and regular
 November/December flocks of 50 - 100 in the Imatong Mountains are
 probably Palaearctic breeders on southward passage.
34. *Threskiornis aethiopica* Sacred Ibis 63
 LM NBR
 The nominate race is a very common non-breeding visitor in the
 dry season, likely to occur at any site. It breeds in North Sudan.
 There are no records during the period May - July.

Sub-family Plataleinae

35. *Platalea alba* African Spoonbill 70
 LM AM? NBR
 An uncommon wanderer to swamps and rivers with permanent water.
 There might be a breeding colony at the Juron River at Jebel
 Kasangor, where this species was noticed in large numbers in
 February.
36. *Platalea leucorodia* Spoonbill 69
 PMW
 The nominate race is a rare Palaearctic visitor, known only from two
 records: Aweil in March (2 birds), Juba in January (2 birds).

Family PHOENICOPTERIDAE Flamingos

- Phoenicopus minor* Lesser Flamingo 72
 This species might occur in the extreme southeast near Lake
 Turkana. There is no definite record.
37. *Phoenicopus ruber* Greater Flamingo 71
 PMW AM? NBR
 The race *roseus*, regular in numbers on the White Nile at Khartoum,
 is only an occasional visitor south to Malakal and Kaka. Khartoum
 records include a bird ringed as a nestling in Iran, but African
 breeders might occur, especially near Lake Turkana in the southeast.

Family ANATIDAE Ducks, Geese

Ducks and Geese are generally rather uncommon in South Sudan. The Aweil/River Lol area is the only region where large numbers are regular. The main wintering range of most Palaearctic species is north of a line from Aweil to Malakal and the Sobat River.

Sub-family Anserinae

38. *Dendrocygna bicolor* Fulvous Whistling Duck 95
 R? LM NBR
 Generally less common than the next species with which it often associates, favouring the same habitat. A record of 50 - 200 at Aweil during late August and early September is noteworthy.
39. *Dendrocygna viduata* White-faced Whistling Duck 94
 R LM BR (6)
 Common resident throughout, favouring sandy rivers in wooded grassland. Breeding is reported from the Aweil/River Lol area.

Sub-family Anatinae

40. *Alopochen aegyptiaca* Egyptian Goose 99
 R? LM NBR
 An uncommon but widely distributed non-breeding visitor in the dry season. It probably breeds around Aweil.
41. *Anas acuta* Pintail 92
 PMW
 The second most numerous Palaearctic duck wintering in the northern part of South Sudan; regularly seen at Malakal, and the Sobat River. There are no records from further south.
42. *Anas clypeata* Shoveler 80
 PMW
 An uncommon Palaearctic winter visitor occurring in small numbers at wetlands throughout, mostly on larger lakes.
43. *Anas crecca* Teal 87
 PMW
 The nominate race is an uncommon Palaearctic visitor to smaller ponds from January to March. It ranges south to Tonj and Bor, always in small numbers.
44. *Anas erythrorhynchus* Red-billed Duck 91
 AM? V
 A bird mentioned by Cave & Macdonald from the Kengen River near Pibor is the only record. Its status is uncertain.
45. *Anas hottentota* Hottentot Teal 90
 AM NBR
 Collected once at Gogrial in 1956 (Nikolaus 1984).
46. *Anas penelope* Wigeon 86
 PMW
 This Palaearctic visitor is common in North Sudan, but winters in the South Sudan only on larger open waters. Cave & Macdonald included Aweil and the Kengen River in its wintering range.

17. *Anas querquedula* Garganey 88
PMW PMP
The most common Palaearctic duck. Numerous flocks of 50 - 100 are regular around Aweil - Malakal. It occurs in smaller numbers on larger waters throughout, both in winter and on passage. A bird ringed at Juba (November) was shot in USSR (56°00'N, 71°30'E) (October).
18. *Anas sparsa* Black Duck 84
R BR (1, 2)
The race *leucostigma* is resident on wooded mountain streams in the Imatong Mountains, and along the Kajiko River south of Yei.
19. *Aythya fuligula* Tufted Duck 78
PMW
Cave & Macdonald recorded this Palaearctic species south to Malakal. There are no recent records.
20. *Aythya nyroca* White-eyed Pochard 75
PMW
A rare Palaearctic visitor known only from a single flock at Aweil in March (Cave & Macdonald 1955).
21. *Netta erythrophthalma* Southern Pochard 77
AM NBR
The race *brunnea* is possibly an overlooked non-breeding visitor from southern Africa, recently recorded from the Upper Nile swamps, frequenting open water.
22. *Nettapus auritus* African Pygmy Goose 97
R? LM AM? NBR
Locally common in the rainy season at deep pools with reeds and floating vegetation, including up to 30 at Aweil in August - September. There are records of single birds from Aweil, Meshra-er-Req, Pibor, Lake Nibor, Maridi and Yei.
23. *Plectropterus gambensis* Spur-winged Goose 100
R? LM AM? BR (9)
The nominate race is fairly common on swampy grasslands throughout. Most birds move north in the rainy season, from May. In August - September large concentrations with flocks of 50 - 200 at Aweil are noteworthy.
24. *Pteronetta hartlaubii* Hartlaub's Duck 93
R BR (7, 8)
Resident along forest streams in Zandeland, near the Zaire border, where it was reported to be locally common (Cave & Macdonald 1955). Present status unknown.
25. *Sarkidiornis melanotos* Knob-billed Goose 98
AM LM R? NBR
Locally common, favouring larger swamps. In northern areas it is most frequent and numerous from mid-June to late October, possibly as a breeding visitor. Daily counts of 100 - 300 at Aweil in August - September are noteworthy. Three ringed birds from Zimbabwe have been recovered in southern Darfur.

56. *Thalassornis leuconotus* White-backed Duck 74
 R? LM? NBR
 There is an unlabelled specimen in the Khartoum Museum, probably from Gogrial in 1956. Otherwise, there is only one recent sight record, from the Upper Nile region.

The following ducks might occur, but have not been recorded so far :

<i>Anas capensis</i>	Cape Wigeon	89
<i>Anas strepera</i>	Gadwall	85
<i>Anas undulata</i>	Yellow-billed Duck	83
<i>Aythya ferina</i>	European Pochard	76

Family SAGITARIDAE Secretary Bird

57. *Sagittarius serpentarius* Secretary Bird 103
 R? LM NBR
 Confined to the extreme southeast, east of Kapoeta, frequenting *Acacia* savanna country. It is likely that breeders from North Sudan enter the grasslands of the Sudd during the dry season.

Family ACCIPITRIDAE Birds of prey (Vultures, Eagles, Hawks etc.)

Sub-family Aegyptiinae

58. *Gypohierax angolensis* Palm-nut Vulture 161
 R? LM NBR
 Not uncommon in the forest region along the Zaire border. Records are for July - February, suggesting that there are seasonal movements, perhaps necessitated by a lack of ripe palm nuts. The record from Didinga in Snow (1978) is unsubstantiated.
59. *Gyps africanus* White-backed Vulture 107
 R LM BR (11, 12)
 Resident throughout, more common in the south. Breeds commonly in tall trees in the open savanna between Kapoeta and Boma. In the rainy season records are restricted to larger towns like Juba, Wau and Torit, where regular food is available at slaughter houses. This species is more numerous than the next in most areas, though the position is reversed north of Malakal.
60. *Gyps rueppellii* Rüppell's Vulture 106
 R LM BR (11, 12)
 Mainly restricted to the area east of Maridi - Malakal, but also occurs in the extreme west at Boro and Kafia Kingi. Large breeding colonies are known on precipitous rocks at Amadi and Boma. Most records are for October - April; May - September records are all from around nesting sites.
61. *Neophron monachus* Hooded Vulture 111
 R LM? NBR
 A common resident throughout, except east of Kapoeta where it is replaced by the Egyptian Vulture. Numbers are much reduced during the rains.

52. *Neophron percnopterus* Egyptian Vulture 110
R? PMW LM NBR
Uncommon north of Malakal and in the Ilemi Triangle, replacing the hooded Vulture in the more arid regions.
53. *Torgos tracheliotus* Lappet-faced Vulture 108
R LM BR (1, 11, 12)
Restricted to the *Acacia* grasslands of the southeast, ranging west to the Nile in the dry seasons. Common near Boma where it breeds on tall trees on open plains. Birds from North Sudan wander to northern areas in the dry season.
54. *Trigonoceps occipitalis* White-headed Vulture 109
R LM BR (11, 12)
Scarce in the forest region along the Zaire border, but otherwise fairly common throughout. It breeds in large trees on the Boma Plateau. Numbers are much reduced during the rainy season.
- Sub-family Gypaetinae
- Gypaetus barbatus* Lammergeyer 162
No records. The race *meridionalis* is likely to occur in the Imatong and Didinga Mountains along the border with Uganda.
- Sub-family Circinae
55. *Circus aeruginosus* Eurasian Marsh Harrier 184
PMW PMP
The Palaearctic nominate race is regular around Aweil and the Sudd on passage in late August and September. Numbers winter in December - February but there are few records for October - November and March. Always associated with swampy places.
56. *Circus macrourus* Pallid Harrier 183
PMW
A Palaearctic visitor to open grassland throughout from December to March. Common in the drier parts of the southeast, but rather uncommon elsewhere. No obvious evidence of passage.
57. *Circus pygargus* Montagu's Harrier 182
PMW
A Palaearctic visitor to open grassland throughout from December to March, typically in wetter places than those favoured by the Pallid Harrier.
58. *Circus ranivorus* African Marsh Harrier 184
V? R? NBR
Status not clear. An immature bird caught and ringed near Juba in October supports recent sight records in the Sudd (Nikolaus 1979).
- Sub-family Polyboroidinae
59. *Polyboroides radiatus* Harrier Hawk 185
R BR (1, 2)
The race *typus* is an uncommon resident of better wooded savanna. It also occurs in the large tracts of forest along the Zaire border, and in the forested parts of the Imatong Mountains. There are no records from open grasslands of the Sudd and the Ilemi Triangle.

Sub-family Circaetinae

70. *Circaetus cinerascans* Banded Snake Eagle 157
 R BR (1 - 3)
 A common resident of forests in the Yei, Imatong, Didinga and Boma areas. There is no record from the Zande forests along the Zaire border.
71. *Circaetus cinereus* Brown Snake Eagle 153
 R LM BR (1)
 An uncommon bird of more heavily wooded savanna. There are records from Wau, Amadi, Kajo Kaji, Torit and Didinga
72. *Circaetus gallicus* Short-toed Snake Eagle 152, 154, 155
 (Including Beaudouin's Snake Eagle *C.g. beaudouini* and Black-chested Snake Eagle *C.g. pectoralis*)
 PMW R? LM BR (1)
 The nominate Palaearctic race is an uncommon visitor throughout, from October to March, recorded from Boma, Aweil, Juba, Torit, Kapoeta and the Nile north of Malakal. The few records of the race *beaudouini* are from Aweil, the Sudd and the Lol River in January and March; status uncertain, but probably an uncommon seasonal visitor to wetter grasslands. The race *pectoralis* is uncommon in the area southeast of Juba, mostly frequenting *Acacia* grassland; all records are for the period October - May.
73. *Terathopius ecaudatus* Bateleur 159
 R LM NBR
 A widespread and common resident, frequenting all kinds of habitats but preferring open wooded savanna. It breeds at the end of the rainy season.

Sub-family Accipitrinae

74. *Accipiter badius* Shikra 176
 R LM NBR
 The race *sphenurus* is a fairly common resident in savanna and *Acacia* woodland throughout. Local northward movements during the rains are reported.
75. *Accipiter brevipes* Levant Sparrowhawk 175
 PMW
 The only record of this Palaearctic migrant is a specimen from Boma in December, but recent observations in North Sudan suggest wintering east of the Nile (Nikolaus 1984).
76. *Accipiter melanoleucus* Great Sparrowhawk 174
 R NBR
 The nominate race is an uncommon resident of heavily wooded savanna. There are recent records from Li Rangu, Zande, Gilo, the Upper Talanga and the Imatong Mountains.
77. *Accipiter minullus* Little Sparrowhawk 170
 R? NBR
 The nominate race is a widespread but uncommon resident. Most records are from the *Acacia* woodlands in the southeast and north of Malakal. West of the Nile recorded only from Juba and Amadi.

78. *Accipiter rufiventris* Rufous Sparrowhawk 173
R? V? NBR
The nominate race was observed once in April 1980 in the Imatong Mountains at Gilo (Nikolaus 1982). A record for the Kapoeta/Boma Hills region given by Snow (1978) is an error.
79. *Accipiter tachiro* African Goshawk 177
R? NBR
Uncommon in forested regions. The race *unduliventer* occurs at Boma Hills, while birds from the Imatong Mountains are assigned to *sparsimfasciatus*.
80. *Accipiter tousenelii* Red-chested Goshawk W 162
R? NBR
The race *canescens* is probably a rare resident of the forests along the Zaire border. It has been collected by Traylor (1977) (Traylor & Archer 1982).
81. *Aquila nipalensis* Steppe Eagle 138
PMW PMP
The race *orientalis* winters regularly in small numbers from late December to February. Passage birds are noted east of the Nile from late September to December and in April.
82. *Aquila pomarina* Lesser Spotted Eagle
PMP
The nominate Palaearctic race has been recorded only east of the Nile. Precise status is uncertain however, because of former confusion with Tawny Eagle and Steppe Eagle.
83. *Aquila rapax* Tawny Eagle 139
LM? AM NBR
The race *belisarius* is probably only a non-breeding visitor from North Sudan. It is fairly common throughout all savanna regions from October to May.
84. *Aquila verreauxi* Verreaux's Eagle 136
R NBR
This species which favours rocky escarpments, has been reported from Imatong at Mt. Kinyeti and Upper Talanga, and from the Uganda side of the Didinga Mountains.
85. *Aquila wahlbergi* Wahlberg's Eagle 142
R? AM? NBR
A non-breeding visitor from December to April, it was seen in large numbers together with Black Kites and Tawny Eagles in late March - early April after heavy rains in the Amadi/Yei region. It might breed in the Torit/Kapoeta area where it is recorded throughout the year.
86. *Butastur rufipennis* Grasshopper Buzzard 158
R AM BR (3, 5)
A locally very common breeding visitor in the dry season in any habitat except rain forest. Birds leave for the North Sudan from the end of May and return in October. Large numbers congregate about bush fires.

87. *Buteo augur* Augur Buzzard 167
 R NBR
 The nominate race is resident at Gilo/Imatong Mountains; also recorded from Boma, Didinga and Kajo Kaji. A bird of the high-lands, favouring forest edges and cultivation.
88. *Buteo auguralis* Red-necked Buzzard 168
 R? LM BR (2)
 Not uncommon in better wooded savanna south of Wau - Torit, and along the Ethiopian border. Some birds might move north during the rains.
89. *Buteo buteo* Common Buzzard 163
 PMW PMP
 The race *vulpinus* is a locally common Palaearctic visitor and passage migrant mainly east of the Nile. Passage birds are seen regularly in Imatong in October and March - early April.
90. *Buteo rufinus* Long-legged Buzzard 166
 PMW
 The Palaearctic nominate race is an uncommon winter visitor from October to March, favouring dry hills. Rare in the Sudd and further south.
91. *Buteo tachardus* Mountain Buzzard 165
 R? LM NBR
 The race *oreophilus* is fairly common in the Imatong, Dongotona and Didinga Mountains. There are no records between mid-May and the beginning of August.
92. *Hieraaetus pennatus* Booted Eagle 145
 PMW PMP
 The nominate race is an uncommon Palaearctic winter visitor. Records are from Renk, Maridi, Juba, Torit, Kapoeta and the Kenya/Sudan border from November to March. Passage has been noticed in November and March.
93. *Hieraaetus spilogaster* African Hawk Eagle 143
 R? LM NBR
 An uncommon dry season visitor favouring well-wooded hilly country. Recorded from Boro, Wau, Juba, Torit and Boma.
94. *Kaupifalco monogrammicus* Lizard Buzzard 150
 R LM BR (3)
 An uncommon resident in better wooded savanna and forest edges in the southern part, rare in the west, north to Wau.
95. *Lophaetus occipitalis* Long-crested Hawk Eagle 149
 R LM BR (4)
 Common throughout. A bird of wooded savanna and cultivation. Absent from the grasslands of the Sudd. Some move north during the rains.
96. *Melierax gabar* Gabar Goshawk 178
 R NBR
 A fairly common resident throughout in any kind of woodland except in the forests of Imatong and Zandeland. Less common than Dark Chanting Goshawk.

97. *Melierax metabates* Dark Chanting Goshawk 180
 R LM? BR (4, 5)
 The nominate race is a common resident throughout in any kind of woodland except the green forests in Zandeland and Imatong.
98. *Polemaetus bellicosus* Martial Eagle 146
 R LM NBR
 An uncommon resident of the savanna along the western, southern and eastern borders.
99. *Stephanoaetus coronatus* Crowned Eagle 147
 R BR (1)
 A rare resident of the forest region, which has been reported breeding on the south side of the Imatong Mountains. Occasionally seen at Gilo. An old record from Bahr-el-Ghazal (Butler 1908) is unlikely.
100. *Urotriorchis macrourus* Long-tailed Hawk 181
 R? LM NBR
 First recorded in January 1982 at Bengengai (J.C. Hillman *in litt.*)
 A rare forest species; precise status unclear.

Sub-family Milvinae

101. *Haliaeetus vocifer* Fish Eagle 160
 R BR (1, 10 - 12)
 A common resident throughout at any permanent water.
102. *Milvus migrans* Black Kite 132
 PMW PMP R LM AM? BR (1, 2, 12)
 The Palaearctic race *migrans* is a winter visitor and passage migrant throughout from October to April. The Yellow-billed Kite *M.m. parasitus* is a breeding visitor and resident. After breeding, most birds depart in April and May and return in August. Enormous numbers concentrate around bush fires and drying swamps. The race *aegyptius* probably also occurs. Distribution and relative abundance of the different races are not well documented.

Sub-family Perninae

103. *Pernis apivorus* Honey Buzzard 135
 PMW? PMP?
 Status uncertain but apparently rare; only one definite record, from Bengengai in November (J.C. Hillman *in litt.*).
104. *Chelictinia riocourii* Swallow-tailed Kite 131
 LM AM? BR (4)
 Only recorded from the Ilemi Triangle, where it arrives, presumably from East Africa, in February for breeding. A bird was seen flying south over Gilo in November. Birds from North Sudan occasionally come south to Malakal in the dry season.
105. *Elanus caeruleus* Black-shouldered Kite 133
 R? LM AM? NBR
 The nominate race is common throughout open savannas in the dry season from November to April. Around Juba, birds are recorded also from August to October.

106. *Macheiramphus alcinus* Bat Hawk 134
 R LM? NBR
 Recorded from Juba (January), Lol River (March) and Aweil (August).
 Always associated with a large supply of small bats.
- Family PANDIONIDAE Osprey
107. *Pandion haliaetus* Osprey 186
 PMW
 The nominate race is a rare visitor to permanent lakes and the Nile. Records are from Malakal (February), Lake Nibor and Nimule (January), and Maridi (August).
- Family FALCONIDAE Falcons
108. *Falco alopec* Fox Kestrel 125
 R LM NBR
 A locally common resident on mountains and cliffs, most common in the southeast. Local movements are indicated by its appearance in the non-breeding season far from rocky mountains.
109. *Falco ardosiaceus* Grey Kestrel 127
 R? LM NBR
 Frequents open savanna and possibly breeds throughout; locally common during the dry season; undertakes short range movements which are not well understood.
110. *Falco biarmicus* Lanner Falcon 113
 R LM BR (1, 2)
 Now an uncommon resident, its numbers having decreased in the last thirty years. The race *abyssinicus* is reported in the southeast from Bor to Nimule and Lake Turkana. Birds at Malakal, Kaka and Aweil (north of 8°N) belong to the race *tanypterus*.
111. *Falco cherrug* Saker Falcon 121
 PMW
 The nominate race is a rare Palaearctic visitor in the east, recorded south to Kajo Kaji, but not recently (Cave & Macdonald 1955).
112. *Falco chicquera* Red-necked Falcon 122
 R? LM BR (1 - 4)
 The race *ruficollis* is a widespread breeding visitor which is locally fairly common; it nests in *Borassus* and sometimes *Hyphaene* palms. Records are all from January to April and July/August; movements require clarification.
113. *Falco cuvieri* African Hobby 116
 R LM? NBR
 A rare resident of hilly savanna country in the southern part; reported from Boma, Nataporoputh Hills, Lado, Yei, Mt. Baginzi and Tonj.
114. *Falco fasciinucha* Taita Falcon 114
 R? LM? NBR
 Rare; it has been reported from high lava cliffs northwest of Lokitaung on the Kenya/Sudan border (Dowsett 1977).

Falco naumanni Lesser Kestrel

Status uncertain; no definite record for South Sudan but perhaps overlooked due to confusion with Common Kestrel. In North Sudan it is common on passage and winters locally, the two species often being seen together around grass fires.

115. *Falco pelegrinoides* Barbary Falcon 112
PMW

Probably a rare visitor to the arid extreme southeast, recorded from Kapoeta in February 1969 (Nikolaus pers. obs.).

116. *Falco peregrinus* Peregrine Falcon 112
PMW? R? LM NBR

Reported in open savanna north of Malakal in February and around Juba in January, February and October. The status of the various races needs to be clarified; *minor* is probably a rare resident and wanderer, recently identified at Aweil in August; the Palaearctic *calidus* may occur.

117. *Falco subbuteo* Hobby 115
PMP

The nominate Palaearctic race is a rare passage migrant, reported from Juba in November and Nimule in April.

118. *Falco tinnunculus* Kestrel 123
PMW PMP? R NBR

Nearly all records refer to Palaearctic migrants of the nominate race, wintering between December and February, and rarely in November and March. Common in the dry grass savanna north of Malakal and in the southeast, but uncommon in the grasslands of the Sudd and River Lol. The resident race *rufescens* has been reported from Boma.

- Falco vespertinus* Red-footed Falcon 119

There are no definite records, but it might be expected on passage in western areas since Lynes reported pronounced autumn passage at Darfur.

119. *Polihierax semitorquatus* Pygmy Falcon 129
R LM NBR

Fairly common resident in the dry thorn-bush country of the Ilemi Triangle (east of Kapoeta) and west to Juba as an uncommon wanderer in the dry season. It is known to take over nests of White-headed Buffalo Weaver for breeding and nesting.

The following species might occur :

<i>Accipiter ovampensis</i>	Ovampo Sparrowhawk	172
<i>Aquila clanga</i>	Greater Spotted Eagle	140
<i>Hieraaetus dubius</i>	Ayres' Hawk Eagle	144
<i>Melierax poliopterus</i>	Pale Chanting Goshawk	179
<i>Aviceda cuculoides</i>	Cuckoo Hawk	130
<i>Falco concolor</i>	Sooty Falcon	117
<i>Falco eleonora</i>	Eleonora's Falcon	118
<i>Falco rupicoloides</i>	White-eyed Kestrel	124

Family PHASIANIDAE Quails, Francolins

120. *Coturnix chinensis* Blue Quail 213
R? LM? AM? BR (9)
A rare bird of wetter short grassland. Recorded only twice from Juba and from Yei, where it was breeding at the end of the rains. (Cave & Macdonald 1955; Nikolaus pers. obs.).
121. *Coturnix coturnix* Quail 211
R? AM? PMW? NBR
Uncommon in short grassland; recorded from Aweil (March, September), Lado (January) and Juba (September, October, November). It is not clear to what extent these birds belong to the nominate Palaearctic race or, especially in the south, to the African race *erlangeri*.
122. *Coturnix delegorguei* Harlequin Quail 212
R? LM AM BR (8, 9, 12)
The nominate race is in some years a very common visitor to *Acacia* short grassland from May to December east of Torit. Occasional wanderers are reported from Aweil, Tonj, Kongor and Malakal. Numbers and movements are irregular.
123. *Francolinus clappertoni* Clapperton's Francolin 202
R BR (9, 11)
A fairly common resident of open savanna. Three races occur: *heuglini* northwards from Bor and Tambura, *gedgi* in the south from the Nile at Juba to Didinga, and *cavei* on the Boma Plateau.
124. *Francolinus icterorhynchus* Heuglin's Francolin 201
R BR (10 - 12)
The nominate race is a common resident of wooded savanna west of the Nile from Kajo Kaji to Tonj and Kafia Kingi.
125. *Francolinus lathami* Forest Francolin 189
R NBR
The race *schubotzi* is a rare resident of the gallery forest in Zandeland.
126. *Francolinus leucoscepus* Yellow-necked Spurfowl 210
R NBR
The race *infuscatus* is a fairly common resident in the extreme southeast from Kapoeta to Boma and Lake Turkana, frequenting dry grass plains.
127. *Francolinus levillantoides* Smith's (Archer's) Greywing 196
R NBR
The race *archeri* is a rare resident of the extreme southeast and is only known from specimens from Natoporoputh Hills and Nagishot, Didinga Mountains. A species frequenting dry rocky slopes with sparse grass.
128. *Francolinus schlegelii* Banded Francolin 193
R NBR
Uncommon resident in the western Bahr-el-Ghazal west of Wau and Gogrial, favouring thickly wooded savanna with clearings. Invariably found in association with "Ka" trees *Isoberlinia doka*. Probably breeding September - October.

129. *Francolinus sephaena* Crested Francolin 195
 R BR (6, 12)
 The race *grantii* is a common resident in the southeast from Juba to Boma and Didinga, favouring bushy savanna. Recent records north to Wau and Malakal may have been young Clapperton's Francolin, but needs confirmation
130. *Francolinus squamatus* Scaly Francolin 204
 R BR (3, 11)
 The race *schuetti* is locally common in the extreme south from Didinga to Zande, favouring secondary growth and cultivation adjoining mountain and gallery forests up to 2400 m.
131. *Ptilopachus petrosus* Stone Partridge 214
 R BR (10)
 Common resident south west of Kapoeta - Aweil. Two races occur: *florentiae* in the hilly country of Equatoria; *brehmi* in the iron-stone country of Bahr-el-Ghazal. The exact limits of the two races are not clear.

Family NUMIDIDAE Guinea-fowls

132. *Guttera edouardi* Crested Guinea-fowl 217
 R BR (3, 11, 12)
 The race *sethsmithi* is an uncommon resident of the Imatong foot-hills and Zande gallery forests.
133. *Numida meleagris* Tufted Guinea-fowl 216
 R BR (8 - 10)
 The race *major* is a common resident of the savanna country south of Lat. 10°N and favours Acacia woodland. Often near water and in large flocks. North of Malakal it is replaced by the nominate race.

The following species might occur :

- Francolinus streptophorus* Ring-necked Francolin 193
 (In Nimule, Imatong, Didinga).
- Acryllium vulturinum* Vulturine Guinea-fowl 219
 (In the extreme southeast).

Family TURNICIDAE Button Quails, Quail Plover

134. *Ortyxelos meiffreni* Quail Plover 367
 R? NBR
 Rare: recorded only once, in the dry grassland of the Boma plains in February (Nikolaus pers. obs.).
135. *Turnix sylvatica* Button Quail 365
 R? LM? AM? NBR
 The race *lepurana* is very little recorded. It favours overgrown cultivation and open grassland. Records are from Wau, the Bahr-el-Ghazal, and recently near Renk in February. Always single birds or couples.

Family GRUIDAE Cranes

136. *Antropoides virgo* Demoiselle Crane 246
PMW
No recent records, but small parties have been reported wintering (October to March) as far south as the Sobat River, where they frequented dry *Acacia* grassland and river margins.
137. *Balearica pavonina* Crowned Crane 244
R LM BR (8 - 10)
The race *ceciliae* is a common resident throughout except along the Zaire border and in the extreme southeast; favours the drier margins of rivers and cultivation, often in large flocks. Nests in marshy areas of the Sudd and the River Lol. The race *gibbericeps* might occur as a visitor from Uganda along the southern border.

Family RALLIDAE Rails, Crakes

138. *Crex crex* Corncrake 223
PMW? PMP
Probably an overlooked and uncommon passage migrant, only once recorded (Hartlaub 1881). Numbers recorded in the North Sudan would suggest that it regularly passes through the South.
139. *Crex egregia* African Crake 224
LM? AM? NBR
Only recorded from the Yei area and Juba from July to September, frequenting reeds and moist grassland.
140. *Gallinula angulata* Lesser Moorhen 240
R? LM AM? NBR
Rare: recorded from Tambura, Kagelu and Kajo Kaji, frequenting moist places in woodlands.
141. *Limnocorax flavirostra* Black Crake 225
R BR (7, 11)
A locally not uncommon resident in marshy grasslands and swamps throughout, where it is more frequently seen than other crakes. No records from the forest regions in Zande and Imatong, the arid southeast, or along the Ethiopian border.
142. *Porphyrio alleni* Allen's Gallinule 239
R? LM BR (7, 8)
Locally common during the rainy season in reedbeds and large marshy places. Very common at Aweil in August. Also reported from Geigar (September), Malakal (June) and Meshra er Req (March). It is not clear whether most birds leave the South during the dry season or whether they move to permanent swamps along the big rivers.
- Porzana parva* Little Crake 229
No definite records but occurrences in North Sudan and Uganda indicate that this secretive bird may be wintering or passing through.
143. *Porzana porzana* Spotted Crake 227
PMW? PMP
Probably a fairly common passage migrant and winter visitor, even though records are few. Reported on passage in October and March.

144. *Rallus caerulescens* African Water Rail 222
 R? NBR
 Rare: probably a resident at Gilo/Itibol where it has been recently observed in March and December in thick secondary growth along the Kinyetti River at 2000 m.

145. *Sarothrura affinis* Chestnut-tailed Pygmy Crake 230
 R? BR (4)
 The race *antonii* was collected once (in breeding condition) at Kipia. A bird of short montane grassland.

146. *Sarothrura elegans* Buff-spotted Pygmy Crake 236
 R? LM NBR
 Recorded from Kagelu, near Yei, in October and from Torit in January. A secretive bird of open forest and thickets.

147. *Sarothrura pulchra* White-spotted Pygmy Crake 235
 R? NBR
 The race *centralis* has been recorded only once, at Bengengai in Zande (Cave & Macdonald 1955). A rare secretive bird of swampy localities in deep forests; probably a resident.

The following species might occur :

<i>Gallinula chloropus</i>	Moorhen	239
<i>Porphyrio porphyrio</i>	Purple Gallinule	237
<i>Porzana pusilla</i>	Baillon's Crake	228
<i>Fulica cristata</i>	Red-knobbed Coot	242

Family OTIDIDAE Bustards

148. *Eupodotis hartlaubii* Hartlaub's Bustard 253
 R? LM NBR
 Locally common and probably resident in *Acacia* short grass savanna in the extreme southeast and north of Malakal.

149. *Eupodotis melanogaster* Black-bellied Bustard 258
 R? LM NBR
 Fairly common in the dry season throughout the broad-leafed tall grass savanna. During the rains most birds move to the North.

150. *Eupodotis ruficrista* Buff-crested Bustard 257
 R? LM NBR
 The race *gindiana* is uncommon to rare, frequenting dry thornbush savanna in the extreme southeast. Movements require clarification.

151. *Eupodotis senegalensis* Senegal Bustard 256
 R BR (4)
 The race *canicollis* is a fairly common resident of the dry grassland in the extreme southeast. The nominate race might occur in the northwest.

152. *Neotis denhami* Denham's Bustard 251
 R? LM BR (1, 2)
 The nominate race is the commonest large bustard of open grassland and tree savanna in the South Sudan. It is replaced by the Kori Bustard in the arid southeast and by the Arabian Bustard in the

North. However, all three species might be seen together along the Ethiopian border. Birds leave during the rains for the North. Local movements require clarification, especially in relation to those of Kori and Arabian Bustard.

153. *Neotis heuglini* Heuglin's Bustard 252
 R? NBR
 Status unknown. According to Snow (1978) recorded from the extreme southeast near Lake Turkana.
154. *Otis arabs* Arabian Bustard 249
 LM BR (2, 3)
 The race *butleri* is not uncommon during the dry season when it visits open grassland with few trees or bushes. It occurs from the Nile east to the Ethiopian border and south to Lake Turkana, where it overlaps with the Kori Bustard. Birds move north during the rains.
155. *Otis kori* Kori Bustard 250
 R? LM NBR
 The race *struthiunculus* is a common dry season visitor to the arid region of the extreme southeast. Commonly seen in open savanna on the Boma Plateau. Movements require clarification.

Family JACANIDAE Jacanas

156. *Actophilornis africanus* Jacana 264
 R LM BR (7, 8)
 A very common resident throughout on swamps and open water with floating vegetation. Local movements are associated with seasonal change of habitat.
157. *Microparra capensis* Lesser Jacana 265
 R? LM NBR
 Rare: recorded from Kodok, Aweil, Shambe, Pibor and the south of the Zeraf River. Most records are in the rainy season. Like the Jacana it is associated with floating vegetation. Recent records from Khartoum suggest seasonal movements within the Sudan.

Family ROSTRATULIDAE Painted Snipe

158. *Rostratula bengalensis* Painted Snipe 297
 R? LM AM? NBR
 The nominate race is a very local bird of swamps with mudbanks and thick vegetation. Common from Juba south to Kajo Kaji, and once recorded from Kodok. Records are all from the period August to April.

Family CHARADRIIDAE Plovers

159. *Charadrius alexandrinus* Kentish Plover 269
 PMW PMP
 Common Palaearctic visitor to the White Nile in the North Sudan, but inland records further south are rare. One was ringed at Aweil in September 1979.

160. *Charadrius asiaticus* Caspian Plover 277
PMW PMP
A Palaearctic visitor to semi-arid savanna and grassland, wintering very commonly in North Sudan south to Malakal; occasionally reported from Tonj, Li Rangu and Lado. Expected in the extreme southeast on spring and autumn migration.
161. *Charadrius dubius* Little Ringed Plover 267
PMW PMP
The race *curonicus* is an uncommon but regular Palaearctic visitor from October to April. It favours muddy riverbanks and open grassland near rivers. A young bird ringed in East Germany (August) was controlled in the same autumn (November) at Juba.
162. *Charadrius forbesi* Forbes' Plover 273
LM AM? NBR
Uncommon in open ironstone clearings in tall-grass savanna of Bahr-el-Ghazal west of the river Jur. Several pairs, observed near Wau and Tambura in July and August, showed breeding behaviour. Possibly a breeding visitor during the rainy season.
163. *Charadrius hiaticula* Ringed Plover 266
PMW PMP
The race *tundrae* is a regular Palaearctic visitor in small numbers throughout, frequenting the banks of large rivers. Records at Juba from August to November probably refer to passage birds; records from Wau, Lake No, Sobat, Malakal and Renk are from winter and spring up to May.
Charadrius leschenaultii Great Sandplover 275
A Palaearctic species which might occur along the Nile.
164. *Charadrius mongolus* Mongolian Sandplover 274
PMW
The race *atrifrons* is a rare Palaearctic visitor, recorded once in January 1978 near Juba (Nikolaus pers. obs.); eight birds stayed during day-time at a muddy swamp and flew to dry grassland at night.
165. *Charadrius pecuarius* Kittlitz's Sandplover 271
PMW? LM AM? NBR
Uncommon to rare; the nominate race is recorded in small numbers on short grassland along the Nile at Juba from August to October and in March.
166. *Charadrius tricollaris* Three-banded Plover 272
LM NBR
The nominate race is an uncommon to rare visitor to rivers in the southeast; reported only in November along the Nile near Juba and in December at a sandy river near the Kenya border.
167. *Pluvialis squatarola* Grey Plover 279
PMW PMP
A rare Palaearctic visitor, reported from a muddy riverbank near Juba in October and January (Nikolaus pers. obs.).
168. *Vanellus albiceps* White-headed Plover 290
R? NBR
Rare; recorded only from sandy rivers in Bahr-el-Ghazal (Ibba, Numatina, Sue, Busseri).

169. *Vanellus coronatus* Crowned Lapwing 281
 R LM? NBR
 The nominate race is a fairly common resident in the extreme south-east where it frequents dry open thorn bush country; usually in small numbers.
170. *Vanellus crassirostris* Long-toed Lapwing 293
 R LM? NBR
 The nominate race is common, but confined to the larger permanent waters with floating vegetation, such as the Sobat, Bahr-el-Ghazal and the Nile north to Renk.
171. *Vanellus leucurus* White-tailed Plover 185
 PMW
 A rather uncommon Palaearctic visitor from October to March, frequenting swampy grassland. So far recorded only from Aweil but should occur along the Nile south to Malakal.
172. *Vanellus senegallus* Wattled Plover 291
 R LM? BR (3)
 The nominate race is a fairly common resident throughout, preferring damp localities in savanna and sometimes open grassland. There is indication of some northward migration in the rainy season.
173. *Vanellus spinosus* Spur-winged Plover 286
 PMW? R? LM BR (2, 3)
 A common resident throughout along open waters and swamps. In the non-breeding season it occurs usually in small parties, and shows local movements.
174. *Vanellus tectus* Black-headed Plover 292
 R LM? BR (3, 4, 6, 7)
 The nominate race is a common resident of less arid grassland and Acacia savanna in southeastern and northern areas. Seasonal movements probably occur.

Family SCOLOPACIDAE Sandpipers, Snipes

WADERS

Palaearctic waders are more or less dependent on marshy grassland or mudbanks, which are widespread at the time of autumn migration and up to November/December. Later in winter and during spring migration, such habitat has largely dried up, so that wader numbers in autumn are much greater than in spring.

Sub-family Tringinae

175. *Actitis hypoleucos* Common Sandpiper 312
 PMW PMP
 A common Palaearctic visitor throughout to any type of wetland from July to April. Passage birds are noticed from July to November and again in March and April. Birds ringed in South Africa and Zimbabwe in December were recovered respectively near Torit in April and at Yambio in March

176. *Tringa erythropus* Spotted Redshank 316
PMW PMP
An uncommon to rare Palaearctic winter visitor, mainly from December to March, to the Nile and large rivers in northern Bahr-el-Ghazal. Common at Aweil from late August.
177. *Tringa glareola* Wood Sandpiper 314
PMW PMP
A common Palaearctic visitor and passage migrant to swampy grassland throughout from August to April. Most birds move further south between November and January because of habitat deterioration in the dry season. A bird ringed near Juba in October was recovered in the U.S.S.R. (66°30'N, 70°38'E) in June.
178. *Tringa nebularia* Greenshank 318
PMW PMP
A common Palaearctic visitor in small numbers throughout to sandy rivers and swampy waters, from August to the beginning of May.
179. *Tringa ochropus* Green Sandpiper 313
PMW PMP
A common Palaearctic passage migrant and uncommon winter visitor, mainly to pools in savanna and forest streams, but more widespread on passage. Recorded from August to April with passage from August to October and March to April.
180. *Tringa stagnatilis* Marsh Sandpiper 317
PMW PMP
A common Palaearctic visitor to marshy grassland along the Nile and sandy rivers in Bahr-el-Ghazal from August to April; uncommon south of Wau and Bor.
181. *Tringa totanus* Redshank 315
PMW PMP
A common Palaearctic visitor to marshy grassland along the Nile and sandy rivers in Bahr-el Ghazal from August to April; uncommon south of Wau and Bor.
182. *Xenus cinereus* Terek Sandpiper 311
PMP
Recorded only once from the Nile near Juba in November 1978 on muddy grassland (Nikolaus pers. obs.).

Sub-family Scolopacinae

183. *Gallinago gallinago* Common Snipe 298
PMW PMP
The nominate race is a common Palaearctic visitor to marshy grasslands throughout. Most common during autumn passage from late August to November; rare after February. Two birds ringed at Juba in October were controlled at the same site in November and January of the same winter.
184. *Gallinago media* Great Snipe 299
PMP
Recorded from Wau, Juba and Lado, between September and November and in April, when regularly seen on grassy marshes near larger rivers. A bird ringed near Juba in November was recovered in Kirov region U.S.S.R (58°15'N, 51°E) in August.

185. *Gallinago nigripennis* African Snipe 300
 R? AM? NBR
 The race *aequatorialis* is an uncommon highland species, recorded from Gilo in the Imatong Mountains (April, June and October), where it might breed. A bird at Lado was probably a wanderer.
186. *Lymnocyptes minimus* Jack Snipe 302
 A sporadic winter visitor to swampy grassland from October to January. Common around Juba in November-December 1978, but absent the following year.
- Sub-family Calidritrinae
187. *Calidris alba* Sanderling 308
 PMP
 A rare Palaearctic passage migrant, recorded once from Juba in September 1975 at a flooded gravel pit (Nikolaus pers. obs.).
188. *Calidris alpina* Dunlin 304
 PMW
 A rare winter visitor along the Nile south to Bor which is the southern limit of its wintering range.
189. *Calidris ferruginea* Curlew Sandpiper 303
 PMW PMP
 A regular Palaearctic passage migrant and uncommon winter visitor, recorded mainly from large muddy river banks around Juba and at Aweil rice scheme from August to January.
190. *Calidris minuta* Little Stint 305
 PMW PMP
 A regular Palaearctic passage migrant and winter visitor from August to March, frequenting open mudflats and small pools mainly along and east of the Nile where flocks of up to 100 birds occur together. Small numbers occur on spring passage until May. Rare in Bahr-el-Ghazal. A bird ringed at Juba in November was controlled at Lake Magadi (Kenya) in January.
191. *Calidris temminckii* Temminck's Stint 306
 PMW
 An uncommon Palaearctic winter visitor to marshy grassland; recorded from October to January in small numbers along the Nile south of Bor.
192. *Limicola falcinellus* Broad-billed Sandpiper 307B
 PMP
 A rare Palaearctic visitor; recorded once on wet mud at Aweil rice scheme in August 1978 (Nikolaus 1979).
193. *Limosa limosa* Black-tailed Godwit 319
 PMW
 An uncommon visitor in small numbers; recorded on wet grassland along the Nile and Sobat Rivers and at Aweil from August to March (mainly November to February).
194. *Philomachus pugnax* Ruff 309
 PMW PMP
 Occurs in very large numbers on open marshy grassland along the Nile and around Aweil/River Lol from August to April. From January

onwards numbers decrease greatly due to reduction of suitable habitat in the dry season. Birds ringed in Aweil in August/September were recovered in Algeria in April and U.S.S.R. (62°N, 117°E) in May.

Sub-family Arenariinae

195. *Arenaria interpres* Turnstone 310
 PMP
 The nominate race is a rare Palaearctic passage migrant recorded at Juba in September and October.

Family RECURVIROSTRIDAE Stilts, Avocets

196. *Himantopus himantopus* Black-winged Stilt 296
 PMW AM? LM? NBR
 The nominate race is a common visitor to swamps and rivers throughout from September to April, presumably of Palaearctic origin. Some birds remain in the South Sudan from May to August. No breeding records can be traced.
197. *Recurvirostra avosetta* Avocet 295
 PMW AM? NBR
 A rare Palaearctic winter visitor. Recorded twice along the Nile near Juba in November; also from Baro River, Bahr-el-Ghazal and Kodok.

Family BURHINIDAE Thicknees

198. *Burhinus capensis* Spotted Thicknee 262
 R LM? BR (5 - 7)
 The race *maculosus* is a fairly common resident of dry savanna often noticed at night or in the late evening, but resting under cover during daytime.
- Burhinus oedicephalus* Stone Curlew 260
 Status unknown: the nominate race should occur as is known from northern Uganda, northern Kenya and Gambela, Ethiopia.
199. *Burhinus senegalensis* Senegal Thicknee 261
 R LM? BR (3, 4)
 The race *inornatus* is a common resident throughout, except west of Wau, where it is rare. Always associated with water; often seen in small groups resting along riverbanks during daytime, and becoming active mainly at night.

Family GLAREOLIDAE Coursers, Egyptian Plover, Pratincoles

Sub-family Cursoriinae

200. *Cursorius cursor* Cream-coloured Courser 325
 R? LM? NBR
 The race *littoralis* is uncommon in open arid Acacia grassland in the extreme southeast near Lake Turkana.
201. *Cursorius temminckii* Temminck's Courser
 R LM? NBR
 The nominate race is uncommon; normally found in small flocks in

semi-arid or stony scrub country and in short grassland in the area southeast of Juba - Boma Hills. Records are from October, February and March.

202. *Pluvianus aegyptius* Egyptian Plover 334
 R? LM BR (1 - 4)
 A common breeding visitor in the dry season to rivers with sand-banks. No records for the southeast, the Sudd or the southwestern border region. Most birds move to the North during the rainy season. Birds ringed in Gambela, Ethiopia, in November and April were recovered in North Sudan at Kosti (August) and Khartoum (September).
203. *Rhinoptilus chalcopterus* Violet-tipped Courser 329
 LM AM? BR (1 - 3)
 An uncommon and secretive bird throughout, normally in small flocks and breeding in open savanna during the dry season from September to May. No information about movements during the rainy season.
204. *Rhinoptilus cinctus* Heuglin's Courser 328
 LM? V? NBR
 Status unknown; the nominate race was recorded prior to 1865 between Juba and Lake Turkana. There is only one recent record from Kapoeta in December 1960 (P.J. Dare *in litt.*).

Sub-family Glareolinae

205. *Glareola nordmanni* Black-winged Pratincole 331
 PMW PMP
 Probably a regular but little recorded Palaearctic visitor, easily confused with the Common Pratincole *G. pratincola*. Passage records are from Juba in March and Kajo Kaji in April. There would appear to be a wintering population on the Sudan/Ethiopian border. Up to 5000 were recorded during December/January 1972-73 roosting on sandbanks of the Baro River together with Common Pratincoles.
206. *Glareola nuchalis* White-collared Pratincole 333
 LM? V? NBR
 In the past the nominate race was recorded twice along the Nile, at Nimule and Lado in October, but there are no recent records.
207. *Glareola pratincola* Common Pratincole 330
 PMW R? LM AM? NBR
 Regularly recorded between September and March on or near swampy grassland. Large flocks of up to 5000 birds were seen, together with Black-winged Pratincole *G. nordmanni* roosting on sandbanks of the Baro River (Sudan/Ethiopian border) in December and January. The nominate race is a Palaearctic winter visitor recorded south to Malek, while the African race *limbata* is a common non-breeding visitor from the North Sudan. Birds seen at Juba might be of the race *fuelleborni* which occurs from Lake Turkana west to Uganda. The status of the different races needs clarification.

Family LARIDAE Gulls, Terns

Sub-family Larinae

Larus cirrocephalus Grey-headed Gull 142
No definite record for South Sudan. Since it occurs regularly in autumn south of Khartoum, it is to be expected at Lake No and along the White Nile.

208. *Larus fuscus* Lesser Black-backed Gull 340

PMW

The nominate race is a rare Palaearctic visitor to the Nile; recorded from Malakal and Juba in December.

209. *Larus ridibundus* Black-headed Gull 345

PMW PMP

Single immature birds are uncommon Palaearctic visitors; have been recorded wintering on lakes and small rivers at Malakal, Aweil, Lake Nibor, Terakeka and Maridi. At Lake No, thirty birds including a few adults stayed around a fishing camp in February. The great increase in numbers visiting Kenya and North Sudan has not so far been recorded in the South Sudan.

Sub-family Sterninae

210. *Chlidonias hybridus* Whiskered Tern 362

PMW PMP

The nominate race is an uncommon Palaearctic visitor in small numbers to lakes and swamps in the Sudd; recorded from Malakal and Lake Nibor from December to February.

211. *Chlidonias leucopterus* White-winged Black Tern 361

PMW PMP

The most common tern, occurring on passage from late August to November at any swamp and open water, especially along the Nile. Winters in small numbers at the swamps and lakes of the Sudd, and commonly until March on the Sobat/Baro River and on the Nile north of Malakal.

Chlidonias niger Black Tern 360

There are no definite records.

212. *Gelochelidon nilotica* Gull-billed Tern 349

PMW PMP

The nominate race is a fairly common Palaearctic visitor in small numbers to rivers and swamps throughout. Passage is noticed along the Nile in September/October and in March.

213. *Sterna caspia* Caspian Tern 350

PMW

A few records of single birds from the Nile around Malakal and on the Sobat/Baro River during January/February. This is the southern limit of the wintering range on the Nile.

Family RHYNCHOPIDAE Skimmers

214. *Rhynchops flavirostris* Skimmer 364

LM AM?

NBR

Recorded in small numbers from Sobat River and the Nile in the dry

season (January/February). Moves away during the rainy season, probably to North Sudan to breed.

Family PTEROCLIDAE Sandgrouse

215. *Pterocles exustus* Chestnut-bellied Sandgrouse 369
 R? LM NBR
 Two races occur: the nominate race from the North Sudan visits the dry extreme north around Renk in the dry season; The race *olivascens* is probably resident in the extreme southeast and regularly visits waterholes.
216. *Pterocles lichtensteinii* Lichtenstein's Sandgrouse 373
 R? LM? NBR
 The race *sukensis* is occasionally reported from the extreme south-east in dry, stony open thornbush country.
217. *Pterocles quadricinctus* Four-banded Sandgrouse 375
 R? LM BR (2, 3)
 The most common Sandgrouse in the South Sudan; found in open savanna throughout except in the arid extreme southeast where it is replaced by other species. Most birds leave, probably moving to the North Sudan, during the rainy season.

Family COLUMBIDAE Pigeons, doves

Sub-family Columbinae

218. *Aplopelia larvata* Lemon Dove 398
 R BR (4, 11)
 The nominate race is a common resident of the mountain forests and forest edges of Imatong, Dongotona and Didinga. The smaller Ethiopian race *bronzina* is reported from the forest of Boma Hills.
219. *Columba albinucha* White-naped Pigeon 381
 R? LM? NBR
 Probably rare: recorded only from Bengengai forest in 1981/82 (Hillman in litt.).
220. *Columba arquatrix* Olive Pigeon 380
 R LM? BR (1, 11, 12)
 The nominate race is a common resident in the forests of the Imatong, Dongotona and Didinga Mountains. Most numerous during the dry season after breeding.
221. *Columba delegorguei* Bronze-naped Pigeon 383
 R? NBR
 The race *sharpii* is recorded from the forest regions of the Imatong, Dongotona and Didinga Mountains where it is probably resident. Less common than the Olive Pigeon, which prefers the higher altitudes.
222. *Columba guinea* Speckled Pigeon 379
 R BR (1 - 5)
 The nominate race is a common resident throughout in savanna and towns except in the Sudd region, the green forests of the south, and the arid extreme southeast. Breeds in *Borassus* palms, buildings and rocks.

223. *Columba uncinata* Afep Pigeon 382
 R? NBR
 Probably rare: known only from gallery forests on the Aloma Plateau south of Yei (Macdonald & Cave 1948).
224. *Oena capensis* Namaqua Dove 393
 R LM NBR
 Very common from December to March throughout except in forests; often on roads and cultivation. Most birds move north before the rains, possibly for breeding, but in Gilo the species was recorded in August. Seasonal movements are not well understood.
225. *Streptopelia capicola* Ring-necked Dove 388
 R? LM? NBR
 The race *tropica* occurs in the arid region southeast of Boma/Torit, where its distribution overlaps with that of the Vinaceous Dove.
226. *Streptopelia decipiens* Mourning Dove 387
 R BR (1 - 4, 6 - 10)
 The nominate race occurs from Malakal northwards and along the Ethiopian border; the race *logonensis* is present in the south and west. Fairly common in savanna and cultivation, always near water.
227. *Streptopelia lugens* Dusky Turtle Dove 385
 R LM? BR (8, 9)
 The nominate race is common in the Imatong and Didinga Mountains and Boma Hills around forest edges and cultivation at high altitudes. Commonest from October to December and rare during June/July.
228. *Streptopelia semitorquata* Red-eyed Dove 386
 R BR (9 - 11)
 Common in the better wooded savanna and gallery forests of Boma and southern Equatoria, and west of Wau north to Boro River. Absent from the primary forest on Imatong.
229. *Streptopelia senegalensis* Laughing Dove 392
 R LM BR (1 - 4, 10 - 12)
 The nominate race is common throughout in open savanna, cultivation and villages, often concentrating in large numbers on harvested fields.
230. *Streptopelia turtur* Turtle Dove 384
 PMW
 The wintering range reaches South Sudan in the dry savanna at Malakal and the Boro River. Wintering birds occur regularly in large numbers around Renk when they fly to the Nile for drinking. These are mostly of the nominate race but *arenicola* could occur.
231. *Streptopelia vinacea* Vinaceous Dove 390
 R LM BR (1, 3, 4, 12)
 The most common ring-necked Dove in savanna throughout South Sudan, except in the arid southeast where it is replaced by the Ring-necked Dove.
232. *Turtur abyssinicus* Black-billed Wood Dove 396
 R LM? BR (1, 12)
 Very common throughout in any savanna and cultivation except in forest and gallery forest, where it is replaced by the Blue-spotted Wood Dove.

233. *Turtur afer* Blue-spotted Wood Dove 395
 R NBR
 A fairly common resident of better wooded savanna and gallery forest in the southwest; reported from Boma, Imatong, Juba, Zande, Tonj, Wau and Boro River.

234. *Turturchalcospilos* Emerald-spotted Wood Dove 397
 R NBR
 Recorded only in the southeast, from Didinga and south of Boma, where it replaces the Black-billed Wood Dove.

235. *Turtur tympanistria* Tambourine Dove 394
 R BR (1 - 4, 8)
 A common resident in heavily wooded savanna and forest at Didinga, Dongotona, Imatong, Aloma Plateau, Aza Forest and Zande.

Sub-family Treroninae

236. *Treron australis* Green Pigeon 401
 R? BR (11)
 The race *uluensis* is fairly common in the gallery forests along the Uganda and Zaire border west of the Nile, feeding on fig trees. Birds around Imatong are assigned to the race *gibberifrons*. Two records given by Snow (1978) for the extreme southeast need confirmation.

237. *Treron waalia* Bruce's Green Pigeon 400
 R? LM NBR
 Locally common in savanna with large fig trees, but absent along the Zaire border and the open grass plains of the Sudd. There are no records for July and August and only a few during September to December.

Family PSITTACIDAE Parrots, lovebirds

238. *Agapornis pullaria* Red-headed Lovebird 452
 R NBR
 Uncommon in better wooded savanna in the southwest; recorded from Kajo Kaji, Yei, Maridi, Bengengai, from Tambura north to Wau and Tonj, and at the Boro River. The racial status is unclear, as both the nominate race and *ugandae* could occur.

239. *Poicephalus crassus* Niam Niam Parrot 446
 R NBR
 Rare in the extreme southwest where recorded only a few times from forested localities around Yambio.

240. *Poicephalus meyeri* Brown Parrot 449
 R NBR
 Fairly common in better wooded savanna southwest of Gogrial - Amadi - Kapeta, except for a small area between Rumbek and Bengengai which separates its distribution into two areas. All birds have been assumed to be of the nominate race, but populations along the southern border could belong to the race *saturatus*.

241. *Psittacula krameri* Rose-ringed Parrakeet 450
 R BR (8)
 The nominate race is fairly common but rather local in all types of

savanna from the Kenya border to Juba, Lado and Maridi, in the region Tonj - Wau - Boro River, and north to the Bahr-el Arab. There are no records from the southwest, but it might be overlooked.

Family MUSOPHAGIDAE Turacos

242. *Corythaeola cristata* Great Blue Turaco 437
R NBR
Fairly common resident in the foothill forests of Imatong up to 1800m and in the gallery forests of Zande. Sometimes in parties of up to twenty birds.
243. *Corythaixoides leucogaster* White-bellied Go-away Bird 440
R LM NBR
Fairly common in *Acacia* savanna in the southeast. Some birds move as far west as Maridi and Mvolo in the dry season from January to March. There is an isolated record from *Acacia* woodland at Ayod south of Malakal in February (G. Nikolaus pers. obs.).
244. *Crinifer zonurus* Eastern Grey Plantain Eater 438.
R BR (12)
Common throughout savanna southwest of Kapoeta - Lado - Rumbek - Aweil - Kafia Kingi.
245. *Musophaga rossae* Ross' Turaco 436
R NBR
Rare: there are only a few records from Didinga Mountains, Aloma Plateau and Zande, where it occurs in larger gallery and "bowl" forests; not recorded from Imatong.
246. *Tauraco leucolophus* White-crested Turaco 433
R NBR
Fairly common and probably resident southwest of Didinga - Lado - Tonj - Nyamyell in better wooded savanna, gallery forests and forest edges at low altitude.
247. *Tauraco leucotis* White-cheeked Turaco 425
R NBR
The nominate race is a common resident in better wooded savanna on Boma Hills.
248. *Tauraco schuetti* Black-billed Turaco 430
R NBR
The race *emini* is quite common in the higher forests of the Imatong and Dongetona Mountains, and uncommon in the forests of Mt. Lotuke, Didinga. The race *finschi* is rather uncommon in the Zande forests and on the Aloma Plateau.

Family CUCULIDAE Cuckoos, Coucals

Sub-family Cuculinae

249. *Chrysococcyx caprius* Didric Cuckoo 417
R LM BR (10)
Fairly common in dry savanna throughout. Mainly resident, although some move north during the rainy season. Parasitic mainly on weavers, bishops and sparrows.

250. *Chrysococcyx cupreus* Emerald Cuckoo 416
R LM BR (4 - 6)
The nominate race is common, calling from March to October, in the forest regions of Imatong, Aloma Plateau, Aza Forest and Zande. Records from Imatong in December and January suggest that it is mainly resident. Wanderers have been reported from Wau (July) and Meshra er Req (May), but these records need confirmation. Parasitic on flycatchers, shrikes, bulbuls, warblers and sunbirds.
251. *Chrysococcyx flavigularis* Yellow-throated Green Cuckoo 419
R? LM? NBR
A rare forest species known only from a sight record from Bengengai in March 1977 (Traylor & Archer 1982).
252. *Chrysococcyx klaas* Klaas' Cuckoo 418
R LM? BR (1, 10 - 12)
Uncommon in savanna during the dry season; reported from Gogrial, Ayod, Maridi, Boma and from Juba to Kapoeta. In Imatong it is a fairly common resident in secondary growth and forest edges. Birds ringed in Gilo in May 1978 and July 1979 were controlled there in December 1980 and January 1980 respectively. Reported hosts are flycatchers, warblers, sunbirds and weavers.
253. *Clamator glandarius* Great Spotted Cuckoo 413
PMW? LM R? NBR
Uncommon in savanna; recorded all months except July, October and December, at Boma, Juba, Torit, Kajo Kaji, Opari, Li Rangu, Rumbek, Tonj, Wau, Aweil, Gogrial and Renk. Status and movements are not clear because both African and Palaearctic birds could occur. Parasitic mainly on Pied Crow, but also on Cape Rook and Piapiac.
254. *Clamator jacobinus* Black and White Cuckoo 415
PMW? AM? LM? BR (8, 10)
The race *pica* is a fairly common visitor to savanna from April to October, mainly southwest of Kapoeta-Bor-Aweil. Breeding is reported from Torit (breeding condition in October) and Wau (August). There is one December record from the Kenya border in dry *Acacia* savanna. The South African race *serratus* should occur, but the racial status needs clarification. Parasitic on babblers.
255. *Clamator levallantii* Levallant's Cuckoo 414
LM? AM? NBR
An uncommon visitor in March/April and July/September to the better wooded savanna of the southwest where recorded north to Wau. A lack of records for May and June suggests movement to the north to breed. Parasitic on babblers.
256. *Cuculus canorus* Eurasian Cuckoo 404
PMP
Recorded only on autumn passage, when small numbers occur regularly in savanna between late August to early November with a peak in September. The races *canorus* and *telephonus* have occurred.
257. *Cuculus clamosus* Black Cuckoo 407
R? LM? AM? NBR
Status unknown. The nominate race was once collected from the Laboni Forest in Imatong in March 1936 (specimen in Khartoum Museum). Parasitic on bulbuls, robin chats and small shrikes.

258. *Cuculus gularis* African Cuckoo 404
 AM? LM? R? NBR
 Uncommon in savanna; reported at Bahr-el Zeraf (June), Yambio (December), Aloma Plateau (March), Lado (April) and Torit (October - November). Hosts are drongos, scrub robins and glossy starlings.
259. *Cuculus solitarius* Red-chested Cuckoo 406
 R LM AM? BR (3 - 6)
 A common resident of savanna and forests mainly south of 5 N, but also recorded from Boro River in April. It is stimulated to call by rain, otherwise unobtrusive. Robin chats are the main hosts.
260. *Pachycoccyx audeberti* Thick-billed Cuckoo 412
 AM? R? NBR
 The race *validus* is rare, recorded only from gallery forests in Zande; probably resident.

Sub-family Phoenicophaeinae

261. *Ceuthmochares aereus* Yellowbill 424
 R NBR
 The nominate race is a little recorded resident of dense thickets and riverine forest west of the Nile in the extreme south.

Sub-family Centropinae

262. *Centropus grillii* Black Coucal 420
 R? NBR
 An uncommon bird of swamps and marshes, reported from Aweil, Tonj, Tambura, Lado, Juba, north of Torit and Boma. Birds in breeding plumage suggest that nesting occurs during the late rainy season (August/September).
263. *Centropus monachus* Blue-headed Coucal 421
 R BR (6, 8)
 The race *fischeri* occurs commonly throughout and intergrades with the nominate race along the Ethiopian border. It inhabits dense parts of secondary growth and forest clearings up to 3000 m in Imatong, as well as papyrus and grass swamps with dense thickets near rivers and lakes in the Sudd. Birds with blue rather than violet glossed heads restricted to papyrus and reeds in the Sudd have been described as '*C. heuglini*' and might prove to be a valid species.
264. *Centropus senegalensis* Senegal Coucal 422
 R BR (3, 10)
 The nominate race is an uncommon resident but widely distributed throughout, frequenting thickets near swamps and also dry thick bushland. Less common than the Blue-headed Coucal.
265. *Centropus superciliosus* White-browed Coucal 423
 R NBR
 The nominate race is a fairly common resident in *Acacia* savanna with scrubby thickets. It occurs mainly in the southeast and north of Renk.

Family TYTONIDAE Barn Owls

266. *Tyto alba* Barn Owl 528
 R BR (1, 2)
 The race *affinus* is uncommon throughout, frequenting towns, and rocks and hollow trees in savanna. Reported from Lake No, Aweil, Yei, Juba, Torit and Jebel Kasangor.

Family STRIGIDAE Owls

267. *Asio capensis* African Marsh Owl 532
 R? LM? AM? NBR
 The nominate race occurs in swamps in the Sudd region. Butler reported very common in papyrus along the Nile and Bahr-el-Ghazal in January and February. Otherwise records are few. Status and movements unclear.
268. *Bubo africanus* Spotted Eagle Owl 543
 R BR (1, 2)
 The race *cinarascens* is common throughout in savanna and towns, where hills and large trees are available. Often noted at night sitting on roads.
269. *Bubo lacteus* Verreaux's Eagle Owl 544
 R BR (1)
 Widely distributed; locally common at forest edges of Imatong and Didinga up to 2500m, and in gallery forest at Aloma Plateau and Zande. A few occur along the larger rivers, the White Nile, Sobat, Bahr-el-Ghazal and Jur where large trees are available.
270. *Ciccaba woodfordii* African Wood Owl 533
 R BR (12)
 Two races occur: *umbrina* on the Boma Plateau, and *bohndorffi* along the southern border from the Didinga Mountains to Zande. Probably not uncommon in forest and gallery forest at low altitude.
271. *Glaucidium perlatum* Pearl-spotted Owlet 538
 R NBR
 There are a few records of the race *licua* from *Acacia* woodland, in The southeast and north of Malakal and Wau.
272. *Otus leucotis* White-faced Scops Owl 536
 R NBR
 The nominate race is widespread, but little recorded, in *Acacia* savanna and open woodland near Boro River, Bahr-el-Zeraf, Lado and the Ilemi Triangle.
273. *Otus scops* Scops Owl 534
 PMW
 Winters commonly in North Sudan but recorded at Tonj in January and at Juba and Malakal in March. Habitat ranges from open savanna and gallery forest to tall reeds. In addition to the nominate race *pulchellus* could occur.
274. *Otus senegalensis* African Scops Owl 534
 R? NBR
 The nominate race is common throughout in savanna southwest of Kapoeta - Lado - Tonj - Bahr el Arab, and on the Boma Plateau. Its call is most frequently heard in the dry season.

275. *Scotopelia peli* Pel's Fishing Owl 546
 R NBR
 Rare: Snow (1978) mentions one record from the Akobo River, and it has recently been photographed at Mbili on the Jur River (J.S. Fairclough in litt.).

The following species might occur, but have not been recorded so far :

- Asio flammeus* Short-eared Owl 531
 (open grassland and swamps of the Sudd)
Asio otus Long-eared Owl 530
 (Ethiopian border)
Glaucidium tephronotus Red-chested Owlet 539
 (Imatong and Zande)

Family CAPRIMULGIDAE Nightjars

276. *Caprimulgus aegypticus* Egyptian Nightjar 561
 PMW
 The nominate race is an uncommon winter visitor from November to March to arid localities north of Malakal; recorded from Kaka and near Lake No.
277. *Caprimulgus clarus* Slender-tailed Nightjar 565
 R BR (3 - 5)
 A common resident east of the Nile in open wet grassland and dry short grassland with scattered bushes. Overlaps along the Nile with Long-tailed Nightjar.
278. *Caprimulgus climacurus* Long-tailed Nightjar 565
 R? LM BR (2 - 4)
 Two races breed: A common savanna species which hunts at night over swamps and marshes, *nigricans* mainly north of 7°N and *sclateri* south of 7°N but mainly west of the Nile. The nominate race which breeds in North Sudan, is probably only a dry season visitor. All three races have been reported from Lado. Both local races move north within their range to breed. Overlaps in the east with Slender-tailed Nightjar.
279. *Caprimulgus donaldsoni* Donaldson-Smith's Nightjar 552
 LM NBR
 Seasonally common in dry *Acacia* scrub in the Ilemi Triangle in February and June; absent in November and December. Status and movements little known.
280. *Caprimulgus europaeus* European Nightjar 547
 PMP?
 Collected by Emin Pasha at Lado (Hartlaub 1881; date and race unrecorded) No other records.
281. *Caprimulgus fraenatus* Dusky Nightjar 548
 R? LM? NBR
 Uncommon; recorded only from the extreme southeast, frequenting thorn scrub in rocky country.

282. *Caprimulgus inornatus* Plain Nightjar 556
 AM LM? R? NBR
 An intra-African migrant, which breeds in dry savanna in North Sudan between April and October. Passage noted regularly in March/April and October/November. Status in the extreme south from Zande to Boma is unclear; this may be the northern limit of the non-breeding range, but there is also probably a resident population in northern Uganda and northern Kenya.
283. *Caprimulgus natalensis* White-tailed Nightjar 557
 R NBR
 The race *chadensis* is very locally distributed on wet grassland, recorded only from Lake No to Malakal, along the Sobat River to the Ethiopian border, and along the southern border from Maridi to Kajo Kaji. The lack of records from the Sudd, where its typical habitat is abundant, is unexpected.
- Caprimulgus nubicus* Nubian Nightjar 553
 This species must be rejected from the South Sudan list. The two specimens from Juba in January 1979 (Nikolaus 1979) were re-identified as juvenile Slender-tailed Nightjars.
284. *Caprimulgus pectoralis* Fiery-necked Nightjar 551
 R? NBR
 The race *nigriscapularis* frequents better wooded savanna and gallery forest along the southern border from Imatong to Zande, where it is fairly common, and possibly also in Boma. All records to date are from March to July, but it might prove to be resident.
285. *Caprimulgus poliocephalus* Montane Nightjar 558
 R BR (2, 3)
 A common resident of grassland and overgrown cultivation above 1500 m on the Didinga, Dongotona and Imatong Mountains.
- Caprimulgus rufigena* Rufous-cheeked Nightjar 550
 No records, but this intra-African Migrant should occur in the west on passage or as a non-breeding visitor from April to August.
286. *Caprimulgus stellatus* Star-spotted Nightjar 555
 R? LM? NBR
 The race *simplex* is rare in dry *Acacia* scrub in the southeast where reported from Torit and near the Kenya border in May.
287. *Caprimulgus tristigma* Freckled Nightjar 554
 R BR (4)
 The nominate race is fairly common but extremely local, and is restricted to rocky hills up to 2000 m. Recorded from Kajo Kaji to Imatong and Boma; records around Tambura might refer to the West African race *sharpei*.
288. *Macrodipteryx longipennis* Standard-winged Nightjar 563
 AM LM R? BR (2 - 5)
 A common breeding visitor to savanna throughout from August to May. Most numerous, especially on roads at night, in April/May, when it moves to the north; it returns south in August. Males are in breeding plumage from November to April.

289. *Macrodipteryx vexillarius* Pennant-winged Nightjar 564
AM NBR
This intra-African migrant is a common non-breeding visitor from February to August to broad-leaved tree savanna in the southwest. It ranges north to Torit, Lado and Wau, and occasionally to Darfur in the North Sudan. Males leave in August with fully developed pennants.

Family APODIDAE Swifts

290. *Apus aequatorialis* Mottled Swift 641
AM? NBR
There are a few records of passage birds along the southern border: from Kapoeta (50 moving north in February); Maridi, Yei, Juba and the Aloma Plateau; single birds moving south during July/August; Upper Talanga (Imatong); 100 moving south during October. These movements support observations by Lynes of summering in Darfur.
291. *Apus affinis* Little Swift 643
R? LM? AM? PMW? NBR
Status uncertain. Only a few records, probably of the nominate race: many were seen at Jebel Kasangor and Boma in February including flocks roosting in caves, a few frequented bridges at Nzara and Tambura in August; a few at Juba in October, coming to drink in the morning with Common and Alpine Swifts.
292. *Apus apus* Common Swift 636
PMP PMW
Common throughout in fairly large numbers on autumn and spring passage during late August/September and April/early May. Only a few remain to winter. The seasonal status of the two races, nominate and *pekinensis* needs clarification.
293. *Apus caffer* White-rumped Swift 644
R? LM NBR
Probably resident. The race *streubelii* was found roosting together with Red-rumped Swallows under a bridge north of Juba. Small numbers sometimes occur with other swifts around Imatong and Aweil from late August to October.
294. *Apus melba* Alpine Swift 640
PMW PMP
The nominate race is a regular but uncommon passage migrant from September to November and from February to April, mainly along the Nile and eastwards to the Ethiopian border.
295. *Apus pallidus* Pallid Swift 638
PMW V?
This species, probably the race *illyricus* has been collected by Traylor at Bengengai, far south of its known winter quarters at Jebel Marra (Traylor & Archer 1982).
296. *Cypsiurus parvus* Palm Swift 646
R LM BR (3 - 5)
The nominate race is a very common resident throughout wherever there are palms, except along the southern border where it is replaced by *myochrous*. There is some north-south movement, often together with other swifts.

297. *Schoutedenapus myoptilus* Scarce Swift 639
 LM AM? NBR
 A common breeding visitor to the forest region of Imatong from late March to late September. It probably breeds in cliffs above 2000 m.

The following species might occur but have not yet been recorded :

- Apus barbatus* African Black Swift 636
Apus horus Horus Swift 645
Apus niansae Nyanza Swift 637
Telacanthura ussheri Mottle-throated Spinetail 647

Family COLIIDAE Mousebirds

298. *Colius striatus* Speckled Mousebird 566
 R BR (3, 4)
 A common resident in broad-leaved tree savanna and forest edges, though less common in the north and absent from the northwest and the Ilemi Triangle. Replaced by the Blue-naped Mousebird in Acacia woodland. Three races occur: *leucotis* at Boma and along the Ethiopian border, *leucophthalmus* in Zande along the Zaire border and *jebelensis* elsewhere.

299. *Urocolius macrourus* Blue-naped Mousebird 568
 R BR (3, 4, 9)
 Fairly common in Acacia savanna and arid scrub country. The race *griseogularis* occurs to about 9°N, north of where it is replaced by the nominate race; *pulcher* occurs in the southeast, east of Torit and south of Boma.

Family TROGONIDAE Trogons

300. *Apaloderma narina* Narina's Trogon 570
 R? NBR
 Uncommon, but easily overlooked in forest and better wooded savanna. The nominate race occurs from Kajo Kaji to Didinga and Boma. The race *brachyurum* should occur west of the Nile in Zande, but has not yet been recorded.

Family ALCEDINIDAE Kingfishers

Sub-family Cerylinae

301. *Ceryle maxima* Giant Kingfisher 466
 R LM BR (11)
 Uncommon along permanent rivers in better wooded savanna and forest. Records from Boro River, River Lol, Wau, Tambura, Maridi, Aloma Plateau, Juba, and in the Imatongs up to 2000 m.
302. *Ceryle rudis* Pied Kingfisher 465
 R LM BR (1, 8)
 Common at rivers, lakes and pools throughout except in the south-east where records are few. Performs fairly long distance movements which need clarification.

Sub-family Alcedininae

303. *Alcedo cristata* Malachite Kingfisher 470
 R? LM?? NBR
 The nominate race is common at swamps and rivers throughout except in the Ilemi Triangle and the forests of Imatong.
304. *Alcedo quadribrachys* Shining Blue Kingfisher 469
 R? NBR
 The race *guentheri* is uncommon, frequenting forest streams along the Zaire border from the Aloma Plateau to Zande.
305. *Alcedo semitorquata* Half-collared Kingfisher 468
 R NBR
 Uncommon: recorded only from Boma, where it was collected along a fast-running woodland stream (Nikolaus 1979).
306. *Halcyon badia* Chocolate-backed Kingfisher 478
 R? NBR
 Uncommon in the forests of Zande, where collected only at Bengengai (Traylor & Archer 1982).
307. *Halcyon chelicuti* Striped Kingfisher 479
 R? LM NBR
 The nominate race is common throughout, mainly in the drier parts of savanna and in cultivation; this species is not dependent on water. Most birds move north from May to September.
308. *Halcyon leucocephala* Chestnut-bellied Kingfisher 477
 LM AM? BR (3, 4)
 The nominate race occurs in savanna throughout during the dry season. It arrives along the southern border between late November and January, and moves north to breed on sandy river banks during March/April. In Darfur, breeding birds are present from April to to June. Southward movement during May/June is seldom noticed. There are single records from Juba in July and September which might refer to the southern race *pallidiventris*.
309. *Halcyon malimbica* Blue-breasted Kingfisher 475
 R NBR
 The nominate race is fairly common in gallery and bowl forests on the Zaire border where recorded from Aloma Plateau, Aza Forest, Zande and Tambura.
310. *Halcyon senegalensis* Woodland Kingfisher 473
 LM AM NBR
 The nominate race is fairly common and widely distributed. Present in southern areas only in the dry season (September to March) and in northern areas from March to July; it arrives in Darfur in June. The southern race *cyanoleuca* is occasionally reported in the rainy season from Torit (May) and Upper Nile Province. The status and movements of the two races need clarification.
311. *Ispidina lecontei* Dwarf Kingfisher 472
 R? NBR
 Uncommon in the forests of Zande where collected by Traylor (Traylor & Archer 1982).

312. *Ispidina picta* Pygmy Kingfisher 471
 R AM BR (3, 4)
 The nominate race is a common resident, mainly in better wooded savanna and gallery forest, and is absent or rare in the Sudd region. It is not dependent on water. The Intra-African migrant race *natalensis* was common in June/July during its non-breeding season in the lowland forests of Imatong and the Aloma Plateau.
- Family MEROPIDAE Bee-eaters
313. *Merops albicollis* White-throated Bee-eater 486
 LM AM NBR
 A fairly common non-breeding visitor from North Sudan. It frequents various types of savanna, mainly along the southern borders, from January to March, also recorded occasionally during April to June, August and November.
314. *Merops apiaster* Eurasian Bee-eater 481
 PMP PMW?
 A common passage migrant during September to November and March/April. There are records also for August, January and February.
315. *Merops breweri* Black-headed Bee-eater W 447
 R? LM? NBR
 There are several sight records of this forest species from Bengengai during February and March 1977 - 82 (Traylor & Archer 1982, Hillman in litt.)
316. *Merops bulocki* Red-throated Bee-eater 492
 R LM BR (1 - 3, 12)
 The race *frenatus* is a fairly common resident in broad-leaved tree savanna southwest of Torit - Lado - Tonj - Wedweil - Kafia Kingi and around Kongor and the Baro and Akobo Rivers on the Ethiopian border. It breeds in high river banks.
317. *Merops hirundineus* Swallow-tailed Bee-eater 496
 R? LM NBR
 The race *heuglini* is locally quite common in better wooded savanna southwest of the line Kapoeta - Bor - Boro River, but it also frequents wooded areas on the Ethiopian border at the Baro River. Information is needed on seasonal movements; records are scarce from April to August when birds are presumably further south.
318. *Merops nubicus* Carmine Bee-eater 484
 R LM BR (3)
 The nominate race is a common breeding visitor in the dry season from November to April. During the rains most birds depart for North Sudan, but it is recorded all the year round at Juba. Breeds in very large colonies in steep river banks.
319. *Merops oreobates* Cinnamon-chested Bee-eater 489
 R BR (1, 2)
 A common resident of forest edges, clearings and cultivation on the Imatong and Didinga Mountains above 1800 m.
320. *Merops orientalis* Little Green Bee-eater 483
 R LM BR (3)
 A fairly common resident of wooded savanna with sandy soil;

recorded from Kapoeta to Lado, at Amadi, and north to Malakal and Renk as well as at Aweil and Wedweil. The racial situation is not clear; records along the Nile and to the east possibly all refer to *cleopatra* but western birds at least are *viridissimus*.

321. *Merops persicus* Blue-cheeked Bee-eater 482
PMW
The nominate Palaearctic race is a common winter visitor to swamps rivers of the Sudd region, mainly from October to March. The first birds arrive in the Aweil - Malakal region in late August.
322. *Merops pusillus* Little Bee-eater 488
R BR (3)
Fairly common throughout, mainly in open bushed grassland. Three races occur: the nominate race west of about 30 E and *ocularis* east of 30 E, itself replaced by *cyanostictus* in the extreme south-east. The exact range of these races needs further clarification.
323. *Merops superciliosus* Madagascar Bee-eater 482
AM NBR
An uncommon to rare non-breeding visitor, reported from Kapoeta in May/June and at Bongo south of Juba in August.
324. *Merops variegatus* Blue-breasted Bee-eater 490
R NBR
The race *lafresnayii* is fairly common in grassy savanna on the Boma Plateau. The race *loringi* could occur around the Didinga Mountains since it has been reported from Kidepo National Park in Uganda.

Family CORACIIDAE Rollers

325. *Coracias abyssinica* Abyssinian Roller 458
LM NBR
Seasonally common from October to April throughout savanna and dry open grassland with bushes. In the Torit - Lake Turkana region it also occurs in May/June. There are thus distinct seasonal movements. The breeding grounds are probably in North Sudan and Ethiopia.
- Coracias caudata* Lilac-breasted Roller 460
The race *lorti* occurs in dry *Acacia* savanna around Lake Turkana but has not yet been recorded from the Sudan.
326. *Coracias cyanogaster* Blue-bellied Roller 462
R? LM? NBR
Status unknown; recorded rarely in woodland between Kajo Kaji and Maridi, but not recently.
327. *Coracias garrulus* Eurasian Roller 457
PMP
The nominate race is a fairly common Palaearctic autumn passage migrant in October/November in dry *Acacia* savanna east of the Nile. There are no spring records.
328. *Coracias naevia* Rufous-crowned Roller 461
LM AM? BR (5)
The nominate race is uncommon in dry woodland. Small numbers are recorded from Wau, Tonj, Amadi, Lado, Kajo Kaji and the southeast.

Breeding birds (Wau to Darfur) return from the south during March to May, and leave again in late July. Non-breeders are present in the southeast from November to March. There are no records from August to October.

329. *Eurystomus glaucurus* Broad-billed Roller 463

LM AM? BR (4, 5)

The race *afer* is a common breeding visitor from March to August in better wooded savanna north of Wau. In August 1978 very strong southward passage (up to 500 birds in one afternoon) was noticed at various places west of the Nile. The race *suahelicus* and the nominate Malagasy race could occur as visitors.

Family UPUPIDAE Hoopoe

330. *Upupa epops* Hoopoe 517

PMW PMP LM AM? BR (10)

A fairly common winter visitor in savanna throughout, but mainly in dry *Acacia* woodland. Except in the southeast nearly all records are from December to March and refer to nominate Palaearctic visitors. The race *senegalensis* which breeds uncommonly in the North Sudan and in the extreme southeast is probably a non-breeding visitor throughout. More information is required on racial status and movements. The southern race *africana* can be expected since it has recently been recorded in Darfur in July (G. Nikolaus pers. obs.).

Family PHOENICULIDAE Wood Hoopoes

331. *Phoeniculus aterrimus* Black Wood Hoopoe 523/525

R NBR

P. cavei (Macdonald 1946) is treated as synonymous with this species. The race *emini* is an uncommon resident, usually in pairs, and frequents dry open country throughout.

332. *Phoeniculus bollei* White-headed Wood Hoopoe 522

R NBR

The race *jacksoni* is a common resident of the Imatong forests, mainly above 2000 m, but occasionally in the foothills; normally in small parties.

333. *Phoeniculus minor* Abyssinian Wood Hoopoe 527

R NBR

The race *cabanisi* is a fairly common resident in the dry woodland southeast of Boma - Lado - Juba.

334. *Phoeniculus purpureus* Green Wood Hoopoe 519

R LM? NBR

The race *niloticus* is a common resident and wanderer throughout in all types of woodland, but absent from the forest along the southern border and the open Sudd region; less common in the north. Usually moves around in noisy troops. The race *marwitzi* could occur along the Uganda and Kenya borders.

Family BUCEROTIDAE Hornbills

Sub-family Bucerotinae

335. *Bycanistes brevis* Silvery-cheeked Hornbill 501
R? NBR
Fairly common and probably resident in better woodland and gallery forest on the Boma Hills.
336. *Bycanistes subcylindricus* Black and White Casqued Hornbill 500
R LM NBR
The race *subquadratus* is a fairly common resident in better wooded savanna and gallery forest southwest of Didinga, Amadi, Aweil and Boro River. There is indication of northward movement during the rains.
337. *Ceratogymna atrata* Black-wattled Hornbill 503
R NBR
An uncommon resident of bowl and gallery forests of the Aloma Plateau and at Bengengai.
338. *Tockus alboterminatus* Crowned Hornbill 509
R NBR
The race *geloensis* is an uncommon resident in better wooded savanna and gallery forests of the Boma Hills.
339. *Tockus camurus* Red-billed Dwarf Hornbill 514
R BR (2)
The nominate race is a fairly common resident of gallery forest in Zande, often near swampy ground.
340. *Tockus erythrorhynchus* Red-billed Hornbill 505
R LM? NBR
The nominate race is a common resident throughout in woodland and wooded savanna, absent only from the forest regions of the south and open grasslands from the Sudd to the Ethiopian border.
341. *Tockus fasciatus* Pied Hornbill 510
R BR (3)
The nominate race is a fairly common resident of gallery and bowl forests along the Zaire border from the Aloma Plateau to Aza Forest and Bengengai.
342. *Tockus flavirostris* Yellow-billed Hornbill 506
R NBR
The nominate race is an uncommon resident in hilly dry *Acacia* savanna in the extreme southeast.
343. *Tockus hemprichii* Hemprich's Hornbill 512
R? LM? NBR
Rare but possibly resident; recorded from Didinga, Jebel Kasangor and Boma, frequenting rocky hills in semi-arid country.
344. *Tockus jacksoni* Jackson's Hornbill 508
R NBR
A common resident in dry *Acacia* woodland in the southeast, east of Torit.

345. *Tockus nasutus* Grey Hornbill 504
 R LM BR (1, 2)
 The nominate race is a common resident throughout in any kind of woodland, absent only from the forests in the south and the open grasslands of the Sudd. Some birds move north during the rains. Appears to concentrate seasonally in roosting areas giving the impression of migration.
- Sub-family Bucoracinae
346. *Bucorvus abyssinicus* Abyssinian Ground Hornbill 516
 R? LM? BR (6 - 8)
 Uncommon, usually in pairs or small parties, in open woodland or tree savanna southwest of Kapoeta - Jebel Kasangor - Lake Nibor - Tonj - Aweil - Bahr-el-Arab. It occurs up to 2000 m on the Didinga Mountains.
- Family CAPITONIDAE Barbets, Tinkerbirds
347. *Gymnobucco bonapartei* Grey-throated Barbet 586
 R NBR
 The race *cinereiceps* is a locally common resident of forest and forest edge up to 2500 m, usually frequenting treetops in small noisy parties. Recorded at Bengengai, Imatong and the Dongotona Mountains.
348. *Lybius bidentatus* Double-toothed Barbet 573
 R BR (10)
 Fairly common resident in better wooded savanna, gallery forest and forest edge at low altitude, frequenting large fig trees. The race *aethiops* occurs at Boma and *aequatorialis* from Didinga to Lado and Tambura.
349. *Lybius guifsobalito* Black-billed Barbet 584
 R NBR
 A fairly common resident in better wooded savanna east of Malakal - Li Rangu, except in the arid extreme southeast and the open grasslands of the Sudd; often in gardens.
350. *Lybius hirsutus* Hairy-breasted Barbet 582
 R NBR
 The race *ansorgii* is an uncommon resident along the Zaire border from the Aloma Plateau to Aza Forest and Zande, frequenting fig trees in gallery and bowl forest.
351. *Lybius lachrymosus* Spotted-flanked Barbet 584
 R NBR
 The nominate race is uncommon in better wooded savanna and gallery forest at low altitude. Recorded to date only from the Didinga Mountains, but it could occur west to Kajo Kaji.
352. *Lybius leucocephalus* White-headed Barbet 576
 R NBR
 The nominate race is a fairly common resident of well wooded savanna. It occurs mainly west of the Nile, but occasionally east to Kidepo River.

533. *Lybius leucomelas* Red-fronted Barbet 585
 R BR (1, 8, 11)
 The nominate race is an uncommon resident of dry bushland and dry open woodland. Apparently restricted to the area Juba - Didinga - Bor, but it could possibly occur north to Shambe.
534. *Lybius melanocephalus* Black-throated Barbet 583
 R BR (12)
 The race *stigmatothorax* is a common resident of dry thornbush country in the extreme southeast, frequenting thick bushes.
535. *Lybius rolleti* Black-breasted Barbet 572
 R NBR
 A fairly common resident in better wooded savanna southwest of Kapoeta - Tonj - Bahr-el-Arab, frequenting large fig trees.
536. *Lybius vieillotii* Vieillot's Barbet 578
 R BR (7, 8)
 The nominate race is locally common but widely distributed in dry woodland with thick bushes, mainly west of the Nile. Recorded from Amadi, Raffile, Wau, Tonj, Aweil, Gogrial and from Malakal to Renk.
537. *Pogoniulus chrysoconus* Yellow-fronted Tinkerbird 595
 R BR (4.8)
 The nominate race is a common resident throughout in dry woodland with thick bushes, except in the arid extreme southeast and the forests along the southern border.
538. *Pogoniulus leucolaima* Lemon-rumped Tinkerbird 596
 R BR (1, 11, 12)
 The race *mfumbiri* is a common resident of forest and forest edge up to 2200m along the southern border, from Bengengai to Aza Forest, the Aloma Plateau, Imatong, Dongotona and Didinga.
539. *Pogoniulus pusillus* Red-fronted Tinkerbird 594
 R NBR
 The race *affinis* is a common resident on rocky hills in dry woodland from the foothills east of Imatong to Lake Turkana.
536. *Trachylaemus purpuratus* Yellow-billed Barbet 604
 R BR (3)
 The race *elgonensis* is a fairly common resident of forest, forest clearings and secondary growth on the Imatong Mountains up to 2200m; also recorded at Bengengai.
531. *Trachyphonus darnaudii* d'Arnaud's Barbet 603
 R BR (1, 2)
 The nominate race is a very common resident of dry open savanna, even found in gardens. It occurs, usually in pairs, in the region southeast of Boma - Bor - Amadi - Juba.
532. *Trachyphonus erythrocephalus* Red and Yellow Barbet 601
 R BR (5)
 The race *versicolor* is a fairly common resident in the dry scrub country with hard red soil in the extreme southeast from Torit to the Ethiopian border.

363. *Indicator exilis* Least Honeyguide 610
 LM BR (1, 3)
 The race *pachyrhynchus* is a fairly common but local breeding visitor from October to March, frequenting forest, secondary growth and well-wooded savanna. It is recorded from Bengengai, Bongo River, near Aweil, north of Boro River and Imatong. It is parasitic on tinkerbirds.
364. *Indicator indicator* Black-throated Honeyguide 605
 R? LM NBR
 The nominate race is locally common in woodlands, gardens and open bush savanna throughout. Recorded from November to April and exceptionally between May and September on the Uganda border. Large numbers caught in April north of the Boro River might have been on migration to North Sudan. Parasitic on hole nesting birds such as barbets, woodpeckers, bee-eaters and starlings.
365. *Indicator maculatus* Spotted Honeyguide 607
 R LM? BR (3)
 The race *stictithorax* is a fairly common resident in the forests of Bengengai (Traylor & Archer 1982). There is one recent January sight record from Gilo (Nikolaus 1982). Probably parasitic on woodpeckers.
366. *Indicator minor* Lesser Honeyguide 608
 R? LM NBR
 Uncommon throughout in any type of woodland. Some apparently move north during the rains. Racial status and distribution are not clear, but *diadematus* probably occurs throughout except where perhaps replaced by *riggenbachii* along the southern border. Parasitic on small barbets, woodpeckers, swallows and kingfishers.
367. *Indicator variegatus* Scaly-throated Honeyguide 606
 R NBR
 A fairly common resident of forests and secondary growth at Imatong, Dongotona and Didinga up to 3000 m. Parasitic on woodpeckers.
368. *Prodotiscus insignis* Cassin's Honeybird 613
 The nominate race is an uncommon breeding visitor to the Imatongs, from January to May, frequenting secondary growth and forest at about 2000 m. Parasitic on white-eyes and probably small warblers.

Family PICIDAE Woodpeckers, Wrynecks

Sub-family Jynginae

369. *Jynx ruficollis* Red-throated Wryneck 635
 R? NBR
 Rare: the nominate race has been recorded once from Ibba River (Woodman 1936).
370. *Jynx torquilla* Wryneck 634
 PMW
 The nominate race is an uncommon winter visitor in open woodland and dry *Acacia* savanna from November to March. Recorded from Kapoeta to Yei, Lado, Tali Post and Meshra er Req.

Sub-family Picinae

371. *Campethera abingoni* Golden-tailed Woodpecker 622
 R? NBR
 The race *chrysura* is uncommon in forest edge and well wooded savanna along the Zaire border at Maridi, Yambio and near Tambura.
372. *Campethera caroli* Brown-eared Woodpecker 615
 R BR (3)
 The race *budongoensis* is a fairly common resident of forest at low altitude, recorded from Imatong, the Aloma Plateau and Bengengai.
373. *Campethera nivosa* Buff-spotted Woodpecker 616
 R NBR
 The race *herberti* occurs in forest at Bengengai where it is apparently rare.
374. *Campethera nubica* Nubian Woodpecker 619
 R BR (3, 4)
 The nominate race is a fairly common resident in any kind of woodland, mainly east of the Nile, where it replaces the Fine-spotted Woodpecker. West of the Nile the only records are from Lado to Yei and the Aloma Plateau.
375. *Campethera cailliautii* Little Spotted (Green-backed) Woodpecker 614
 R NBR
 The race *permista* is an uncommon resident in forest and secondary growth in Zande where recorded from Zakure Forest and Bengengai.
376. *Campethera punctuligera* Fine-spotted Woodpecker 621
 R BR (3, 4)
 The race *balia* is a fairly common resident in better wooded savanna west of the Nile, where recorded from Maridi, Tambura, Rumbek, Wau to the Boro River and Kafia Kingi. A record from Torit needs confirmation.
377. *Dendropicos fuscescens* Cardinal Woodpecker 623
 R BR (12)
 Locally common along the eastern and southern border. Two races occur: *lepidus* in better wooded savanna and forest clearings up to 2200 m along the southern border from Tambura to Imatong and Didinga, and on the Ethiopian border north of Boma; *hemprichii* in the dry Acacia savanna east of Kapoeta.
378. *Dendropicos poecilolaemus* Uganda Spotted Woodpecker 626
 R NBR
 Little recorded, but possibly not uncommon in open wooded savanna and cultivation. So far only recorded from Kajo Kaji, Maridi and Numatina River.
379. *Mesopicos goertae* Grey Woodpecker 630
 R NBR
 A fairly common resident of woodland, cultivation and forest edge on Imatong and Didinga up to 2200 m. The race *centralis* occurs throughout except along the Ethiopian border where it is replaced by *spodocephalus*.

380. *Mesopicos xantholophus* Yellow-crested Woodpecker 632
 R NBR
 Apparently rare in forests along the Zaire border where recorded only from the Aloma Plateau and Bengengai.
381. *Picoides obsoletus* Brown-backed Woodpecker 627
 R BR (3)
 The nominate race is fairly common in dry open wooded savanna throughout.
382. *Thripias namaquus* Bearded Woodpecker 629
 R BR (3)
 The race *schoensis* is rather uncommon and local in well wooded savanna. Recorded from Raga, Gogrial, Tonj, the Imatong foothills and Boya Hills.

Family ALAUDIDAE Larks

383. *Calandrella brachydactyla* Short-toed Lark 684
 PMW
 Locally common in small parties in the arid savanna north of Malakal from late October to early March. Three Palaearctic races could occur: nominate, *hermonensis* and *longipennis*.
384. *Eremopterix leucotis* Chestnut-backed Sparrow Lark 679
 R? LM? BR (1, 11, 12)
 Fairly common on short grassland in the dry season, from September to March, especially after recent burning. Found along the Nile north of Juba in the Sudd and around Tonj and Wau. Probably moves north during the rainy season. The nominate race occurs throughout most of the range, but is probably replaced by *melanocephala* north of Kodok.
385. *Eremopterix nigriceps* White-fronted Sparrow Lark 680
 R NBR
 The race *albifrons* is locally common in the extreme north around Renk on arid short grass plains, showing a preference for sandy ground.
386. *Eremopterix signata* Chestnut-headed Sparrow Lark 681
 R LM? NBR
 The race *harrisoni* is a common resident in the extreme south east, frequenting open stony savanna.
- Galerida cristata* Crested Lark 675
 In sandy and stony semi-desert country the following races could occur: *somaliensis* near Lake Turkana and *isabellina* around Renk.
387. *Galerida modesta* Sun Lark 677
 R LM BR (7)
 Fairly common south-west of Torit - Rumbek - Aweil favouring open ironstone clearings with short grass in woodland and also around Malakal. Mainly resident, but with local movements in the dry season. The race involved is *bucoloca*, except along the Zaire border where the nominate race occurs.

388. *Mirafraga albicauda* Northern White-tailed Bush Lark 655
 R? NBR
 Rather local on black cotton soil in the north around Kaka and Kodok, and along the Sobat river.
389. *Mirafraga cantillans* Singing Bush Lark 654
 R? LM? NBR
 The race *marginata* is locally common in short grassland and open *Acacia* savanna, from Juba and Lado east to the Ethiopian border. The race *chadensis* may occur around Renk.
390. *Mirafraga cordofanica* Kordofan Bush Lark 656
 R? NBR
 Rare with only one record, from Lake No (H & M 1970). Probably occurs elsewhere along the northern border in red sandy soil areas.
391. *Mirafraga erythropygia* Red-tailed Bush Lark 657
 LM BR (1, 3)
 An uncommon and little recorded breeding visitor from December to April, often noticed on recently burnt grassland. Recorded from Boma, Mongalla, Kajo Kaji, Malakal, Wau and Riki River. It moves to open bush savanna in the North Sudan during the rainy season.
392. *Mirafraga hypermetra* Red-winged Bush Lark 657
 R NBR
 Locally common in bushed grassland in the south east. The race *kidepoensis* occurs along the Uganda border from Rejaf to Lake Turkana. The race *kathangorensis* is known only from the area around Jebel Kathangor but might occur east to the Ethiopian border.
393. *Mirafraga poecilosterna* Pink-breasted Lark 665
 R? LM? NBR
 Uncommon in *Acacia* short grass savanna in the extreme southeast, east of Kapoeta. There seem to be some local movements for recent records are confined to May/June.
394. *Mirafraga rufocinnamomea* Flappet Lark 660
 R BR (6 - 8)
 Fairly common throughout in tall grass savanna and open woodland. Five races occur, but their exact ranges need clarification: *sobatensis* from the Baro/Sobat River across to Lake No, and possibly *omoensis* around Boma; *torrida* along the Uganda border west to Kajo Kaji; *tigrina* west of it, mainly along the Zaire border, possibly up to Wau. It is not clear whether birds around Boro/Riki River also belong to *tigrina* or whether they mark the southern limit of *furensis*.

Family HIRUNDINIDAE Swallows, Martins, Rough-wings

395. *Delichon urbica* House Martin 1074
 PMP PMW
 The nominate race is a rather uncommon Palaearctic passage migrant throughout in late September/October and April/early May but a regular and common winter visitor at higher altitudes of the Imatong, Dongotona and Didinga Mts.

396. *Hirundo abyssinica* Striped Swallow 1065
 R? LM BR(4, 11, 12)
 Rather local in better wooded savanna and forest clearings. The race *bannermanni* might be resident in the extreme northwest from the Boro and Riki Rivers south to Wau; the nominate race possibly breeds in April around Boma; *unitatis* breeds in the Kajo Kaji - Didinga region in November/December and migrates, probably southwards, in April/May when there is a regular passage through Gilo.
397. *Hirundo aethiopica* Ethiopian Swallow 1057
 R? LM BR (4 - 6)
 The nominate race is fairly common in the Juba - Lado - Maridi region, mainly from August to March, and north of Wau - Aweil - Malakal, mainly from May to September, frequenting rivers and towns. More information is needed about status and movements.
398. *Hirundo daurica* Red-rumped Swallow 1062
 PM? R? LM BR(1, 6)
 The race *domicella* is fairly common and probably resident along small rivers at low altitude throughout, breeding under bridges. The race *emini* occurs in the Imatong, Dongotona and Didinga Mts. from September to February apparently as a non-breeding visitor. The Palaearctic race *rufula* could occur.
399. *Hirundo fuligula* African Rock Martin 1073
 R BR (7, 8)
 The race *fusciventris* is an uncommon resident in suitable rocky localities at Kajo Kaji, Imatong, Dongotona, Didinga and Boma.
400. *Hirundo griseopyga* Grey-rumped Swallow 1066
 LM? BR (1)
 The nominate race is a very local breeding visitor from December to April to small rivers in better wooded savanna, reported from Boro River, Wau, Rumbek and Amadi.
401. *Hirundo rustica* Swallow 1054
 PMP PMW
 The nominate race is a common passage migrant and winter visitor throughout, but never in very large numbers. It occurs from late August to April. Passage is noticed from September to November and in April.
402. *Hirundo semirufa* Rufous-chested Swallow 1064
 LM NBR
 The race *gordoni* is locally uncommon in the Kajo Kaji - Mongalla - Imatong region from November to April, probably as a non-breeding visitor. Birds at Boro River in late March and April could be on passage to breeding grounds in Darfur.
403. *Hirundo senegalensis* Mosque Swallow 1063
 R LM BR (1, 2, 12)
 The race *saturator* is a fairly common resident on the Imatong, Dongotona and Didinga Mts. up to 2000m, breeding from December to February. The species also occurs at rivers and lakes elsewhere, mainly from January to April and in July and August; these records could refer to *saturator* or to the nominate race, which breeds in Darfur from June to September.

404. *Hirundo smithii* Wire-tailed Swallow 1061
 R? BR (2 - 4, 11)
 The nominate race is rather uncommon and local in the southwest where it breeds under bridges. Recorded from Kajo Kaji, Lado, Maridi, Tambura, Aweil, Raga and Boro River.
405. *Psalidoprocne albiceps* White-headed Roughwing 1080
 R LM? BR (8, 9)
 The nominate race is a common resident of forest and forest edge on Imatong, Dongotona and Didinga up to 3000 m. Less common around Kajo Kaji and on the Aloma Plateau.
406. *Psalidoprocne pristopectera* Black Roughwing 1076
 R? LM NBR
 Rather uncommon along forest edges and in better wooded savanna. White underwinged birds of the race *obaginea* occur along the Zaire border from the Aloma Plateau to Tambura. Birds around Boro/Riki River in April which appeared to have grey underwings were probably of the race *chalybea*.
407. *Riparia cincta* Banded Martin 1070
 LM? AM? NBR
 Small numbers occur in the Juba area between May and July; these could involve the race *suahelica* and/or *erlangeri*.
408. *Riparia paludicola* African Sand Martin 1069
 LM AM? NBR
 An uncommon non-breeding visitor, mainly along the southern border from December to February. Records undoubtedly involve the race *dubius*, but *minor* from North Sudan could also occur.
409. *Riparia riparia* Sand Martin 1068
 PMP PMW
 The nominate race is probably the most numerous Palaearctic winter visitor to the Sudd region, arriving from late August and leaving in late April, favouring open waters and open swampy grassland. Uncommon elsewhere, and occurring mainly on passage during September/October and late March/April.

Family DICRURIDAE Drongos

410. *Dicrurus adsimilis* Drongo 1088
 R BR (6, 11)
 The race *divaricata* is a common resident of woodland at low altitude throughout; *coracinus* is a forest bird confined to the Zaire border.
411. *Dicruris ludwigii* Square-tailed Drongo 1089
 R NBR
 The race *sharpei* is rather uncommon, confined to forest thicket up to 2000 m on the southern slopes of the Imatongs.

Family ORIOLIDAE Orioles

412. *Oriolus auratus* African Golden Oriole 1165
 LM? AM NBR
 The nominate race is a common dry season visitor throughout to better wooded savanna. Birds move north from May to September.

413. *Oriolus larvatus* Black-headed Oriole 1167
 R? NBR
 The race *rolleti* is fairly common in forests, better wooded savanna and dry woodland southeast of Juba, Bor and Boma.
414. *Oriolus nigripennis* Black-winged Oriole 1169
 R NBR
 Typically a bird of lowland forests but common on Imatong up to 2000 m, also recorded from Zande and Bengengai.
415. *Oriolus oriolus* Golden Oriole 1164
 PMP
 The nominate race is a common Palaearctic passage migrant in North Sudan from September to November. In the South, recorded only along the southern border and the Nile. Frequenting better wooded savanna and large trees in gardens, but only during March, suggesting that they overfly on autumn passage.

Family CORVIDAE Crows, Piapiac

416. *Corvus albus* Pied Crow 1172
 R LM? NBR
 Widespread in the dry season in any open country. Most common and probably resident around towns, though breeding is not yet recorded. Rare in the forest region along the Zaire border and in the arid extreme southeast.
417. *Corvus capensis* Cape Rook 1173
 R LM BR (1, 2, 12)
 Locally common in the dry season in open grass plains with scattered trees, mainly east of the Nile north to Melut. Most birds seem to move to the drier southeast in the rainy season.
418. *Corvus rhipidurus* Fan-tailed Raven 1177
 R? NBR
 A typical resident of rocky hills southeast of Juba - Boma; rare further north along the Ethiopian border.
419. *Corvus ruficollis* Brown-necked Raven 1171
 R? NBR
 The race *edithae* is an uncommon resident in the arid plains east of Kapoeta. An apparent hybrid between the Pied Crow and this species has been seen at Jebel Kasangor.
420. *Ptilostomus afer* Piapiac 1179
 R LM? NBR
 Common in small parties in open country throughout, but absent along the Ethiopian border. Often together with cattle and around villages where they favour *Borassus* palms for roosting.

Family PARIDAE Tits

421. *Parus albiventris* White-bellied Tit 1155
 R BR (1)
 A common resident of the Imatong, Dongotona and Didinga Mts. above 1600 m, frequenting forest edges and secondary growth.

422. *Parus funereus* Dusky Tit 1157
 R BR (3)
 A rather uncommon resident in lowland forests on Imatong, sometimes occurring in forest edge and secondary growth up to 2000 m at Gilo.
423. *Parus leucomelas* Black Tit 1154
 R NBR
 The race *guineensis* is a fairly common resident of open bushed savanna with scattered trees, less common east of the Nile. The nominate race could occur north of the Sobat River.

Family REMIZIDAE Penduline Tits

424. *Remiz musculus* Mouse-coloured Penduline Tit 1161
 R BR (12)
 An uncommon resident in semi-arid *Acacia* short grass savanna in the southeast from Lalo to the Kenya border.
425. *Remiz parvulus* Yellow Penduline Tit 1163
 R NBR
 The nominate race is an uncommon resident of better wooded savanna west of the Nile.
426. *Remiz punctifrons* Sennar Penduline Tit 1162
 R NBR
 Uncommon in dry woodland, replacing the Yellow Penduline Tit in the drier habitat north of Malakal.

Family SALPORNITHIDAE Spotted Creeper

427. *Salpornis spilonota* Spotted Creeper 1283
 R NBR
 The race *emini* is a rare and local resident of open woodland from Kajo Kaji to Tambura.

Family TIMALIIDAE Babblers

428. *Alcippe abyssinica* African Hill Babbler 740
 R BR (1, 11, 12)
 The nominate race is a common resident of thick forest undergrowth and forest edge above 1800 m on the Imatong, Dongotona and Didinga Mts.
429. *Ptyrticus turdinus* Thrush Babbler 734
 R NBR
 The nominate race is rather uncommon, frequenting the undergrowth of gallery and bowl forests from Yei to Bengengai.
430. *Trichastoma albipectus* Scaly-breasted Illadopsis 737
 R BR (8, 10)
 A rather uncommon species of lowland forest undergrowth, known from Imatong and Bengengai.
431. *Trichastoma fulvescens* Brown Illadopsis 735
 R BR (8)
 The race *ugandae* is a common resident of the undergrowth of lowland forest from Imatong to Bengengai.

432. *Trichastoma puveli* Puvel's Illadopsis W 667
 R BR (3)
 Only recorded from Bengengai where it is perhaps a fairly common resident of forest undergrowth. (Traylor & Archer 1982).
(Trichastoma rufipennis Pale-breasted Illadopsis) 736
 This species must be removed from the Sudan list; the skins listed from Imatong by Cave and Macdonald have been re-identified as immature Scaly-breasted Illadopsis.
433. *Turdoides leucopygius* White-rumped Babbler 728
 R NBR
 The race *omoensis* is fairly common on the Boma Hills, frequenting thick bushes and woodland at higher elevation.
434. *Turdoides plebejus* Brown Babbler 723
 R BR (1. 11. 12)
 The race *cinereus* is fairly common in any broadleaved tree savanna southwest of Didinga - Lado - Rumbek - Aweil. The nominate race could occur north of Kodok.
435. *Turdoides rubiginosus* Rufous Chatterer 732
 R BR (1, 2, 8, 9, 12)
 The nominate race is common in dry savanna where it moves around in noisy parties. Recorded north to Jongolei and Boma and west to Aweil.
436. *Turdoides tenebrosus* Dusky Babbler 727
 R NBR
 Uncommon to rare, frequenting well-wooded hilly districts. Recorded from Boma, Jebel Lado, Kajo Kaji, Mt. Baginzi, Li Rangu and Boro River.

Family CAMPEPHAGIDAE Cuckoo Shrikes

437. *Campephaga flava* Black Cuckoo Shrike 1081
 LM NBR
 Uncommon and silent, frequenting forests, gallery forests and better woodland along the southern border, mainly at higher altitudes in January/February and September/October. Movements need clarification
438. *Campephaga phoenicea* Red-shouldered Cuckoo Shrike 1083
 R? LM BR (6)
 Fairly common from December to April, but otherwise uncommon frequenting better wooded savanna, mainly along the southern border but with records north to Wau and the Sobat River. More information is needed about movements and relationship to the Black Cuckoo Shrike
439. *Campephaga quiscalina* Purple-throated Cuckoo Shrike 1084
 R? BR (4, 5)
 The race *martini* is fairly common and probably resident on Imatong in forest and secondary growth from 1600 to 2600m. Reported also from the Aloma Plateau and less certainly from Didinga Mts.
440. *Coracina caesia* Grey Cuckoo Shrike 1068
 R? BR (8)
 The race *pura* is very common in the dry season on the Imatong, Dongotona and Didinga Mts, inhabiting forest, forest edge and cultivation from 1600 to 2500m.

441. *Coracina pectoralis* White-breasted Cuckoo Shrike 1085
R? LM BR (2, 3)
Fairly common in better wooded savanna with tall trees southeast of Didinga - Rumbek - Wedweil, and also on the Boma Plateau. Possibly only a breeding migrant; recorded January - March and once September.
- Family PYCNONOTIDAE Bulbuls
442. *Andropadus curvirostris* Cameroon Sombre Greenbul 774
R BR (1, 8, 10)
The nominate race is a common forest resident up to 1600 m on Imatong and in the gallery forests in Zande. Occasionally wanders to 2000 m at Gilo in November/December.
443. *Andropadus gracilirostris* Slender-billed Greenbul 771
R BR (8)
The race *congensis* is a rather uncommon bird of forest canopy, occurring from Imatong to the Aloma Plateau and Zande up to 1600 m.
444. *Andropadus latirostris* Yellow-whiskered Greenbul 776
R BR (1, 3, 5 - 7)
A common resident of forests up to 2700 m from Didinga to Zande, but not recorded from Aza Forest. There is a distinct movement on Imatong where birds descend to lower altitude in April/May after breeding. Lowland birds west of the Nile breed in the rainy season.
445. *Andropadus virens* Little Greenbul 775
R BR (9, 10)
The race *holochlorus* is a common resident of bowl and gallery forests west of the Nile at Aloma Plateau, Bengengai and Zande.
446. *Baeopogon indicator* Honeyguide Greenbul 751
R NBR
The nominate race is a fairly common resident of lowland and gallery forests along the southern border, reported from Imatong, the Aloma Plateau, Aza Forest and Bengengai. Usually confined to high tree tops.
447. *Bleda eximia* Green-tailed Bristlebill 747
R NBR
The race *ugandae* is fairly common, frequenting thick undergrowth in gallery and bowl forests at Bengengai.
448. *Bleda syndactyla* Bristle-bill 746
R BR (1, 3, 6, 8, 10)
The race *woosnami* is a common resident in lowland forest undergrowth on Imatong, the Aloma Plateau and at Bengengai.
449. *Chlorocichla flavicollis* Yellow-throated Leaflove 749
R BR (9)
The race *soror* is a fairly common resident of green woodland along the Zaire border from Yei to Zande, showing a preference for Mango trees.
50. *Chlorocichla laetissima* Joyful Greenbul 770
R BR (3, 10)
A fairly common though rather local resident of forest, forest edge and secondary growth on Imatong from 1000 - 2000 m.

451. *Chlorocichla simplex* Simple Greenbul W 694
R? NBR
A secretive species of dense lowland forest known only from Bengengai (Traylor & Archer 1982).
452. *Criniger calurus* Red-tailed Greenbul 745
R NBR
The race *emini* is known only from Bengengai (Traylor & Archer 1982), but is probably an uncommon resident of thick forest undergrowth elsewhere in Zande.
453. *Nicator chloris* Nicator 1148
R BR (7, 8)
The nominate race is a common resident of lowland bowl and gallery forests, recorded from Imatong, the Aloma Plateau, Yambio and Bengengai.
454. *Phyllastrephus albigularis* White-throated Greenbul 761
R BR (6 - 8)
A common resident of lowland forest undergrowth up to about 1200 m, at Imatong, Aza Forest and Bengengai. Not yet recorded from the Aloma Plateau.
455. *Phyllastrephus baumanni* Toro Olive Greenbul 760
R BR (5)
The race *hypochloris* appears to be a fairly common forest resident up to 1600 m on the southern side of Imatong and on the Aloma Plateau.
456. *Phyllastrephus placidus* Olive Mountain Greenbul 758
(treated as separate from *P. fischeri*)
R BR (1 - 3, 12)
The race *sucosus* is a common resident of forested ravines and secondary growth from 1400 to 2000 m on the Imatong, Dongotona and Didinga Mts.
457. *Phyllastrephus scandens* Leaflove 750
R BR (10)
The race *orientalis* is a fairly uncommon resident in better woodland and gallery forests along the Zaire border from the Aloma Plateau and Yei to Zande.
458. *Phyllastrephus strepitans* Northern Brownbul 754
R? BR (1, 2)
Uncommon, probably resident, frequenting thickets in wooded savanna and arid scrub country in the southeast from Juba and Lado to the Ethiopian border.
459. *Pycnonotus barbatus* Common Bulbul 744
R BR (4 - 6)
The race *shoanus* is a common resident in any type of habitat along the Ethiopian border. The other white-vented race *arsinoe* reaches South Sudan around Renk. Between Malakal and Renk and on the Jebel Marra the yellow and the white-vented forms occur together without interbreeding.
460. *Pycnonotus tricolor* Yellow-vented Bulbul 742
R BR (1, 4, 8 - 10)
A common resident in most types of habitat, although uncommon in the

Sudd, and replaced by the White-vented Bulbul along the Ethiopian border. North of Malakal both species are found together.

Family TURDIDAE Thrushes, Robins etc.

461. *Alethe diademata* Fire-crested Alethe 901
 R NBR
 The race *woosnami* is fairly common in gallery forests and forest edge at Bengengai.
462. *Alethe poliocephala* Brown-chested Alethe 903
 R BR (1 - 3)
 The race *giloensis* is a common resident in Imatong up to 2500 m. Lowland birds move to higher altitude in the dry season.
463. *Cercomela familiaris* Red-tailed Chat 871
 R BR (1)
 An uncommon and very local resident on rocky hills with open woodland. The nominate race is recorded from Boro River, from Wau to Tambura and from Boya Hills, while Boma birds are assigned to *omoensis*.
Cercomela scotocerea Brown-tailed Rock Chat 870
 The race *turkana* should occur in the arid rocky bushland of the Ilemi Triangle.
464. *Cercotrichas galactotes* Rufous Bush Chat 934
 PM LM? NBR
 The nominate race is an uncommon winter visitor to dry *Acacia* shrub and open woodland as far south as Malakal.
465. *Cercotrichas leucophrys* White-browed Scrub Robin 910/911
 R? LM NBR (12)
 The white-winged race *leucoptera* is a breeding resident in dry *Acacia* shrub country in the southeast. In the dry season some wander west to Lado where they meet the red-backed race *zambesiana*. The latter is recorded from Lado (Sept.), Raffile (Sept.), Yei (Nov.), and Kajo Kaji.
Cercotrichas podobe Black Bush Robin 916
 Should occur in dry *Acacia* shrub around Renk.
466. *Cichladusa guttata* Spotted Morning Thrush 909
 R BR (4)
 The nominate race is a common resident in thick dry bushland along the Nile, east of it north to the Sobat River, and rarely west to Wau.
467. *Cossypha albicapilla* White-crowned Robin Chat 891
 R NBR
 The race *omoensis* is a fairly common resident in better wooded savanna and green bushes on Boma Hills.
468. *Cossypha caffra* Robin Chat 893
 R BR (1,3)
 The race *iolaema* occurs as a common resident of secondary growth and forest above 1600 m on Imatong, and more rarely on the Dongotona and Didinga Mts.

469. *Cossypha cyanocampter* Blue-shouldered Robin Chat 889
 R BR (10)
 The race *bartteloti* is a rather uncommon resident on Imatong frequenting thick damp forest undergrowth at Gilo, Upper Talanga and Lotti.
470. *Cossypha heuglini* White-browed Robin Chat 884
 R BR (3)
 The nominate race is a common resident in well wooded savanna south of the Sobat and Bahr-el-Arab Rivers.
471. *Cossypha natalensis* Red-capped Robin Chat 890
 LM? AM BR (1, 2, 3)
 The race *garguensis* is seasonally common in lowland forests from Imatong to Bengengai, arriving at the beginning of March, breeding in the rainy season and leaving about October. Birds from Didinga and Boma represent a different population which probably stays from November to May and breeds in January/February.
472. *Cossypha niveicapilla* Snowy-headed Robin Chat 892
 R? LM? BR (3, 7)
 A fairly common but local species of forest edge and dense vegetation near streams. Recorded southwest of Torit - Lado - Wau - Kafia Kingi.
473. *Cossypha semirufa* Rüppell's Robin Chat 885
 R? LM? NBR
 The nominate race has only once been recorded, in thick damp scrub at Boma in November (C & MacD 1955).
474. *Dryocichloides poliopterus* Grey-winged Ground Robin 887
 R? LM? NBR
 The nominate race is a fairly common but very local bird of thick forest undergrowth near water up to 1800m. Possibly only a non-breeding visitor to Imatong and the Aloma Plateau from July to October.
475. *Luscinia luscinia* Sprosser 992
 PMP
 A little noticed Palaearctic autumn passage migrant east of the Nile. Recorded only once, in dense *Acacia* thicket east of Torit in November (pers. obs.).
476. *Luscinia megarhynchos* Nightingale 921
 PM
 The nominate race is an uncommon Palaearctic visitor from November to early April, frequenting thickets in better wooded savanna and wintering south of Boro River - Tonj - Lado - Kapoeta.
Luscinia suecica Bluethroat 919
 Could occur in thickets near water along the northern boundary.
477. *Monticola rufocinerea* Little Rock Thrush 853
 LM? NBR
 The nominate race has been recorded from the Imatong and Didinga Mts. between November and March, from open slopes with scattered trees at about 2000m. There are no recent records.

478. *Monticola saxatilis* Rock Thrush 850
PMW PMP?
An uncommon Palaearctic winter visitor to rocky hills in open savanna from December to March, mainly south of Boma - Amadi - Wau - Boro River. Even though there is a marked autumn movement in North Sudan from September to November, passage is not noticed in the South.
- Monticola solitarius* Blue Rock Thrush 851
This Palaearctic species is to be expected in the arid rocky country round Renk.
479. *Myrmecocichla nigra* Sooty Chat 880
R? LM NBR
Uncommon in open bushed tree savanna from Wau to Tambura in August/September and from Maridi to Kajo Kaji and Opari in July, September, December and March/April.
480. *Oenanthe bottae* Red-breasted Wheatear 867
R? LM AM? NBR
The race *heuglini* is seasonally very common on open short grass plains, and very partial to burnt grass patches. Movements are not clear. All records are from east of the Nile. Occurrence in the east and northeast is from December to February and probably involves North Sudan breeders, but May/June records from the extreme southeast could involve a population moving south to Uganda.
481. *Oenanthe deserti* Desert Wheatear 857
PMW
The nominate race is a regular Palaearctic winter visitor in December/January to open arid country south to Renk.
482. *Oenanthe hispanica* Black-eared Wheatear 858
PMW
The race *melanoleuca* is a fairly common winter visitor to semi-arid open grassland and bushland south to Malakal from December to February; once recorded from Lado in December.
483. *Oenanthe isabellina* Isabelline Wheatear 855
PMW
A common Palaearctic winter visitor, mainly east of the Nile, where it tends to frequent drier open grassland habitat than the Northern Wheatear. No distinct passage is noticeable.
484. *Oenanthe oenanthe* Northern Wheatear 854
PMW PMP?
The most common Palaearctic wheatear, wintering throughout in suitable open grassland and bush habitat from September to the beginning of April (mainly November - March). Passage, though expected, is not noticeable. The nominate race and *libanotica* occur, the latter probably more restricted to the North Sudan.
485. *Oenanthe pleschanka* Pied Wheatear 859
PMW PMP?
The nominate race is an uncommon Palaearctic winter visitor to the semi-arid southeast in open rocky grassland. Recorded from Lado and Torit in December, Kapoeta in December and January and along the Kenya border in November, December and January.

486. *Pentholaea albifrons* White-fronted Black Chat 874
 R? LM NBR
 The race *clericalis* is an uncommon dry season visitor to open grassy woodland from November to March, recorded from Yei to Torit and from Lado to Kajo Kaji.
487. *Phoenicurus phoenicurus* Redstart 917
 PMW
 The nominate race is a common Palaearctic winter visitor throughout in any type of woodland, from October to early April. The white-winged *samamiscus* has been collected near Wau in January and at Yei in November.
488. *Pogonocichla stellata* White-starred Forest Robin 915
 R NBR
 The race *pallidiflava* is a common resident on Imatong, but uncommon on the Dongotona and Didinga Mts. It frequents forest above 2000 m.
489. *Saxicola rubetra* Whinchat 883
 PMW PMP
 A common Palaearctic passage migrant throughout from October to December and in March/early April in any open grassy habitat. Some regularly winter along the southern border south of 6°N.
490. *Saxicola torquata* Stonechat 882
 R * PMW? BR (4)
 The race *albofasciata* is a common resident on the Imatong, Dongotona and Didinga Mts., frequenting open grassland above 1800 m. Palaearctic winter visitors (*variegatus*) could occur in the extreme north.
491. *Sheppardia aequatorialis* Equatorial Akalat 898
 R BR (1, 4, 11)
 The race *acholiensis* is a common resident of primary forest undergrowth on Imatong above 1800 m, breeding in the dry season.
492. *Stiphrornis erythrothorax* Forest Robin 900
 R? NBR
 The race *xanthogaster* is recorded from Bengengai (Traylor & Archer 1982). It appears to be an uncommon resident of forest undergrowth.
493. *Stizorhina fraseri* Rufous Thrush 910
 R? NBR
 The race *vulpina* is an apparently uncommon species of gallery forest, where it perches on branches overhanging rivers. Recorded only from the Aloma Plateau and Bengengai (Cave & MacDonald 1955).
494. *Thamnolaea cinnamomeiventris* Cliff Chat 876, 878
 R NBR
 A rather uncommon and local species of rocky hills and cliffs. The race *subrufipennis* has been recorded from Boma, Jebel Kasangor, Jebel Lado and Kajo Kaji, while *coronatus* has its southern limits at Boro River.

495. *Turdus abyssinicus* Olive Thrush 841
 R BR (1 - 4)
 The nominate race is a common resident on Imatong, frequenting forest, forest edge and secondary growth above 1600 m. Also reported from the Dongotona and Didinga Mts.
496. *Turdus fischeri* Spotted Ground Thrush 846
 R? NBR
 A single example of a newly described race *maxis* was netted on Imatong in November 1979 in the Lotti primary lowland forest (Nikolaus 1982a).
497. *Turdus pelios* African Thrush 840
 R BR (1 - 3)
 A common resident in low altitude woodland throughout. Two races occur: *pelios* north of about Tambura - Shambe - Baro River and *centralis* south of this line.
498. *Turdus piaggiae* Abyssinian Ground Thrush 845
 R BR (7)
 An uncommon resident near damp forest streams at higher altitudes. The nominate race is recorded from Boma, and *hadii* from the Imatong and Dongotona Mts.

Family SYLVIIDAE Warblers

499. *Acrocephalus arundinaceus* Great Reed Warbler 943
 PMP
 A rare Palaearctic passage migrant; only twice recorded, both birds in early April at Gilo (pers. obs.). Probably overflies, at least in autumn, as it is fairly common in North Sudan during September/October.
500. *Acrocephalus baeticatus* African Reed Warbler 946
 R? LM? NBR
 The race *cinnamomeus* is rare but widely distributed in any kind of tall grass and reeds. Recorded from Wau (February), Juba (January), Lado and Bahr-el-Zeraf. These could be non-breeding visitors from Jebel Marra.
501. *Acrocephalus gracilirostris* Lesser Swamp Warbler 957
 R NBR
 The race *jacksoni* is uncommon in reeds and tall grass over water; sometimes also in papyrus. Recorded along the Nile from Malakal to Juba.
502. *Acrocephalus griseldis* Basra Reed Warbler 943
 PMP
 A rare Palaearctic autumn passage migrant, caught once in late August at Juba in tall reed along the Nile (Nikolaus 1979a).
- Acrocephalus palustris* Marsh Warbler 945
 Not recorded, but it should occur along the Ethiopian border as an autumn passage migrant.
503. *Acrocephalus rufescens* Greater Swamp Warbler 955
 R NBR
 The race *niloticus* is a common resident in the papyrus vegetation of the Sudd, less common in reeds over water. Reported from Maridi, Juba and near Torit.

504. *Acrocephalus schoenobaenus* Sedge Warbler 947
PMW PMP
There are very few records of this Palaearctic migrant, from reedbeds along the Nile at Lado and Juba (October, November, January, April) and from dense bush vegetation beside water at Gilo (January, March, September). It is perhaps more common in the reed and sedge vegetation of the Sudd than available records indicate.
505. *Acrocephalus scirpaceus* Reed Warbler 944
PMW PMP
Little recorded, but probably not uncommon as a Palaearctic winter visitor and autumn passage migrant to reeds, tall grass and *Acacia* bushland. All available records are from the Nile valley and east of it from October to March. Most are assignable to the race *fuscus*, but nominate birds also occur.
506. *Apalis cinerea* Grey Apalis 973
R BR (2, 7)
The nominate race is common on the Didinga, Dongotona and Imatong Mts., mainly on forest edges above 1800 m, favouring large *Acacias*.
507. *Apalis flavida* Yellow-breasted Apalis 979/980
R NBR
An uncommon species represented by two races: *malensis* in the dry stony *acacia* grassland of the extreme southeast, and *caniceps* in low bushy grassland at Tambura and Mbili near Wau.
508. *Apalis jacksoni* Black-throated Apalis 982
R NBR
The nominate race is uncommon on the Didinga and Imatong Mts. above 1900 m, frequenting the tops of large forest trees, with a marked preference for *Albizia*.
509. *Apalis pulchra* Black-collared Apalis 977
R BR (1, 3, 4, ?1 - 12)
The nominate race is very common on Imatong in dense thickets above 1800 m.
510. *Apalis rufogularis* Buff-throated Apalis 972
R? NBR
The race *nigricens* is apparently rare, and known only from the Aloma Plateau and Bengengai, where it frequents large forest trees.
511. *Bathmocercus rufus* Black-faced Rufous Warbler 1053
R BR (7)
The race *vulpinus* is fairly common on Imatong above 1800 m and on the Aloma Plateau, frequenting dense thickets and forest undergrowth bordering rivers.
512. *Bradypterus alfredi* Bamboo Warbler 951
R NBR
Uncommon and recorded only at Gilo in grassy secondary growth. Quite often noticed after a large fire had damaged most of the low vegetation in the area. (Nikolaus 1982).

513. *Bradypterus baboecala* Little Rush Warbler 948
R NBR
An apparently local and uncommon resident in reedbeds, papyrus and tall floating grass in the Sudd. The race *sudanensis* is described from Lake No; otherwise there is but a single record from Juba.
514. *Bradypterus cinnamomeus* Cinnamon Bracken Warbler 952
R BR (5)
The race *cavei* is a very common resident in dense forest shrubbery and thick secondary growth on the Imatong and Dongotona Mts. above 1800 m.
515. *Camaroptera brevicaudata* Grey-backed Camaroptera 1011
R BR (2, 7, 10, 11)
The race *abbessinica* is very common throughout in any type of thick bush vegetation. North of Renk it is replaced by the nominate race.
516. *Camaroptera chloronota* Olive-green Camaroptera 1010
R BR (2, 3)
The race *toroensis* is known only from Bengengai, where it frequents forest undergrowth and is possibly not uncommon (Traylor & Archer 1982).
Camaroptera superciliaris Yellow-browed Camaroptera 1012
A lowland forest species; a record from Kombo in the extreme southeast needs confirmation.
517. *Camaroptera simplex* Grey Wren Warbler 968
R? NBR
The nominate race is fairly common in the extreme southeast, frequenting thicker bushes in dry *Acacia* grassland.
518. *Chloropeta natalensis* Yellow Warbler 805
R BR (6)
The race *massaica* is a rare resident of tall grass savanna, often near water; recorded from the Imatong foothills west to Maridi.
519. *Chloropeta similis* Mountain Yellow Warbler 806
R BR (4, 10)
A common resident on Imatong in forest undergrowth, forest edge and secondary growth above 1800 m; rare on the Didinga and Dongatona Mts.
- CISTICOLAS : All the *Cisticola* species seem to perform local movements to some extent in response to seasonal changes in habitat. They all breed during the rainy season.
520. *Cisticola aberrans* Rock-loving *Cisticola* 1027
R NBR
The race *petrophila* is restricted to open rocky grassland at high altitude on the Imatong and Dongotona Mts. where it is fairly common.
521. *Cisticola ayresii* Wing-snapping *Cisticola* 1019
R BR (4 - 6)
The race *imatong* is a fairly common resident in the Imatong Mts. on meadows above 2000 m.
522. *Cisticola bodessa* Boran *Cisticola* 1024a
R NBR
Fairly common in tall grass savanna on the Boma Hills, where it is often conspicuous singing from tall trees. These birds are best assigned to the race *caffaensis*.

523. *Cisticola brachyptera* Siffling Cisticola 1037
R BR (7)
Rather uncommon in dense tall grassland with scattered trees. The nominate race is recorded from the southern border, Boro River, Chak-Chak, Wau and Tonga; birds from Boma are nearer to the race *sedlitzi*.
524. *Cisticola cantans* Singing Cisticola 1031
R? LM? BR (7)
The race *concolor* is fairly common but rather local in better wooded thick grassland, often near rivers. Distributed from Gogrial south to Tambura and east to Didinga. Found breeding at Yei where the race *belli* has also been recorded.
525. *Cisticola chiniana* Rattling Cisticola 1024
R BR (6, 9)
The race *fricki* is fairly common, frequenting *Acacia* grassland south-east of Boma-Bor-Yei. A record from Wau needs confirmation.
526. *Cisticola cinereola* Ashy Cisticola 1042
R? LM? BR (12)
Only once recorded, possibly as a breeding visitor to *Acacia* short grassland on the Kenya border (Nikolaus 1979).
527. *Cisticola erythrops* Red-faced Cisticola 1032
R? LM NBR
Locally common in better wooded or bushed thick grassland, often near rivers. The race *pyrhomitra* is reported from Boma, Maridi (February) and Gilo (July to April). The race *sylvia* is recorded from Tambura (February), Yambio (May), Maridi (January) and Tapari River (February). The status of the two races, and the relationship to the Singing Cisticola, where both species occur, is not clear.
528. *Cisticola eximia* Black-backed Cisticola 1021
The nominate race is recorded rather locally in short open grassland from Aweil, Ayod region, Boma Plains and Kajo Kaji.
529. *Cisticola galactotes* Winding Cisticola 1033
R BR (6 - 9)
The race *marginata* is very common along the Nile and in the Sudd, frequenting papyrus and reeds and always associated with water bound vegetation. Other records are from Tarangole (Koss River), Yambio (Sue River), Wau (Sue River), Raga (Lol River) and Bahr-el-Arab.
530. *Cisticola juncidis* Zitting Cisticola 1016
R? LM? BR (5 - 10, 12)
The race *uropygialis* is common in short grassland along the Nile and east of it, but uncommon to the west.
531. *Cisticola lateralis* Whistling Cisticola 1028
R? BR (7)
The race *antinorii* is fairly common in well wooded grassland and low altitude forest edge from Didinga to Benengai and the Boro-Riki River. The type was described from Jur River.
532. *Cisticola nana* Tiny Cisticola 104
R? NBR
Recorded only once, in dry thornbush savanna along the Kenya border (Macdonald & Cave 1948).

533. *Cisticola natalensis* Croaking Cisticola 1036
R? LM? NBR
Locally common in open tall grassland with scattered bushes. Three races occur: *argentea* along the Ethiopian border south of Boma and *strangei* west of about 33 E. North of Lake No and the Sobat the latter is replaced by the race *tonga*.
534. *Cisticola robusta* Stout Cisticola 1035
R? NBR
There is a single record of the race *nuchalis* from tall grass savanna at Shambe (Macdonald & Cave 1948). The status of this species needs clarification.
535. *Cisticola ruficeps* Red-pate Cisticola 1043
R BR (7 - 10)
A common species along the Nile, frequenting thornbush savanna and sometimes open wooded grassland; uncommon elsewhere. The race *mongalla* is replaced by *scotoptera* north of Lake No and the Sobat River.
536. *Cisticola troglodytes* Foxy Cisticola 1038
R BR (4, 7, 9)
Common in grass savanna, and often associated with ironstone country. The nominate race occurs throughout except along the Ethiopian border where it is replaced by *ferruginea*.
537. *Drymocichla incana* Red-winged Grey Warbler 994
R NBR
Rare, and only recorded from Kajo Kaji, Yei, Maridi, Lado, Wau and the Boro River, mainly in thick bushes fringing rivers in open grass savanna.
538. *Eminia lepida* Grey-capped Warbler 993
R BR (11)
Common on the Imatongs in dense damp undergrowth above 1600 m.
539. *Eremomela badiceps* Brown-crowned Eremomela 1007
R NBR
The nominate race (*latukae* rejected by Traylor 1982) is fairly common on the Imatongs, frequenting large lowland forest trees.
540. *Eremomela icteropygialis* Yellow-bellied Eremomela 1003
R NBR
Uncommon in dry savanna scrub. The race *griseoflava* occurs southeast of Juba - Lado, while *alexanderi* is reported from Renk and Wau.
541. *Eremomela pusilla* Green-backed Eremomela 1005
R BR (2, 3)
The race *canescens* occurs fairly commonly throughout in any type of savanna at low altitude
542. *Heliolais erythroptera* Red-winged Warbler 1050
R NBR
The race *jodoptera* is a very local bird of open bushed grassland recorded along the southern border from Li Rangu to Torit and from Wau.

- Hippolais icterina* Icterine Warbler 935
Not yet recorded, but expected as a Palaearctic migrant in open woodland.
- Hippolais languida* Upcher's Warbler 931
Possibly a rare Palaearctic autumn passage migrant, with a single record from Mongalla (Cave & Macdonald 1955) which could not be traced. Confirmation is needed.
543. *Hippolais pallida* Olivaceous Warbler 938
PMW PMP
The Palaearctic race *eleica* is a locally abundant winter visitor from August to March in *Acacia* savanna along the Nile and east of it, but uncommon to rare in the west. The nominate race from Egypt winters south to Malakal from October to February.
544. *Hylia prasina* Green Hylia 1281
R BR (1)
The nominate race is fairly common in thick lowland forest undergrowth on Imatong, the Aloma Plateau and at Bengengai
545. *Locustella luscinioides* Savi's Warbler 939
PMW
A seldom recorded but perhaps not uncommon Palaearctic winter visitor to the Sudd south to Juba, frequenting tall grass, reeds and bushes near water. The nominate race has been found, but *fusca* could also occur.
546. *Macrosphenus flavicans* Yellow Longbill 1014
R BR (3)
The race *hypochondriacus* has been collected at Bengengai (Traylor & Archer 1982). A species of dense forest undergrowth and creepers.
547. *Parisoma lugens* Brown Parisoma 939
R? LM? BR (7)
The race *jacksoni* is rare, recorded only from the Imatong and Didinga Mts., where it has been found in large *Acacia* trees in well wooded grassland up to 2000m.
548. *Phyllolais pulchella* Buff-bellied Warbler 995
R? NBR
Fairly common along the Nile and to the east, frequenting tall trees in *Acacia* savanna. Elsewhere reported only from Maridi and Gorgrial.
549. *Phylloscopus bonelli* Bonelli's Warbler 962
PMW
The race *orientalis* winters rarely south to Malakal in dry *Acacia* bush land.
550. *Phylloscopus collybita* Chiffchaff 960
PMW
The race *abietinus* is a fairly common Palaearctic winter visitor from December to March to the Imatong, Dongotona and Didinga Mts. and the Boma Hills, frequenting forest edges and bushes, mainly above 1800m.
551. *Phylloscopus sibilatrix* Wood Warbler 96
PMW PMP?
An apparently rare Palaearctic visitor from late September to early April to forest edges near the southern border. Recorded only from Bengengai and Imatong (M.A. Traylor *in litt.*).

552. *Phylloscopus trochilus* Willow Warbler 959
PMP PMW?
There are surprisingly few records of this Palaearctic migrant, all from Wau and the region between Juba and Lake Turkana between September and January. There is no evidence of wintering or of spring passage. Birds are referable to the race *acredula*.
553. *Phylloscopus umbrovirens* Brown Woodland Warbler 964
R BR (4, 6, 8)
A fairly common resident of highland forest and forest edge above 1800 m on the Imatong, Dongotona and Didinga Mts.
554. *Prinia leucopogon* White-chinned Prinia 1048
R BR (4, ?1 - 12)
The race *reichenowi* is a common resident on the Imatong in forest and secondary growth. Also recorded from Bengengai.
555. *Prinia somalica* Pale Prinia 1046
R BR (12)
The race *erlangeri* is a fairly common resident in the dry *Acacia* scrub country of the extreme southeast, east of Kapoeta.
556. *Prinia subflava* Tawny-flanked Prinia 1045
R BR (7 - 10)
The nominate race is common in any kind of savanna except north of Lake No and the Sobat River where it is replaced by *deserti* and in the arid extreme southeast where the species is replaced by *P. somalica*.
557. *Schoenicola platyura* Fan-tailed Warbler 969
R? NBR
A species of tall moist grassland. There are only four records of the race *alexinae*, from Gazelle River (type), Nagishot/Didinga, Lotti Forest and Gilo.
558. *Sphenoeacus mentalis* Moustached Warbler 1051
R? NBR
The nominate race is locally common in tall grass savanna southwest of Raga - Wau - Bor - Imatong and at Boma near the Ethiopian border.
559. *Spiloptila rufifrons* Red-fronted Warbler 987
R? BR (12)
Recorded only from the extreme southeast in dry *Acacia* scrub. Birds are probably referable to the race *smithii*.
560. *Sylvia atricapilla* Blackcap 926
PMW PMP?
The nominate race is a very common Palaearctic winter visitor to the Imatong, Dongotona and Didinga Mts., frequenting green bush vegetation and forest above 1600 m from October to April. Not recorded elsewhere even though it should occur on passage. Birds ringed at Imatong have been recovered in Sweden and Lebanon.
561. *Sylvia borin* Garden Warbler 925
PMP
A regular Palaearctic passage migrant from September to November in wooded savanna near Wau and in the Yei - Lado - Didinga - Imatong area. On Imatong it occurs in secondary vegetation around Gilo from September to December and also from late February to the beginning of May.

562. *Sylvia communis* Whitethroat 92
PMW PMP
A rather uncommon Palaearctic winter visitor and passage migrant to dry woodland, favouring thorny vegetation. Recorded from late August to early April. Wintering birds are all of the nominate race, while many passage birds are *icterops* or *volgensis*.
563. *Sylvia curruca* Lesser Whitethroat 92
PMW
The nominate race winters commonly from late September to late February/early March in thornbush vegetation north of 10°N; occasionally recorded south to Juba.
564. *Sylvia hortensis* Orphean Warbler 92
PMW
The race *crassirostris* is a rare Palaearctic visitor to *Acacia* thickets south to Melut between September and February. The main wintering area is in North Sudan.
565. *Sylvia nisoria* Barred Warbler 93
?PMP
Four very fat birds were netted in mid February in *Acacia* woodland near Melut. The only other record is of a bird in late October near Wau (M. Rae). This species is a common passage migrant in North Sudan.
566. *Sylvietta brachyura* Crombec 99
R BR (1, 2, 4, 5, 6, 9, 10, 12)
Common throughout in any kind of savanna. The nominate race occurs at low altitude throughout except in the arid southeast where it is replaced by *leucopsis*. Birds ranging to 2400m on Imatong would be *dilutior* if this race proves to be valid.
567. *Sylvietta virens* Green Crombec 100
R BR (10)
The race *baraka* is a rather uncommon resident of lowland forest thickets, from Bengengai and Lotti Forest/Imatong.
568. *Sylvietta whytii* Red-faced Crombec 99
R BR (2, 12)
The race *abayensis* is rather uncommon and confined to the extreme southeast where it frequents better bush vegetation in acacia short grass savanna.

Family MUSCICAPIDAE Flycatchers

Sub-family Muscicapinae

569. *Artomyias fuliginosa* Sooty Flycatcher 80
R? NBR
Recorded only twice, from the Aloma Plateau (Macdonald & Cave 1948).
570. *Bradornis microhynchus* Grey Flycatcher 793/79
R? LM BR (2, 12)
The race *neumanni* is uncommon in dry bush savanna west to Juba and Lado and north to Jebel Kasangor. All records are from the dry season.

571. *Bradornis pallidus* Pale Flycatcher 792
R? LM? BR (1)
The nominate race is uncommon but occurs, perhaps as a non-breeding dry season visitor, in the open wooded savanna west of Malakal - Ayod - Juba. The only rainy season records are from the Zaire border region. Two other races occur: *parvus* on the Boma Hills and *murinus* on the Didinga Mts. above 1500 m.
572. *Empidonis semipartitus* Silverbird 801
R NBR
A common resident in Acacia savanna and semi-arid bushland northeast of a line from Kajo Kaji to Aweil.
573. *Ficedula albicollis* Collared Flycatcher 779/780
PMP PMW?
A rare and little recorded Palaearctic visitor to better wooded savanna. The race *semitorquata* has been collected at Kajo Kaji (February), Taufika (April) and recently by Traylor at Gilo and Bengengai (March).
574. *Melaenornis chocolatina* White-eyed Slaty Flycatcher 796
R BR (2, 3)
The race *fischeri* is a common resident on the Imatong, Dongtona and Didinga Mts., frequenting forest edge and secondary growth above 1600 m.
575. *Melaenornis edoliodes* Black Flycatcher 798
R BR (4)
The race *lugubris* is a fairly common resident southwest of Boro River - Gogrial - Lado - Didinga Mts., where it inhabits well wooded Savanna and gardens. Also recorded from Boma
576. *Muscicapa adusta* Dusky Flycatcher 781
R BR (1, 2)
The race *pumila* is a very common resident on the Imatong, Dongotona and Didinga Mts., frequenting open highland forest and secondary growth above 1800 m.
577. *Muscicapa aquatica* Swamp Flycatcher 782
R BR (4)
A species of reed, bush and papyrus vegetation near swamps and rivers, where it is locally common. The race *infulata* occurs along the Nile south of Malakal, while the West African nominate race is recorded from Boro River, Wau and Raffile.
578. *Muscicapa caerulescens* Ashy Flycatcher 785
R NBR
The race *brevicaudata* is an uncommon resident in the foothill forests of Imatong. Also recorded from Zande.
579. *Muscicapa comitata* Dusky Blue Flycatcher 803
R? NBR
The race *stuhlmanni* has been recorded only once, from gallery forest on the Aloma Plateau (Macdonald & Cave 1948).
580. *Muscicapa gambagae* Gambaga Flycatcher 778
R? LM? NBR
Recorded from open savanna at Lado in January (Nikolaus), Mvolo in March (B.M.).

581. *Muscicapa striata* Spotted Flycatcher 778
PMP
A common Palaearctic passage migrant from mid-October to early December in open woodland or wooded savanna. A few pass northwards between mid-March and mid-April. Two races occur: the nominate race passes mainly west of the Nile while *neumanni* is predominant east of it.
582. *Myioparus plumbeus* Lead-coloured Flycatcher 788
R BR (12)
The nominate race is a fairly common savanna resident southwest of Gogrial - Bor - Didinga.
- Sub-family Platysteirinae
583. *Batis minor* Black-headed Batis 820
R NBR
The race *erlangeri* is fairly common throughout, especially in well-wooded savanna and large *Acacia* trees along the Nile.
584. *Batis molitor* Chinspot Batis 817
R NBR
Apparently confined to the Didinga Mts., where it is a fairly common resident of secondary growth and well wooded grassland.
585. *Batis orientalis* Grey-headed Batis 818
R NBR
Fairly common in open wooded grassland. Two races occur: the nominate along the Ethiopian border with records confined to the Torit - Imatong - Didinga area, and *chadensis* west of the Nile, extending eastwards to Lado and Renk.
586. *Batis perkeo* Pygmy Batis 819
R? NBR
Recorded only from the Natoporuputh Hills where it is not uncommon in dry *Acacia* scrub (Nikolaus 1973).
587. *Bias musicus* Black and White Flycatcher 819
R? NBR
The nominate race is known only from Bengengai where it is a rare forest species (Macdonald & Cave 1948; Traylor).
588. *Hyliota flavigaster* Yellow-bellied Flycatcher 820
R NBR
The nominate race is a fairly common resident, frequenting tree tops in better wooded savanna southwest of Gogrial - Yei - Imatong and around Boma.
589. *Megabyas flammulata* Shrike Flycatcher 821
R? NBR
The race *aequatorialis* is known only from Bengengai, where it appears to be an uncommon resident in both primary and secondary forests.
590. *Platysteira blissetti* Jameson's Wattle-eye 822
R BR (1)
The race *jamesoni* is a rare resident of dense vegetation in bowl forests at Lotti (Imatong), the Aloma Plateau and Bengengai.

591. *Platysteira castanea* Chestnut Wattle-eye 824
R BR (6, 8)
The nominate race is known as a common resident in thick gallery forest at Bengengai and in lowland forest at Imatong.
592. *Platysteira cyanea* Wattle-eye 822
R BR (1, 2, 4, 7, 12)
The race *nyanzae* is a fairly common resident southwest of Didinga - Lado - Wau - Boro River, frequenting gallery forests and forest clearings up to 2000 m. The race *aethiopica* occurs on the Boma Hills.
- Sub-family Monarchinae
593. *Erranornis longicauda* Blue Flycatcher 827
R BR (3)
The race *teresita* is a common but local resident of gallery forest and forest edges up to 2000 m, recorded southwest of Imatong - Rumbek - Boro River.
594. *Terpsiphone viridis* Paradise Flycatcher 832
R? LM AM NBR
The race *ferreti* (incl. *speciosa*) is common throughout in any woodland and forest up to 2200 m. Status needs further clarification: in the forests along the southern border there are records for most months, but elsewhere it occurs from September to April, probably migrating to breed in the North Sudan during the rainy season. The southern African migratory race *plumbeiceps* was once collected in July on the Aloma Plateau (Nikolaus 1982).
595. *Trochocercus nigromitratus* Dusky Crested Flycatcher 831
R? NBR
Recorded only from Bengengai in March (Traylor & Archer 1982).
A species of forest undergrowth and secondary bush.
596. *Trochocercus nitens* Blue-headed Crested Flycatcher 830
R? NBR?
The nominate race is known only from Bengengai where a single bird was collected in March (Traylor & Archer 1982). A bird of forest undergrowth.
- Family MOTACILLIDAE Wagtails, Pipits, Longclaws
97. *Anthus campestris* Tawny Pipit 702
PMW
The nominate race is a rare Palaearctic winter visitor, recorded in January from Juba (pers. observ.).
98. *Anthus cervinus* Red-throated Pipit 713
PMW
A common Palaearctic winter visitor on wet short grassland throughout, from late October to March. Most numerous along the Nile and in the Sudd region.
99. *Anthus leucophrys* Plain-backed Pipit 704
R? LM? NBR
The race *zenkeri* is uncommon, frequenting savanna with scattered trees and bushes in better wooded parts of the south-west; recorded from Didinga, the Kajo Kaji - Lado - Yei area and from Wau to Raffili.
The race *omoensis* may occur along the Ethiopian border.

600. *Anthus novaeseelandiae* Richard's Pipit 706
 V? LM? NBR
 A rare visitor, known only from Kajo Kaji (Brown 1931) and Mongalla. These birds have been assigned to *cinnamomeus*.
601. *Anthus similis* Long-billed Pipit 703
 V? LM? NBR
 The race *hararensis* has only once been recorded, in dry rocky country in the Natoporoputh Hills in the extreme southeast (Macdonald & Cave 1948).
602. *Anthus trivialis* Tree Pipit 708
 PMW PMP
 The nominate race is a common Palaearctic passage migrant throughout, and occasionally winters in well wooded savanna, forest edge and cultivation up to 3000 m; occasionally in late September and October, but mainly from November to April.
603. *Macronyx croceus* Yellow-throated Longclaw 716
 R? LM? BR (10)
 The nominate race is locally common along the southern border from Imatong to Tambura, and in the Sudd, frequenting damper grassland and grassy swamps. Recorded from Juba, Lado, Meshra er Req and Lake No.
604. *Motacilla aguimp* African Pied Wagtail 691
 R LM? NBR
 The race *vidua* is a locally common resident of streams, swamps and waterholes at low altitude, not recorded from the arid extreme south-east, the Sudd or the region northwest of Wau.
605. *Motacilla alba* White Wagtail 690
 PMW
 The nominate race is an uncommon but regular winter visitor in small numbers, mainly from November to early March. Commoner towards the north; this is a common wintering species in the North Sudan.
606. *Motacilla cinerea* Grey Wagtail 694
 PMW PMP
 A common Palaearctic visitor from September to April on forest edges and along rivers up to 3000 m on the Imatong, Dongotona and Didinga Mts. Small numbers of passage birds occur throughout, mainly from late August to November and from late February to April. One ringed as a pullus in the USSR (43°45'N, 40°00'E) was controlled at Gilo in November.
- Motacilla clara* Mountain Wagtail 692
 There is one record of 'white wagtails' in July on forest roads in the Imatongs (Imatong Tea Project). This could well refer to the occurrence of Mountain Wagtails.
607. *Motacilla flava* Yellow Wagtail 695, 697-700
 PMW PMP
 A common winter visitor and passage migrant up to 2000 m throughout, frequenting savanna, swamps, cultivation and towns, often in association with cattle. Two main races occur: the blue-headed *flava* is common throughout from late September to March, whilst the black-headed *feldegg* is almost always associated with swamps and wet grassland, arriving in mid October. Other types have been reported:

yellow-headed *lutea* at Maridi in March; grey-headed *thunbergi* on spring passage at Gilo in April, black-headed *superciliaris* at Wau in October and blue-headed birds with dark ear coverts (? *dombrowskii*) near Malakal in February.

608. *Tmetothylacus tenellus* Golden Pipit 715
 R? LM? NBR
 Uncommon and probably only a dry season visitor to the extreme south-east frequenting arid grassy *Acacia* scrub. Recorded from Kapoeta, Jebel Kathangor and Boma.

Family MALACONOTIDAE Bush Shrikes

609. *Dryoscopus angolensis* Pink-footed Puffback 1132
 R BR (6)
 The race *nandensis* is an uncommon resident of the Imatong primary forests, ranging up to at least 2400 m. It could occur on the Aloma Plateau.
610. *Dryoscopus gambensis* Puffback 1131
 R LM? BR (2, 12)
 The race *Malzacii* is fairly common, frequenting large trees in better wooded savanna southwest of Didinga - Lado - Tonj - Wedweil. It is regular on Didinga up to 2000 m, and presumed wanderers have been observed at Gilo/Imatong in December. The race *erythraeae* occurs in similar habitat along the Ethiopian border.
611. *Laniarius barbarus* Black-headed Gonolek 1119
 R BR (6, 7)
 The race *erythrogaster* is a common resident in any type of wooded savanna throughout except the forest regions of Zande. Birds in the papyrus swamps of the Sudd are assigned to this species, but this needs confirmation for in East African papyrus swamps it is replaced by *L. mufumbiri*.
612. *Laniarius ferrugineus* Tropical Boubou 1125
 R BR (6 - 9)
 The race *major* is a common resident in well wooded savanna southwest of Didinga - Rumbek - Wedweil, while *aethiopicus* is found along the Ethiopian border.
613. *Laniarius funebris* Slate-coloured Boubou 1121
 R BR (10)
 A fairly common resident in the arid country southeast of Torit, preferring thick bushland and *Acacia* scrub.
614. *Laniarius leucorhynchus* Sooty Boubou 1122
 R NBR
 Recorded only from dense secondary forest at Bengengai. Probably a rare resident elsewhere in Zande.
615. *Laniarius luehderi* Lühder's Bush Shrike 1127
 R BR (1 - 3, 4, 11 - 12)
 The nominate race is an uncommon resident on the Aloma Plateau, but is common on Imatong up to 2800 m, frequenting thick forest undergrowth and dense vegetation in forest clearings.

616. *Malaconotus blanchoti* Grey-headed Bush Shrike 1144
 R? LM? NBR
 The race *catharoxanthus* is rather uncommon but widely distributed in better wooded savanna southwest of Didinga - Lado - Rumbek - Raga.
617. *Malaconotus sulfureopectus* Sulphur-breasted Bush Shrike 1138
 R? LM? BR (8, 9)
 The race *similis* is fairly common in better wooded savanna along the southern border. It is also reported from Boma, Wau, Boro and Riki River. There might be some northward movement during the rains.
618. *Nilaus afer* Northern Brubru 1098
 R BR (4, 5, 11)
 Fairly common and probably resident throughout in open tree savanna, bushland and arid *Acacia* savanna. The race *minor* replaces the nominate race in the arid extreme southeast.
619. *Rhodophoneus cruentus* Rosy-patched Shrike 1147
 R NBR
 Uncommon and only known from arid open low thorn-scrub country near the Kenya border.
620. *Tchagra australis* Brown-headed Tchagra 1134
 R LM? BR (12)
 The race *emini* is locally common in secondary growth, forest edge and gallery forest up to 2200 m along the southern border. Recorded from the Didinga and Imatong Mts., Maridi and Bengengai, and north to at least the Tapari River near Amadi.
621. *Tchagra jamesi* Three-streaked Tchagra 1135
 R BR (12)
 The nominate race is confined to the arid extreme southeast where it is fairly common in thick scrub.
622. *Tchagra minuta* Black-capped Tchagra 1136
 R BR (10)
 The nominate race is an uncommon resident of tall wet grassland, forest edge and swamps, including papyrus. Recorded along the southern border from Imatong to Zande and from Boma, Bor, Wau, Malakal and Lake No.
623. *Tchagra senegala* Black-headed Tchagra 1133
 R BR (5, 9)
 Common at low altitude in any kind of woodland with thickets and bushes, including gardens and forest edges. Four races occur: *remigialis* in the extreme north round Renk, *habessinica* east of the Nile south of Malakal, *notha* west of the Nile and *armena* in the extreme southeast from Kapoeta to the Natoporuputh Hills.

Family LANIIDAE Shrikes

624. *Corvinella corvina* Yellow-billed Shrike 1117
 R NBR
 The race *affinis* is a fairly common resident in better wooded savanna and bushland southwest of Didinga - Lado - Rumbek - Wedweil; usually in small parties. The status and validity of the races *togoensis*, *caliginosa* and *chapini* need clarification.

625. *Lanius collaris* Fiscal 1104
 R BR (5, 6)
 Fairly common resident in better wooded savanna and on forest edges up to at least 2000 m. The race *humeralis* occurs east of the Nile on Imatong, Dongotona, Didinga and Boma and *smithii* west of the Nile from Kajo Kaji to Zande.
626. *Lanius collurio* Red-backed Shrike 1112
 PMP PMW?
 An uncommon Palaearctic autumn passage migrant recorded mainly along the southern border below 2000 m from September to December. There are no spring records. A male collected in full moult at Bahr-el-Ghazal in January is exceptional.
627. *Lanius dorsalis* Taita Fiscal 1107
 Fairly common and probably resident in open *Acacia* savanna in the extreme southeast.
628. *Lanius excubitor* Greater Grey Shrike 1100
 LM? PMW NBR
 A common winter visitor in the open semi-arid region around Renk, frequenting single bushes and trees from late September to February. Three races representing Palaearctic and North Sudan breeders could occur: *leucopygos*, *aucheri* and *pallidirostris*.
629. *Lanius excubitorius* Grey-backed Fiscal 1102
 R LM? BR (4, 5, 9)
 The nominate race is common throughout except in the thicker woodlands and forests in the southwest. It is mainly resident and frequents grassland with scattered bushes and trees.
630. *Lanius gubernator* Emin's Shrike 1113
 R BR (3, 4)
 An uncommon resident in open bushy grassland along the southern border, known only from the Yei - Kajo Kaji - Aloma Plateau area; it could be expected elsewhere between Didinga and Maridi.
631. *Lanius isabellinus* Red-tailed Shrike 1114
 PMW PMP
 The race *speculigerus* is a common Palaearctic winter visitor throughout from November to February, associated mainly with swamp edges and wet grassland. The race *phoenicuroides* could occur.
632. *Lanius minor* Lesser Grey Shrike 1103
 PMP
 An uncommon to rare Palaearctic passage migrant from late August to October, with a single record from Juba in November. Many must overfly since the species is common in late August/September in North Sudan. There appear to be no spring records.
633. *Lanius nubicus* Nubian Shrike 1109
 PM
 A common Palaearctic winter visitor to semi-arid bushland in the extreme north around Renk from late September to February. Also recorded from Magoth in the extreme southeast in February and from Wau in late March (M. Rae).

634. *Lanius senator* Woodchat Shrike 1115
 PMW
 The race *niloticus* is a common Palaearctic winter visitor throughout except in the arid extreme southeast and the forests along the southern border, occurs from mid-September to late April or the beginning of May, preferring open grassland with scattered bushes, especially wetter habitats near swamps. The nominate race may also occur, particularly in the west.
635. *Lanius somalicus* Somali Fiscal 1106
 R? NBR
 Known only from the four records in the extreme southeast given by Hall & Moreau (1970).

Family PRIONOPIDAE Helmet Shrikes

636. *Eurocephalus rueppelli* White-crowned Shrike 1097
 R LM? BR (8, 12)
 A common resident in dry thornbush savanna in the southeast, mainly east of Torit. Regular dry season observations around Juba - Lado might refer to wanderers.
637. *Prionops plumata* Helmet Shrike 1090, 1091
 R? BR (4)
 Fairly common at low altitude throughout except in the Sudd, occurring in small parties in wooded savanna, woodland and forest edge. The racial status is not well understood: curly-crested specimens from Boma, Imatong and Didinga are assigned to *cristatus*, others west of Torit to *cocinnatus*; straight crested birds from the Natoporoputh Hills are assigned to *vinaceigularis*.

Family STURNIDAE Starlings, Oxpeckers

638. *Cinnyricinclus leucogaster* Violet-backed Starling 1184
 LM AM? NBR
 The nominate race is seasonally common at Juba and southwest of Didinga - Lado - Rumbek - Wedweil, occurring in any type of tree savanna from January to April and in September/October: *verreauxi* has been recorded from Didinga in March and Mongalla in August.
639. *Cinnyricinclus sharpii* Sharpe's Starling 1185
 R? LM BR (2, 4)
 Fairly common in forests above 1800m on the Imatong, Dongotona and Didinga Mts., though possibly absent during the rainy season.
640. *Creatophora cinerea* Wattled Starling 1182
 LM AM? NBR
 Fairly common in short grass bushland southeast of Lado - Boma from August to March, and sporadically also at Rumbek and Wau.
641. *Lamprotorornis caudatus* Long-tailed Glossy Starling 1195
 R? LM? NBR
 Common in open tree savanna west of the Sue River (Wau), and more rarely east to Tonj.

642. *Lamprotornis chalcurus* Bronze-tailed Glossy Starling 1189
LM NBR
The race *emini* is fairly common in the dry season in better wooded savanna southwest of Torit - Rumbek - Gogrial - Wedweil. At least some move north to breed.
643. *Lamprotornis chalybaeus* Blue-eared Glossy Starling 1188
LM NBR
The race *cyaniventris* occurs in better wooded savanna but is uncommon; possibly a non-breeding visitor from North Sudan. Status is unclear because of confusion in the field with Lesser Blue-eared and Bronze-tailed Glossy Starlings.
644. *Lamprotornis chloropterus* Lesser Blue-eared Glossy Starling 1190
R? LM NBR
The nominate race is fairly common throughout in better wooded savanna, but mainly southwest of Didinga - Bor - Gogrial and along the Ethiopian border.
645. *Lamprotornis purpureus* Purple Glossy Starling 1191
R? LM BR (4)
The race *amethystinus* is fairly common in better wooded savanna west of the Nile.
646. *Lamprotornis purpuropterus* Ruppell's Long-tailed Glossy Starling 1196
R LM? BR (1, 2)
Common throughout in low altitude tree savanna except west of Aweil - Raffile.
647. *Lamprotornis splendidus* Splendid Glossy Starling 1192
R? NBR
The nominate race is uncommon in more mature woodland and forest edges along the Zaire border; also recorded from Boma.
648. *Onychognathus fulgidus* Chestnut-winged Starling 1200
R? NBR
The race *hartlaubii* is recorded only from Bengengai forest (Hall & Moreau 1970).
649. *Onychognathus morio* Red-winged Starling 1203
R? NBR
Uncommon and local, frequenting cliffs and rocky hills in better wooded savanna. Recorded from Mt. Baginzi to Didinga and on the Boma Hills.
650. *Onychognathus walleri* Waller's Chestnut-winged Starling 1201
R NBR
The race *elgonensis* is common on the Imatong and Dongotona Mts. in forest above 1800 m.
651. *Poeoptera stuhlmanni* Stuhlmann's Starling 1209
R BR (4)
Known only from Imatong, where it is common in forests above 1900 m and nests in dead trees.
652. *Spreo shelleyi* Shelley's Starling 1215
R? LM NBR
At least seasonally common from Kapoeta to Lake Turkana, frequenting arid Acacia bushland and showing a preference for Salvadora trees.

653. *Spreo superbus* Superb Starling 1216
 R? LM NBR
 Common from Torit to Jebel Kasangor and Lake Turkana in semi-arid open short grass savanna; moves west to Juba and Lado between August and January.

Sub-family Buphaginae

654. *Buphagus africanus* Yellow-billed Oxpecker 1217
 R? NBR
 The nominate race associates fairly commonly with cattle and game southwest of Didinga - Lado - Gogrial - Wedweil, and could occur throughout.
655. *Buphagus erythrorhynchus* Red-billed Oxpecker 1218
 R? NBR
 Fairly common with cattle and game southeast of Juba - Boma.

Family NECTARINIIDAE Sunbirds

656. *Anthreptes collaris* Collared Sunbird 1271
 R BR (1, 4, 10, 11)
 The race *garguensis* is common on the Imatong, Dongotona and Didinga Mts. in forest and secondary growth up to 2400m. West of the Nile the race *hypodilus* is recorded from Maridi, Yambio, Bengengai and Tambura in similar habitat.
657. *Anthreptes longuemarei* Violet-backed Sunbird 1274
 R? BR (2)
 The nominate race is rather uncommon in tree savanna, forest edge and bushland west of Kajo Kaji - Lado - Ayod and north to the Bahr-el-Arab.
- Anthreptes metallicus* Eastern Pygmy Sunbird 1236
 R? LM NBR
 May occur north of Malakal. Common in the dry *Acacia* savanna of northeast Sudan.
658. *Anthreptes orientalis* Eastern Violet-backed Sunbird 1274
 R? BR (1, 3, 11, 12)
 Common in dry savanna and arid thornbush country southeast of Juba - Lado - Boma.
659. *Anthreptes platurus* Pygmy Sunbird 1236
 R? LM? BR (2)
 Occurs in dry woodland and *Acacia* savanna throughout. Locally common except in the northeast and west.
660. *Anthreptes rectirostris* Green Sunbird 1272
 R? LM? NBR
 The race *tephrolaema* is known only from the lowland forests of Imatong where it is seasonally fairly common in the forest canopy.
661. *Nectarinia amethystina* Amethyst Sunbird 1261
 R? LM? BR (5)
 The race *doggetti* is fairly common in grassy woodland and bushes on the Didinga Mts. above 1600m, and has once been recorded from Imatong.

662. *Nectarinia chloropygia* Olive-bellied Sunbird 1257
 R? NBR
 The race *orphogaster* is rare, and recorded only from the Aloma Plateau, Yambio and Bengengai (Nikolaus, Cave & Macdonald 1955).
663. *Nectarinia coccinigaster* Splendid Sunbird 1239
 R? NBR
 Uncommon, and recorded only from the Yei and Yambio areas; a bird of thickets and forest edge bushes.
664. *Nectarinia cuprea* Copper Sunbird 1238
 R? BR (7, 8)
 The nominate race occurs in open bushy places in wooded savanna south of Imatong - Lado - Gogrial, but is uncommon.
665. *Nectarinia erythroceria* Red-chested Sunbird 1232
 R? NBR
 Uncommon and confined to vegetation near water along the Nile, north to Lake No and Malakal.
666. *Nectarinia famosa* Malachite Sunbird 1227
 R? BR (5)
 The race *cupreonitens* is rather uncommon and known only from Imatong, where it frequents grassy patches near forest above 1800 m.
Nectarinia habessinica Shining Sunbird 1240
 The race *turkanae* may well occur in the extreme southeast, but is not yet recorded.
Nectarinia hunteri Hunter's Sunbird 1264
 Could occur in the arid extreme southeast, but not yet recorded.
667. *Nectarinia mariquensis* Mariqua Sunbird 1245
 R? BR (12)
 The race *osiris* is uncommon and recorded only from the Didinga Mts. where it frequents better wooded savanna and forest edge up to 2000 m.
668. *Nectarinia olivacea* Olive Sunbird 1269
 R BR (6 - 8)
 The race *ragazzii* is common up to 3000 m in any type of forest from Bengengai to Didinga and on the Boma Hills.
669. *Nectarinia osea* Northern Orange-tufted Sunbird 1248
 R? LM? AM? NBR
 The race *decorsei* has been collected from bushy grassland in the area between Kajo Kaji and Bengengai - Tambura. Records are few and confined to the period October to March.
670. *Nectarinia preussi* Northern Double-collared Sunbird 1256
 R BR (1, 10 - 12)
 The race *kikuyuensis* is a very common resident above 1800 m on the Imatong, Dongotona and Didinga Mts., frequenting forest and secondary growth.
671. *Nectarinia pulchella* Beautiful Sunbird 1233
 R? BR (3, 10, 11)
 The nominate race is common throughout in any kind of savanna; absent only from the forests along the southern border.

672. *Nectarinia rubescens* Green-throated Sunbird 1262
 R? BR (9, 10)
 The nominate race is uncommon in low altitude forest and forest edge from Bengengai to Imatong.
673. *Nectarinia seimundii* Little Green Sunbird 1277
 R? LM? AM? BR (3)
 The race *traylori* was found to be very common at Talanga forest in March, possibly as a non-breeding visitor (Traylor & Archer 1982). A specimen was also collected in breeding condition at Bengengai in March (Traylor *in litt.*). A lowland forest species whose status is unclear.
674. *Nectarinia senegalensis* Scarlet-chested Sunbird 1263
 R? BR (4, 8, 9)
 Common in any type of tree and bush savanna. The race *acite* occurs throughout west of Didinga - Malakal, whilst the Ethiopian race *cruentata* is common around Boma.
675. *Nectarinia tacazze* Tacazze Sunbird 1229
 R? LM BR (1)
 The race *jacksoni* occurs above 1800 m on the Imatong, Dongotona and Didinga Mts. Always very common at flowering *Lobelia*, but its whereabouts during periods when this attraction is absent are not clear.
676. *Nectarinia venusta* Variable Sunbird 1251
 R? LM? BR (1, 11, 12)
 Common in open grassy bushland, forest edge and secondary growth at higher altitude. The race *falkensteini* occurs on the Imatong, Dongotona and Didinga Mts., and exceptionally in *Acacia* scrub along the Kenya border (December), while the race *fazugliensis* has been reported from Boma.
677. *Nectarinia verticalis* Green-headed Sunbird 1266
 R? NBR
 The race *viridisplendens* is common up to 3000 m in better wooded savanna and forests from Didinga to Bengengai and north to Tambura, Wau, Boro and Riki River.

Family ZOSTEROPIDAE White-eyes

678. *Zosterops poliogaster* Montane White-eye 1221
 R BR (1 - 4, 11, 12)
 The race *gerhardi* is endemic to the Imatong Mts., where it is very common in forest and secondary growth above 1800 m.
679. *Zosterops senegalensis* Yellow White-eye 1219
 R? LM BR (2, 3, 7, 9, 12)
 The nominate race is common throughout in any kind of wooded savanna at low altitude.

Family PLOCEIDAE Weavers etc.

Sub-family Ploceinae

680. *Amblyospiza albifrons* Grosbeak Weaver 1358
 R? LM BR (11)
 The race *melanota* is fairly common in well-wooded savanna, and on forest edges at Aza and on Imatong up to 2000 m. Breeds in tall grass and papyrus along the Nile to Shambe.
681. *Anaplectes rubriceps* Red-headed Weaver 1359
 R? BR (8)
 The race *leuconotus* is rather uncommon, but widely distributed in well-wooded savanna and open bushland throughout.
682. *Anomalospiza imberbis* Parasitic Weaver 1405
 R? LM? AM? NBR
 The nominate race has been recorded sporadically in moist grassland between Lado and Kajo Kaji, presumably as a non-breeding visitor.
683. *Euplectes afer* Yellow-crowned Bishop 1369
 R? LM BR (9)
 The race *ladoensis* is very common in swamps and wet grassland in the Sudd. Records from similar habitat at Yambio, Aweil and the Diolet River could refer to the nominate race.
Euplectes albonotatus White-winged Widowbird 1373
 The race *eques* may occur in tall bushed grassland along the Uganda border.
684. *Euplectes ardens* Red-naped Widowbird 1375, 1376
 (including the Red-collared Widowbird)
 R? NBR
 The race *concolor* is fairly common in patches of tall grass in low altitude wooded savanna along the southern border from Imatong to Tambura. The race *laticauda* occurs in similar habitat but at higher altitude on the Boma Hills.
685. *Euplectes axillaris* Fan-tailed Widowbird 1370
 R? LM? BR (8)
 The race *phoeniceus* is fairly common in tall, moist bushed grassland, mainly along the Nile, Sobat and Bahr-el-Ghazal west to Aweil.
686. *Euplectes capensis* Yellow Bishop 1367
 R? LM BR (8 - 10)
 The race *crassirostris* is seasonally common up to 2000 m on the Imatong, Dongotona and Didinga Mts., frequenting forest edges and open grassland. Also recorded from the Aloma Plateau and Maridi.
687. *Euplectes franciscanus* Red Bishop 1363
 R LM BR (9 - 11)
 Locally abundant in the Sudd, and common in tall grassland elsewhere, except in the forested region along the Zaire border.
688. *Euplectes gierowii* Black Bishop 1366
 R? NBR
 The race *ansorgei* is uncommon in well-wooded tall grassland at low altitude along the southern border, and also on Boma.

689. *Euplectes hordeaceus* Black-winged Red Bishop 1365
 R LM BR (8 - 10)
 The race *craspedopterus* is very common throughout in bush and tree savanna, sometimes abundant in ranker vegetation during the dry season.
690. *Euplectes macrourus* Yellow-mantled Widowbird 1372
 R? LM? BR (8 - 10)
 The nominate race is common in well-wooded tall grassland in the southwest. There is one old record from Torit.
691. *Malimbus nitens* Blue-billed Malimbe 1357A
 R? NBR
 Recorded only from forest undergrowth at Bengengai (Traylor & Archer 1982).
692. *Malimbus rubricollis* Red-headed Malimbe 1356
 R? NBR
 The nominate race is fairly common in the lowland forests of Imatong up to 1600 m. Rare elsewhere but recorded from the Aloma Plateau and Bengengai.
693. *Ploceus badius* Cinnamon Weaver 1328
 R LM BR (8, 9)
 The race *axillaris* is endemic to the Sudd region where it is abundant in tall grassland near water. In the dry season wanderers have been recorded from Torit, Maridi and Wau and regularly north to Renk.
694. *Ploceus baglafecht* Baglafecht Weaver 1323
 (separated from Emin's Weaver following van den Elzen & Konig (1983))
 R? BR (3)
 The race *eremodius* is uncommon, but recorded from better wooded savanna at Yei, Maridi, Li Rangu and Tambura. Nominate birds have been recorded from Boma.
695. *Ploceus bicolor* Dark-backed Weaver 1335
 R NBR
 The race *mentalis* occurs uncommonly in forest undergrowth in the Imatong foothills.
696. *Ploceus emini* Emin's Weaver 1347
 (separated from Baglafecht Weaver following van den Elzen & Konig (1983))
 R? LM? BR (6 - 8, 11)
 The nominate race is common up to 2400 m on the Imatong, Dongotona and Didinga Mts., frequenting forest clearings and cultivation. Also recorded from Yambio in March and from the Ethiopian border south of Boma.
697. *Ploceus galbula* Rüppell's Weaver 1329
 LM V? NBR
 Only recorded along the Kenya border in November 1979, when it was fairly common in dry *Acacia* woodland (Nikolaus).
698. *Ploceus heuglini* Heuglin's Masked Weaver 1318
 R? BR (8, 9)
 Uncommon and rather local in open woodland southwest of Wau - Rumbek - Lado - Kajo Kaji, nesting in small colonies in Mango trees.

699. *Ploceus insignis* Brown-capped Weaver 1351
R BR (3, 10)
Fairly common above 1800m in forest and forest edge on the Imatong, Dongotona and Didinga Mts.
700. *Ploceus intermedius* Masked Weaver 1319
LM? BR (4, 9, 12)
The nominate race is mainly a rainy season visitor to semi-arid Acacia savanna southeast of Juba - Boma, locally fairly common.
701. *Ploceus jacksoni* Golden-backed Weaver 1326
R? NBR
Apparently rare, and confined to moist localities in lowland grass savanna around Imatong and Didinga.
702. *Ploceus luteolus* Little Weaver 1324
R? BR (5, 8, 9)
The nominate race is common throughout in any kind of lowland savanna woodland. The race *kavirondensis* may occur along the Uganda border.
703. *Ploceus melanocephalus* Yellow-backed Weaver 1325
R? LM? BR (8)
The race *fischeri* is fairly common in the southwest, north to Wau and east to Imatong. It frequents tall grassland near water, especially during the breeding season.
704. *Ploceus melanogaster* Black-billed Weaver 1346
R BR (1 - 3, 11, 12)
The race *stephanophorus* is common above 1600m on the Imatong, Dongotona and Didinga Mts., frequenting forest undergrowth and cultivation.
705. *Ploceus nigerrimus* Black Weaver 1344
R? BR (8, 11)
The nominate race is fairly common in thick tree savanna and forest edge with open grassland along the Zaire border from Tambura to the Aloma Plateau.
706. *Ploceus nigricollis* Black-necked Weaver 1336
R? LM? NBR
The nominate race is uncommon, and recorded only from savanna country and forest edge at Li Rangu and Bengengai.
707. *Ploceus ocularis* Spectacled Weaver 1337
LM? NBR
The race *crocatu*s is uncommon in low altitude wooded savanna and forest edge along the southern border from Imatong to Tambura, as well as at Lado and the Boro River. All records are from the dry season.
708. *Ploceus rubiginosus* Chestnut Weaver 1327
LM? AM? NBR
The nominate race is rather uncommon but occurs regularly in November/December in the Acacia savanna of the extreme southeast; probably a non-breeding visitor.
709. *Ploceus superciliosus* Compact Weaver 1356
R? LM? NBR
Recorded only from Boma, Imatong, Tambura and Bengengai where it is locally common at low altitude in well wooded grassland

710. *Ploceus taeniopterus* Northern Masked Weaver 1316
 R LM? BR (6 - 11)
 The nominate race is locally abundant along the Nile and Sobat Rivers in tall grassland, nesting in large "sunt" *Acacia* trees. Breeding is also reported along the Zaire border.
711. *Ploceus velatus* Vitelline Masked Weaver 1321
 R LM? BR (6 - 9, 11)
 Common in wooded grassland and cultivation at low altitude. The race *vitellinus* occurs north of 10°N (Melut), while *uluensis* ranges north to Boma, Shambe and Aweil and west to Wau and Tambura.
712. *Quelea cardinalis* Cardinal Quelea 1362
 R? LM? NBR
 The nominate race is uncommon in tall grass savanna under 2000 m in the Maridi - Lado - Imatong area.
713. *Quelea erythrops* Red-headed Quelea 1361
 R? LM BR (7, 8)
 Uncommon in moist grass patches in better wooded savanna. Recorded from Juba to Tambura and north to Wau, probably only as a breeding visitor. A wanderer was collected in dry *Acacia* savanna east of Kapoeta in December.
714. *Quelea quelea* Red-billed Quelea 1360
 R LM BR (9)
 The race *aethiopica* is seasonally abundant along the Nile south to the Sudd, especially in the dry season, but uncommon to rare elsewhere. Breeding colonies have appeared with the recent establishment of large irrigation schemes such as that at Bor. This species is much more common in the dry bush savanna of North Sudan. A bird ringed in Ethiopia north of Lake Turkana (5°05'N, 35°57' E) was recovered after two years 387 km west, near Torit.

Sub-family Bubalornithinae

715. *Bubalornis albirostris* White-billed Buffalo Weaver 1284
 R? LM NBR
 Fairly common in the *Acacia* savanna around Kapoeta and Jebel Kasangor. It is not clear whether these are just non-breeding visitors, as the few birds which occur further west around Juba appear to be.
716. *Bubalornis niger* Red-billed Buffalo Weaver 1285
 R? BR (8, 9)
 The race *intermedius* is fairly common in dry *Acacia* savanna east of Kapoeta.
717. *Dinemellia dinemellia* White-headed Buffalo Weaver 1286
 R LM BR (8, 9, 12)
 The nominate race is common in semi-arid *Acacia* savanna east of Torit; wanderers move west to Juba and Lado in the dry season. There are three old skins from the Li Rangu/Yambio area, collected in April and October, showing clean white feathering unlike that of other Sudan specimens (Woodman - Khartoum Museum).

Sub-family Plocepasserinae

718. *Plocepasser mahali* White-browed Sparrow Weaver 1287
 R BR (2, 6, 9, 10, 12)
 The race *melanorhynchus* is common in dry *Acacia* savanna southeast of Juba, Lado and Jebel Kathangor.
719. *Plocepasser superciliosus* Chestnut-crowned Sparrow Weaver 1288
 R BR (7)
 Common in better wooded savanna southwest of Kafia Kingi - Tonj - Lado - Didinga, and also along the Ethiopian border where it is recorded from Boma.
720. *Pseudonigrita arnaudi* Grey-headed Social Weaver 1291
 R LM? BR (8 - 12)
 The nominate race is locally common in *Acacia* savanna north to Boma and Ayod and west to Lado.

Sub-family Passerinae

- Passer castanopterus* Somali Sparrow 1298
 The race *fulgens* probably occurs in the arid extreme southeast near Lake Turkana.
721. *Passer domesticus* House Sparrow 1293
 R BR (1 - 12)
 The race *arboreus* is common in towns with solid buildings along the Nile south to Malakal where it was probably only recently introduced.
722. *Passer eminibey* Chestnut Sparrow 1306
 R? LM? BR (8 - 10)
 Uncommon and local in *Acacia* country from Juba and Lado east to the Kenya border; also recorded from Malakal and Meshra er Req.
723. *Passer griseus* Grey-headed Sparrow 1300, 1303
 (including Parrot-billed Sparrow)
 R LM? BR (9 - 11)
 The nominate race is very common throughout, except along the Ethiopian border, in any kind of tree and bush savanna. The race *gongonensis* is uncommon in the arid extreme southeast where it tends to replace the nominate race.
724. *Passer luteus* Sudan Golden Sparrow 1304
 LM NBR
 The nominate race is fairly common in the dry season in arid savanna around Renk, and occasionally south to the Sobat River.
725. *Passer motitensis* Rufous Sparrow 1294
 R? BR (9)
 The race *shelleyi* is uncommon and recorded only from Lado and Jebel Kathangor, in semi-arid *Acacia* bushland, sometimes near villages.
726. *Petronia dentata* Bush Petronia 1309
 R LM? BR (1, 2, 12)
 Common throughout at low altitude in tree and bush savanna.
727. *Petronia pyrgita* Yellow-spotted Petronia 1308
 R? NBR
 The nominate race is fairly common in wooded savanna on rocky ground in the arid extreme southeast.

Sub-family Sporopipinae

728. *Sporopipes frontalis* Speckle-fronted Weaver 1311
 R? BR (1, 9, 12)
 Common in dry *Acacia* savanna. The race *emini* occurs southeast of Juba - Lado - Boma, while the nominate race ranges south to Wau and Ayod.

Sub-family Viduinae

Indigo Birds: Identification of Indigobirds is very difficult: information on song and on the bill and foot colour of the live bird may be necessary. It is even more difficult to assign the right estrildid host for some species, where field observations are scanty.

729. *Hypochera camerunensis* "Camerunensis" Indigobird 1439
 R? NBR
 Rare, and only reported from well wooded savanna at Li Rangu; possibly parasitic on the Black-faced Firefinch *Lagonosticta larvata*.
730. *Hypochera chalybeata* Red-billed Firefinch Indigobird 1435
 R? BR (8, 9)
 The race *ultramarina* is fairly common in open tree and bush savanna and cultivation, recorded from Wau, Gogrial, Yei, Amadi and along the Nile from Juba to Renk; parasitic on the Red-billed Firefinch *Lagonosticta senegala*.
731. *Hypochera nigeriae* "Nigeriae" Indigobird 1440
 R? NBR
 Rare in open wooded savanna; recorded from Li Rangu, Mongalla and Wau. Possibly parasitic on the Black-bellied Firefinch *Lagonosticta rara*.
732. *Hypochera wilsoni* Bar-breasted Firefinch Indigobird 1437
 R? BR (9, 10)
 Fairly common in well wooded savanna southwest of Raffile - Lado - Torit; parasitic on the Bar-breasted Firefinch *Lagonosticta rufopicta*.
733. *Vidua fischeri* Straw-tailed Whydah 1443
 R? LM BR (12)
 Seasonally fairly common in semi-arid *Acacia* savanna east of Torit. Mainly parasitic on the Purple Grenadier *Uraeginthus ianthinogaster* but possibly also on the Blue-capped Cordon Bleu *Uraeginthus cyanocephalus* (Nikolaus 1979).
734. *Vidua hypocherina* Steel-blue Whydah 1442
 R? LM? NBR
 Rare, and known only from arid open *Acacia* savanna east of Kapoeta (Nikolaus 1979), where it is mainly parasitic on the Black-cheeked Waxbill *Estrilda erythronotus*.
735. *Vidua interjecta* Northern Broad-tailed Paradise Whydah W1386
 R? LM NBR
 Fairly common in open woodland along the southern border south of Tambura, Lado and Torit; parasitic on the Red-winged Pytilia *Pytilia phoenicoptera*.

36. *Vidua macroura* Pin-tailed Whydah 1441
 R? LM BR (8 - 11)
 Common throughout in any kind of grass savanna; some move north during the rainy season. Parasitic on Waxbills *Estrilda* spp.
37. *Vidua orientalis* Northern Paradise Whydah 1445
 R? LM NBR
 The nominate race is fairly common in open woodland and bush savanna north of Aweil - Malakal. Parasitic on the Green-winged Pytilia *Pytilia melba* (races having red lores).
38. *Vidua paradisaea* Paradise Whydah 1444
 R? LM? BR (9)
 The nominate race is fairly common in open woodland and bush savanna southwest of Juba - Lado - Boma, where it is parasitic on the Green-winged Pytilia *Pytilia melba* (races having grey lores).
- Family ESTRILDIDAE Waxbills etc.
 Sub-family Estrildinae
39. *Amandava subflava* Zebra Waxbill 1421
 R? LM? BR (6 - 9)
 The nominate race is widespread but uncommon in swamps and tall grass savanna near water throughout.
40. *Clytospiza dybowskii* Dybowski's Twinspot 1390
 R? BR (10)
 Rare at open grass patches in lowland forests and recorded from Bengengai, Li Rangu and Lotti Forest.
41. *Clytospiza monteiri* Brown Twinspot 1389
 R BR (9)
 Fairly common in moist thickets west of the Nile, north to Lado, Wau and Kafila Kingi.
42. *Cryptospiza salvadorii* Abyssinian Crimson-wing 1399
 R BR (1, 10 - 12)
 Common up to 3000 m in forest undergrowth and clearings on the Imatong, Dongotona and Didinga Mts. Best regarded as an intergrade between *kilimensis* and the nominate race.
43. *Estrilda astrild* Waxbill 1418
 R? LM? NBR
 The race *peasei* is uncommon in tall grassland along the Nile, Bahr-el-Ghazal and Sobat Rivers.
44. *Estrilda erythronotus* Black-cheeked Waxbill 1427
 R? NBR
 Rare in open Acacia savanna between Torit and the Kenya border; status unknown.
45. *Estrilda melanotis* Yellow-bellied Waxbill 1417
 R? LM BR (10, 11)
 The race *kilimensis* (not *quartinia*) is common up to 2400 m in grassland on the Imatong, Dongotona and Didinga Mts.; seen mainly during the dry season.

746. *Estrilda nonnula* Black-crowned Waxbill 1425
 R LM? BR (10, 11)
 The nominate race occurs up to 2800m in forest clearings and cultivation from Tambura to Didinga. It is very common on the Imatong Mts. but uncommon elsewhere
747. *Estrilda paludicola* Fawn-breasted Waxbill 1422
 R? BR (7, 8)
 Fairly common in open damp wooded grassland within better wooded savanna. The nominate race occurs southwest of Imatong - Lado - Rumbek - Wau - Kafia Kingi and *ochrogaster* along the Ethiopian border from Baro River to Boma.
748. *Estrilda rhodopygia* Crimson-rumped Waxbill 1420
 R? NBR
 The race *centralis* is a rare bird of grass savanna, recorded only from Lado, the Aloma Plateau and Didinga; status unknown.
749. *Estrilda troglodytes* Black-rumped Waxbill 1419
 R? LM? NBR
 Fairly common in grass savanna with thickets near water, mainly west of the Nile.
750. *Lagonosticta larvata* Black-faced Firefinch 1416
 R? BR (9)
 Fairly common in better wooded tall grassland. The race *nigricollis* occurs west of the Nile throughout except in the Sudd, while the nominate race is restricted to the Boma area.
751. *Lagonosticta rara* Black-bellied Firefinch 1426
 R? BR (8 - 11)
 The nominate race is fairly common in better wooded savanna and the edge of gallery forest southwest of Didinga - Rumbek - Kafia Kingi.
752. *Lagonosticta rhodopareia* Jameson's Firefinch 1411, 1412
 R? LM? BR (12)
 The nominate race is fairly common in open grassland at higher altitudes on the Boma Hills and Didinga Mts.
753. *Lagonosticta rubricata* African Firefinch 1411
 R? LM? BR (7)
 Fairly common in tall grass and overgrown cultivation up to 2400m, moving to lower altitude for breeding during the breeding season. the race *hildebrandti* occurs east of the Nile from Imatong to the Didinga Mts. and the Boma Hills; *congica* from the Aloma Plateau to Bengengai.
754. *Lagonosticta rufopicta* Bar-breasted Firefinch 1414
 R? BR (6 - 10)
 The nominate race is fairly common throughout north to the Bahr-el-Arab and Sobat Rivers, frequenting trees and bush savanna and cultivation.
755. *Lagonosticta senegala* Red-billed Firefinch 1413
 R BR (7, 8)
 The race *rhodopsis* is common in grass savanna and cultivation throughout, except in the better wooded savanna and forest area southwest of Juba, Rumbek and Wau. It intergrades with *ruberrima* along the Uganda border.

756. *Mandingoa nitidula* Green-backed Twinspot 1407
LM BR (1)
The race *chubbi* is a common breeding visitor from November to March, Occasionally till May, to thick forest undergrowth and thickets up to 2400m on the Imatong, Dongotona and possibly the Didinga Mts.
757. *Nesocharis capistrata* Grey-headed Olive-back 1428
R? NBR
Apparently uncommon or rare in gallery forests with tall trees. Recorded only from Maridi, Wau and Boro River.
758. *Nigrita canicapilla* Grey-headed Negrofinch 1386
R LM? BR (1, 3, 12)
The race *schistacea* is fairly common in lowland forests and forest clearings up to 2400m on the Imatong Mts.; the race *canicapilla* was once recorded from Bengengai (Traylor & Archer 1982).
759. *Ortygospiza atricollis* Quailfinch 1403
R? LM? NBR
The nominate race is uncommon in dry or wet short grass swamps. Recorded around Kajo Kaji, Juba, Rumbek, Tonj, Aweil and Malakal.
Ortygospiza locustella Locust Finch 1404
The race *uelensis* is recorded along the Zaire border from Bengengai to the Aloma Plateau, but not yet on Sudan territory. In this area or north of it Chapin suggests breeding during the late rainy season in marshy grassland.
760. *Pyrenestes ostrinus* Large-billed Seedcracker 1393-1395
R? NBR
Uncommon in dense thicket, forest edge and forest undergrowth. All three races have been recorded along the Zaire border: *frommi* from Li Rangu, Yei and the Aloma Plateau, *rothschildi* and nominate from Bengengai.
761. *Pytilia afra* Orange-winged Pytilia 1409
R? LM? NBR
Rare in open wooded savanna, and recorded only from Kajo Kaji, Didinga and Rejaf in April (B.M.); status unknown.
762. *Pytilia melba* Green-winged Pytilia 1410
R BR (1, 2, 12)
Common throughout in any kind of bush and open tree savanna. Four races are recognised: *criterior* north of the Bahr-el-Arab and Sobat Rivers, *clanceyi* south of it to Tambura and Shambe, where it intergrades with *belli* to the south; *soudanensis* occurs east of the Nile north at least to Boma.
763. *Pytilia phoenicoptera* Red-winged Pytilia 1408
R? NBR
The nominate race is rather uncommon in open tree savanna with thick bushes, recorded from Kajo Kaji and Lado west to Tambura and Wau.
764. *Spermophaga ruficapilla* Red-headed Bluebill 1391
R BR (1, 4)
The nominate race is fairly common but unobtrusive in thick moist forest undergrowth from Bengengai to Imatong up to 2400m.

765. *Uraeginthus bengalus* Red-cheeked Cordon-bleu 1431
 R BR (9 - 11)
 The nominate race is common throughout at low altitude in any kind of savanna and cultivation except in the arid extreme southeast where it is replaced by the Blue-capped Cordon-bleu *U. cyanocephalus*.
766. *Uraeginthus cyanocephalus* Blue-capped Cordon-bleu 1432
 LM BR (12)
 An uncommon breeding visitor, restricted to semi-arid *Acacia* savanna east of Kapoeta (Nikolaus 1973).
767. *Uraeginthus ianthinogaster* Purple Grenadier 1433
 R LM NBR
 Uncommon in *Acacia* thickets and bush country east of Didinga. In the dry season it wanders west to Torit.
- Sub-family Amadinae
768. *Amadina fasciata* Cut-throat 1402
 R? LM? NBR (11)
 Uncommon in dry *Acacia* savanna. The nominate race is probably only a wanderer south to Wau and Gogrial; birds southeast of Lado - Jebel Kasangor are nearer to *alexanderi*.
769. *Lonchura bicolor* Black and White Mannikin 1380
 LM BR (9, 10)
 The race *poensis* is a common breeding visitor to the Imatong and Didinga Mts., up to 2400 m, favouring small grass patches on forest edges and cultivation. Moves to low altitude from April to July. Once recorded from Maridi in August.
770. *Lonchura cucullata* Bronze Mannikin 1379
 R LM BR (7, 9, 11)
 The nominate race is seasonally common in grassland and cultivation up to 2000 m southwest of Didinga - Lado - Rumbek and the Bahr-el-Arab. Birds from Boma are nearer to *scutata*.
771. *Lonchura fringilloides* Magpie Mannikin 1382
 LM BR (11)
 Fairly common from October to May on the Imatong Mts. in secondary growth and grassy forest edge up to 2000 m; also recorded from Maridi and Bengengai. Status unknown.
772. *Lonchura griseicapilla* Grey-headed Silverbill 1384
 LM NBR
 An uncommon dry season visitor in small parties to the dry savanna and *Acacia* bushland between Juba and the Kenya border.
773. *Lonchura malabarica* Silverbill 1383
 R? BR (12)
 Uncommon, mainly as a dry season visitor, in semi-arid grassland. The race *orientalis* occurs south of Boma - Lado, ranging west to Maridi in the dry season, while *cantans* is found north of the Bahr-el-Arab and Sobat Rivers.

Family FRINGILLIDAE Buntings, Canaries, Seed-eaters

Sub-family Emberizinae

774. *Emberiza cabanisi* Cabanis' Bunting 1467
 R BR (7)
 The nominate race is uncommon in open bush savanna from Tambura to Yei.
Emberiza caesis Cretzschmar's Bunting 1473
 May occur as a Palaearctic winter visitor in arid bush savanna north of Melut.
775. *Emberiza flaviventris* Golden-breasted Bunting 1469
 R? NBR
 The race *kalaharica* (*caryochroa*) frequents open tree savanna along the southern border, with records from Maridi, Yei and Didinga; *flavigastra* should occur north of Melut.
776. *Emberiza forbesi* Brown-rumped Bunting 1471
 R BR (3)
 The nominate race is fairly common throughout in open bush and tree savanna.
Emberiza hortulana Ortolan 1472
 Probably occurs as a Palaearctic winter visitor in green bushland round Renk.
777. *Emberiza poliopleura* Somali Golden-breasted Bunting 1470
 R? BR (4, 12)
 Uncommon in arid *Acacia* bush savanna in the extreme southeast.
778. *Emberiza striolata* House Bunting 1477
 R? LM V? NBR
 The race *saturator* was collected in March 1915 by Butler at Kajo Kaji. Frequents arid stony country around Lake Turkana and in North Sudan.
779. *Emberiza tahapisi* Cinnamon-breasted Rock Bunting 1476
 R LM NBR
 Fairly common and always associated with rocky hills, grass and bushes. Three races probably meet: *goslingi* in the west around the Boro/Riki River, nominate along the southern border from Zande to Boma, and *septemstriata* east of the Nile south to the Sobat River.

Sub-family Carduelinae

780. *Linurgus olivaceus* Oriole Finch 1463
 R BR (1, 2, 11)
 The race *elgonensis* is common in forest and forest clearings above 1800 m on the Imatong Mts.
781. *Serinus atrogularis* Yellow-rumped Seed-eater 1459
 R? NBR
 The race *reichenowi* is uncommon in dry *Acacia* savanna southeast of Kapoeta. There is one recent August record from Tambura (Nikolaus).

782. *Serinus canicollis* Yellow-crowned Canary 1454
 R BR (11)
 The race *flavivertex* is rather uncommon above 1800 m on the Imatong Mts., favouring *Podocarpus* trees in highland forest.
783. *Serinus citrinelloides* Grey-faced African Citril 1464
 R BR (1 - 3, 10 - 12)
 The nominate race is very common up to 2400 m on the Imatong, Dongotona and Didinga Mts., moving to lower altitudes in the rainy season. A species of forest clearings, secondary growth and cultivation.
784. *Serinus dorsostriatus* White-bellied Canary 1449
 R? LM? NBR
 There is one record of the race *maculicollis*, from dry bush savanna in the Natoporoputh Hills (Macdonald & Cave 1948).
785. *Serinus gularis* Streaky-headed Seed-eater 1456
 R? LM NBR
 The race *elgonensis* is uncommon in dry woodland, recorded from Wau and from Tambura to Yei between November and March.
786. *Serinus leucopygius* White-rumped Seed-eater 1458
 R LM? NBR
 The nominate race is rather uncommon in wooded grassland near water, recorded along the Nile from Gogrial, Rumbek and Torit. Extending its range with the development of new irrigation schemes.
787. *Serinus mozambicus* Yellow-fronted Canary 1448
 R? LM? BR (9)
 Common in tree and bush savanna. The race *barbatus* ranges throughout west of Didinga - Ayod, while *grotei* occurs along the Ethiopian border.
788. *Serinus reichardi* Stripe-breasted Seed-eater 1456
 R? LM? NBR
 The race *striatipectus* is uncommon in open tree savanna, recorded from Yambio, Yei, Didinga and Boma from November to May.
789. *Serinus striolatus* Streaky Seed-eater 1461
 R BR (1)
 The nominate race is very common on the Imatong Mts. in secondary scrub, cultivation and forest edge above 1800 m. Also recorded from the Dongotona Mts.

GAZETTEER

Aloma Plateau	ca. 3°45'N, 30°45'E	1200 m
Amadi	5°30'N, 30°22'E	
Aweil	8°45'N, 27°24'E	
Ayod	8°08'N, 28°22'E	
Aza Forest	4°42'N, 29°50'E	700 m
Baginzi, Mt.	4°28'N, 29°01'E	1067 m
Baro River	ca. 8°30'N, 33°30'E	
Bengengai	4°49'N, 27°43'E	
Boma Hills	6°10'N, 34°30'E	1500 m
Bor	6°13'N, 31°33'E	420 m
Boro River	8°27'N, 24°47'E	ca. 800 m
Didinga Mts.	4°25'N, 33°30'E	2795 m
Diofo River	9°03'N, 24°30'E	ca. 800 m
Dongotona Mts.	4°10'N, 33°07'E	2623 m
Gambela, Ethiopia	8°15'N, 34°35'E	
Geigar	11°58'N, 32°45'E	
Gilo/Itibol	4°00'N, 32°51'E	1900 m
Gogrial	8°32'N, 28°05'E	
Ilemi Triangle	ca. 4°35'N, 35°00'E	500 m
Imatong Mts.	3°55'N, 32°55'E	3187 m
Jongolei	6°50'N, 31°15'E	
Juba	4°52'N, 31°30'E	450 m
Kafia Kingi	9°16'N, 24°23'E	600 m
Kagelu	4°03'N, 30°37'E	
Kajo Kaji	3°52'N, 31°40'E	
Kaka	10°35'N, 32°10'E	
Kapoeta	4°45'N, 33°35'E	
Kasangor, Jebel	5°45'N, 31°07'E	1063 m
Kenya/Sudan Border	4°32'N, 34°14'E	700 m
Kinyeti, Mt.	3°55'N, 32°55'E	3180 m
Kipia	3°56'N, 32°57'E	2600 m
Kodok	9°53'N, 32°07'E	
Kongor	7°10'N, 31°20'E	
Kosti	13°10'N, 32°42'E	380 m
Lado	5°06'N, 31°30'E	500 m
Li Rangu	4°45'N, 28°22'E	
Lotti Forest	4°03'N, 32°32'E	1250 m
Malakal	9°30'N, 31°40'E	386 m
Maridi	4°55'N, 29°28'E	
Mbili	7°25'N, 28°12'E	
Melut	10°25'N, 32°15'E	
Meshra er Req	8°25'N, 29°15'E	
Mongalla	5°10'N, 31°47'E	
Muolo	6°05'N, 29°58'E	
Nagishot	4°15'N, 33°35'E	1900 m
Natoporoputh Hills	4°52'N, 34°00'E	800 m

Nibor, Lake	6°52'N, 30°15'E	
Nimule	3°36'N, 32°02'E	
No, Lake	9°23'N, 30°25'E	
Raffili	6°57'N, 27°58'E	
Raga	8°27'N, 25°40'E	
Renk	11°45'N, 32°47'E	
Riki River	8°95'N, 24°35'E	ca. 1000 m
Rumbek	6°48'N, 29°40'E	
Shambe	7°05'N, 30°45'E	
Sobat River	ca. 9°00'N, 32°30'E	
Talanga Forest	4°01'N, 32°43'E	950 m
Tambura	5°35'N, 27°27'E	
Tapari River	5°08'N, 30°52'E	
Tonj	7°17'N, 28°45'E	
Torit	4°25'N, 32°35'E	
Turkana, Lake	4°30'N, 35°55'E	375 m
Wau	7°43'N, 27°58'E	430 m
Wedweil	8°58'N, 27°12'E	
Yambio	4°34'N, 28°25'E	
Yei	4°05'N, 30°40'E	
Zande	4°30'N, 28°30'E	
Zeraf, Bahr el	9°32'N, 31°10'E	

REFERENCES

- ASH, J.S. 1977. Four species of birds new to Ethiopia, and other notes. *Bulletin of the British Ornithologists' Club* 97: 4-9.
- & POMEROY, D.E. 1981. Mapping schemes in the Afrotropical region. *Scopus* 5: 36-38.
- BOWEN, W.W. 1926 & 1931. *Catalogue of Sudan birds*. Khartoum: Sudan Government Museum. Part 1 & 2.
- 1925, 1926, 1928. The Game birds and Waterfowl of the Sudan. *Sudan Notes and Records* 8: 85-95, 9: 107-124, 11: 69-82.
- BROWN, L.H., URBAN, E.K. & NEWMAN, K. 1982. *The birds of Africa*. Vol.1. London and New York: Academic Press.
- BRITTON, P.L. (ed.) 1980. *Birds of East Africa*. Nairobi: EANHNS.
- 1971. On the apparent movements of *Cossypha natalensis*. *Bulletin of the British Ornithologists' Club* 91: 137-144.
- BUTTLER, A.L. 1905. A contribution to the ornithology of the Egyptian Sudan. *Ibis* 8 (5): 301-401
- 1908. A second contribution to the ornithology of the Egyptian Sudan. *Ibis* 9 (4): 205-263.
- 1909. On birds collected by Cpt. E.P. Blencowe in the Bahr-el-Ghazal Province. *Ibis* 10: 71-90.
- CAVE BEY, F.O. 1938. A note on birds not previously recorded from the Southern Sudan. *Sudan Notes and Records* 21: 167-188.
- 1939. A further list of new records of birds for the Southern Sudan. *Sudan Notes and Records* 22: 159-162.
- CAVE, F.O. 1946. Some interesting birds from the Southern Sudan. *Sudan Notes and Records* 26: 225-228.
- & MACDONALD, J.D. 1955. *Birds of the Sudan*. Edinburgh: Oliver & Boyd.
- CHAPIN, J.P. 1930, 1939, 1953. The birds of the Belgian Congo. *Bulletin of the American Museum of Natural History* 65, 75, 75a, 75b. Vol.1-4.
- CLANCEY, P.A. 1975. On the species limits of *Acrocephalus beatius* (Viellot) of Ethiopian Africa. *Arnoldia* 7, No.20: 1-14.
- CUNNINGHAM-VAN SOMEREN, R.G. & SCHIFTER, H. 1981. New races of montane birds from Kenya and Southern Sudan. *Bulletin of the British Ornithologists' Club* 101: 347-354, 355-363.
- DOWSETT, R.J. 1971. The Lesser Grey Shrike *Lanius minor* in Africa. *Ostrich* 42.
- 1973. The River Warbler *Locustella fluviatilis* in Africa. *Zambia Museum Journal* 3 (1972): 69-79.

- DOWSETT, R.J. 1977. The distribution of some falcons and plovers in East Africa. *Scopus* 1: 73-78.
- ELLIOTT, C.C.H. 1972. An ornithological survey of the Kidepo National Park in North Uganda. *Journal of the East Africa Natural History Society and National Museum* Vol.28 No.129: 1-34.
- ERARD, C. 1974. The problem of the Boran *Cisticola*. *Bulletin of the British Ornithologists' Club* 94: 26-38.
- FRIEDMANN, H. 1978. Results of the Lathry's Central African Republic Expedition 1976, ornithology. *Contributions in Science, Los Angeles County Museum of Natural History* No.287: 1-22.
- FRYXELL, J. 1980. Preliminary report on an aerial survey of the Boma National Park region October 1980. New York Zoological Society.
- GRANT, O. 1907. Birds from Sobat and Baro River. *Ibis* 19: 578-614.
- GUILLET, A. 1978. Distribution of the Shoebill Stork *Balaeniceps rex* in the Southern Sudan. *Biological Conservation* 13(1): 39-50.
- HALL, B.P & MOREAU, R.E. 1970. *An atlas of speciation in African passerine birds*. London: British Museum (Natural History).
- HARTLAUB, G. 1881. Beitrag zur Ornithologie der östlich äquatorialen Gebiete Afrikas. *Abhandlungen naturkundliche Vereinigung Bremen* 7: 83-128.
- 1882. Zweiter Beitrag zur Ornithologie der östlich-äquatorialen Gebiete Afrikas. *Abhandlungen naturkundliche Vereinigung Bremen* 8: 184-232.
- KOENIG, A. 1911. Die Ergebnisse meiner Reise nach dem Sudan im Frühjahr 1910. *Verhandlungen vom Internationalen Ornithologen Kongress, Berlin* 1910: 498-545.
- 1926. Die Ergebnisse meiner zweiten Forschungsreise in das Gebiet der Quellflüsse des Nils. *Journal für Ornithologie* 74: 315-361.
- MACDONALD, J.D. & CAVE, F.O. 1948. Recent addition to the Sudan checklist of birds. *Ibis* 90: 240-251.
- 1950. Notes on Sudan birds. *Ibis* 92: 377-379.
- MACKWORTH-PRAED, C.W. & GRANT, C.H.B. 1952, 1955. *African Handbook of birds*. Series 1, Vols.1 & 2. *Birds of eastern and north eastern Africa*. London: Longmans Green & Co.
- 1970, 1973. *African Handbook of birds*. Series 3, Vols 1 & 2 *Birds of west central and western Africa*. London: Longmans Green & Co.
- MADDEN, J.F. 1945. Some common birds of Torit District, Equatoria Province. *Sudan Notes and Records* 26: 285-300.
- MOREAU, R.E. 1966. *The bird faunas of Africa and its islands*. London: Academic Press.

- NIKOLAUS, G. 1979. Notes on some birds new to South Sudan. *Scopus* 3: 68-73.
- _____ 1979. The first record of Basra Reed Warbler in the Sudan. *Scopus* 3: 103-104.
- _____ 1982. Further notes on some birds new to South Sudan. *Scopus* 6: 1-4.
- _____ 1982. A new race of the Spotted Ground Thrush *Turdus fischeri* from South Sudan. *Bulletin of the British Ornithologists' Club* 102: 45-47.
- _____ 1983. An important passerine ringing site near the Sudan Red Sea coast. *Scopus* 7: 15-18.
- _____ 1984. Further notes on birds new or little known in the Sudan. *Scopus* 8: 36-38.
- _____ 1987. Distribution atlas of Sudan birds with notes on habitats and status. *Bonner Zoologische monographien* Nr.25.
- _____ & BACKHURST, G.C. 1982. First ringing report for the Sudan. *Scopus* 6: 77-90.
- PEARSON, D.J. 1979. The races of the Red-tailed Shrike *Lanius isabellinus* occurring in East Africa. *Scopus* 3: 74-78.
- V.PELZELN, A. 1881. Über eine Sendung von Vögeln aus Centralafrika von Emin Bey. *Verhandlungen der K.K. Zoologisch-Botanischen Gesellschaft in Wien* 31: 141-156.
- _____ 1881. Über Dr. Emin Bey's zweite Sendung von Vögeln aus Centralafrika. *Verhandlungen der K.K. Zoologisch-Botanischen Gesellschaft in Wien*. 31: 605-618.
- SCHÜZ, E & BÖHRINGER, R. 1950. Vom Zug des Weissstorchs in Afrika und Asien nach Ringfunden bis 1949. *Die Vogelwarte* 161-226.
- SHARPE, R.B. 1901. On the collection of birds made by Dr. A. Donaldson-Smith on his last expedition to Lake Rudolf and the Nile. *Proceedings of the Zoological Society of London* 1901: 602-622.
- SHELLEY, G.E. 1888. On a collection of birds made by Emin Pasha in Equatorial Africa. *Proceedings of the Zoological Society of London* 1888: 17-50.
- SNOW, D.W. (ed.) 1978. *An atlas of speciation in African non-passerine birds*. London: British Museum (Natural History).
- TRAYLOR, M.A. & ARCHER, A.L. 1982. Some results of the Field Museum 1977 expedition to South Sudan. *Scopus* 6: 5-12.
- TRAYLOR, M.A. 1970. East African *Bradornis*. *Ibis* 112: 513-531.
- URBAN, E.K. & BROWN, L.H. 1971. *A checklist of the birds of Ethiopia*. Addis Ababa: University Press.
- VAN DEN ELZEN, R. & KÖNIG, C. 1983. Vögel des (Süd-)Sudan: taxonomische und tiergeographische Bemerkungen. *Bonner Zoologische Beiträge* 34: 149-196.

- VAURIE, C. 1959. *The birds of the Palearctic fauna. Passeriformes.* London: Witherby.
- _____ 1965. *The birds of the Palearctic fauna. Non-Passeriformes.* London: Witherby.
- WETMORE, A. 1960. A classification for the birds of the world. *Smithsonian Miscellaneous collection* 139 (11): 1-37.
- WHITE, C.M.N. 1960. A check list of the Ethiopian Muscicapidae (Sylviinae). Part 1. *Occasional papers of the National Museums of Southern Rhodesia* 3 (24b): 399-430.
- _____ 1961. *A revised check list of African broadbills, pittas, larks, swallows, wagtails and pipits.* Lusaka: Government Printer.
- _____ 1962a. A check list of the Ethiopian Muscicapidae (Sylviinae). Part II & III. *Occasional papers of the National Museums of Southern Rhodesia* 3 (26B): 653-738.
- _____ 1962b. *A revised check list of African shrikes, orioles, drongos, starlings, waxwings, cuckoo shrikes, bulbuls accentors, thrushes and babblers.* Lusaka: Government Printer.
- _____ 1963. *A revised check list of African flycatchers, tits, tree creepers, sunbirds, white-eyes, honey eaters, buntings, finches and waxbills.* Lusaka: Government Printer.
- _____ 1965. *A revised check list of African non-passerine birds.* Lusaka: Government Printer.
- WOODMAN, H.N. 1936. Rare birds of the Zande District. *Sudan Notes and Records* 19: 184-188.
- _____ 1938. More rare birds of the southern Bahr-el-Ghazal. *Sudan Notes and Records* 21: 315-324.
- _____ 1951. Ducks and geese in the Southern Sudan. *Sudan Wildlife* 2: 36.

INDEX OF SCIENTIFIC NAMES

Numbers refer to species, not pages

- abdimi, Ciconia 25
- aberrans, Cisticola 520
- abingoni, Campethera 371
- abyssinica, Alcippe 428
- abyssinica, Coracias 325
- abyssinica, Hirundo 396
- abyssinicus, Bucorvus 346
- abyssinicus, Turdus 495
- Accipiter 74
- Acrocephalus 499
- Acryllium p.23
- Actitis 175
- Actophilornis 156
- acuta, Anas 41
- adsimilis, Dicrurus 410
- adusta, Muscicapa 576
- aegyptius, Caprimulgus 276
- aegyptius, Pluvianus 202
- aequatorialis, Apus 290
- aequatorialis, Sheppardia 491
- aereus, Ceuthmochares 261
- aeruginosus, Circus 65
- aethiopica, Hirundo 397
- aethiopica, Threskiornis 34
- aegyptiacus, Alopochen 40
- afer, Euplectes 683
- afer, Nilaus 618
- afer, Ptilostomus 420
- afer, Turtur 233
- affinis, Apus 291
- affinis, Sarothrura 145
- afra, Pytilia 761
- africanus, Actophilornis 156
- africanus, Bubo 268
- africanus, Buphagus 654
- africanus, Gyps 59
- africanus, Phalacrocorax 5
- Agapornis 238
- aguimp, Motacilla 604
- alba, Calidris 187
- alba, Egretta 16
- alba, Motacilla 605
- alba, Platalea 35
- alba, Tyto 266
- alboterminatus, Tockus 338
- albicapilla, Cossypha 467
- albicauda, Mirafra 388
- albiceps, Vanellus 168
- albiceps, Psalidoprocne 405
- albicollis, Ficedula 573
- albicollis, Merops 313
- albifrons, Amblyospiza 680
- albifrons, Pentholaea 486
- albigularis, Phyllastrephus 454
- albinucha, Columba 219
- albipectus, Trichastoma 430
- albirostris, Bubalornis 715
- albiventris, Parus 421
- alboterminatus, Tockus 338
- albus, Corvus 416
- Alcedo 303
- alcinus, Macheiramphus 106
- Alcippe 428
- Alethe 461
- alexandrinus, Charadrius 159
- alfredi, Brachypterus 512
- alleni, Porphyrio 142
- alopez, Falco 108
- Alopochen 40
- alpina, Calidris 188
- Amadina 768
- Amandava 739
- Amblyospiza 680
- amethystina, Nectarinia 661
- Anaplectes 681
- Anas 41
- Anastomus 24
- Andropadus 442
- angolensis, Dryoscopus 609
- angolensis, Gypohierax 58
- angulata, Gallinula 140
- Anhinga 6
- Anomalospiza 682
- Anthreptes 656
- Anthropoides 136
- Anthus 597
- Apalis 506
- Apaloderma 300
- apiaster, Merops 314
- apivorus, Pernis 103
- Aplopelia 218
- Apus 290
- apus, Apus 292
- aquatica, Muscicapa 577
- Aquila 81
- arabs, Otis 154
- Ardea 9
- ardens, Euplectes 684
- Ardeola 13
- ardesiaca, Egretta 17
- ardosiaceus, Falco 109
- Arenaria 195
- arnaudi, Pseudonigrita 720
- arquatrix, Columba 220

Artomyias 569
 arundinaceus, Acrocephalus 499
 asiaticus, Charadrius 160
 Asio 267
 astrild, Estrilda 743
 aterrimus, Phoeniculus 331
 atrata, Ceratogymna 337
 atricapilla, Sylvia 560
 atricollis, Ortygospiza 759
 atrogularis, Serinus 781
 audeberti, Pachyoccyx 260
 augur, Buteo 87
 auguralis, Buteo 88
 auratus, Oriolus 412
 auritus, Nettapus 52
 australis, Tchagra 620
 australis, Treron 236
 avosetta, Recurvirostra 197
 axillaris, Euplectes 685
 ayresii, Cisticola 521
 Aythya 49

 baboecala, Brachypterus 513
 badia, Halcyon 306
 badiceps, Eremomela 539
 badius, Accipiter 74
 badius, Ploceus 693
 Baeopogon 446
 baeticatus, Acrocephalus 500
 baglaffeht, Ploceus 694
 Balaeniceps 22
 Balearica 137
 Barbarus, Laniarius 611
 barbatus, Apus p.44
 barbatus, Gypaetus p.15
 barbatus, Pycnonoyus 459
 Bathmocercus 511
 Batis 583
 baumanni, Phyllastrephus 455
 bellicosus, Polemaetus 98
 bengalus, Uraeginthus 765
 benghalensis, Rostratula 158
 biarmicus, Falco 110
 Bias 587
 bicolor, Dendrocygna 38
 bicolor, Lonchura 769
 bicolor, Ploceus 695
 bidentatus, Lybius 348
 blanchoti, Malaconotus 616
 Bleda 447
 blissetti, Platysteira 592
 bodessa, Cisticola 522
 bollei, Phoeniculus 332
 bonapartei, Gymnobucco 347
 bonelli, Phylloscopus 549
 borin, Sylvia 561

 Bostrychia 32
 Botaurus p.8
 bottae, Oenanthe 480
 brachydactyla, Calandrella 383
 brachyptera, Cisticola 523
 brachyura, Sylvieta 566
 Bradornis 570
 Brachypterus 512
 brevicaudata, Camaroptera 515
 brevipes, Accipiter 75
 brevis, Bycanistes 335
 breweri, Merops 315
 Bubalornis 715
 Bubo 268
 Bucorvus 346
 bulocki, Merops 316
 Buphagus 654
 Burhinus 198
 Butastur 86
 Buteo 87
 buteo, Buteo 89
 Butorides 15
 Bycanistes 335

 cabanisi, Emberiza 774
 caerulescens, Muscicapa 578
 caerulescens, Rallus 144
 caeruleus, Elanus 105
 caesia, Coracina 440
 caffer, Apus 293
 caffra, Cossypha 468
 cailliautii, Campethera 375
 Calandrella 383
 Calidris 187
 calurus, Criniger 452
 Camaroptera 515
 camelus, Struthio 1
 camerunensis, Hypochera 729
 Campephaga 437
 campestris, Anthus 597
 Campethera 371
 camurus, Tockus 339
 canicapilla, Nigrita 758
 canicollis, Serinus 782
 canorus, Cuculus 256
 cantans, Cisticola 524
 cantillans, Mirafra 389
 capensis, Anas p.14
 capensis, Asio 267
 capensis, Burhinus 198
 capensis, Corvus 417
 capensis, Euplectes 686
 capensis, Microparra 157
 capensis, Oena 224
 capicola, Streptopelia 225
 capistrata, Nesocharis 757

Caprimulgus 276
 caprius, Chrysococcyx 249
 carbo, Phalacrocorax p.7
 cardinalis, Quelea 712
 caroli, Campethera 372
 caspia, Sterna 213
 castanea, Platysteira 590
 castanopterus, Passer p.91
 caudata, Coracias p.47
 caudatus, Lamprotornis 641
 Centropus 262
 Ceratogymna 337
 Cercomela 463
 Cercotrichas 464
 cervinus, Anthus 598
 Ceryle 301
 Ceuthmochares 261
 chalcopertus, Rhinoptilus 203
 chalcospilos, Turtur
 chalcurus, Lamprotornis 642
 chalybaeus, Lamprotornis 643
 chalybeata, Hypochera 730
 Charadrius 159
 Chelictinia 104
 chelicuti, Halcyon 307
 cherrug, Falco 111
 chinensis, Coturnix 120
 chiniana, Cisticola 525
 chicquera, Falco 112
 Chlidonias 210
 chloris, Nicator 453
 Chlorocichla 449
 chloronota, Camaroptera 516
 Chloropeta 518
 chloropterus, Lamprotornis 644
 chloropus, Gallinula p.25
 chloropygia, Nectarinia 662
 chokolatina, Melaenornis 574
 Chrysococcyx 249
 chrysoconus, Pogoniulus 357
 Ciccaba 270
 Cichladusa 466
 Ciconia 25
 ciconia, Ciconia 26
 cincta, Riparia 407
 cinctus, Rhinoptilus 204
 cinerascens, Circaetus 70
 cinerea, Apalis 506
 cinerea, Ardea 9
 cinerea, Creatophora 640
 cinerea, Motacilla 606
 cinereola, Cisticola 526
 cinereus, Circaetus 71
 cinereus, Xenus 182
 cinnamomeiventris, Thamnolaea 494
 cinnamomeus, Bradypterus 514

Cinnyrinclus 638
 Circaetus 70
 Circus 65
 cirrocephalus, Larus p.33
 Cisticola 520
 citrenelloides, Serinus 783
 Clamator 253
 clamosus, Cuculus 257
 clanga, Aquila p.21
 clappertoni, Francolinus 123
 clara, Motacilla p.78
 clarus, Caprimulgus 277
 climacurus, Caprimulgus 278
 clypeata, Anas 42
 Clytospiza 740
 coccinigaster, Nectarinia 663
 Colius 298
 collaris, Anthreptes 656
 collaris, Lanius 625
 collurio, Lanius 626
 collybita, Phylloscopus 550
 Columba 219
 comitata, Muscicapa 579
 communis, Sylvia 562
 concolor, Falco p.21
 Coracias 325
 Coracina 440
 cordofanica, Mirafr 390
 coronatus, Stephanoaetus 99
 coronatus, Vanellus 169
 corvina, Corvinella 624
 Corvinella 624
 Corvus 416
 Corythaeola 242
 Corythaixoides 243
 Cossypha 467
 Coturnix 120
 coturnix, Coturnix 121
 crassirostris, Vanellus 170
 crassus, Poicephalus 239
 Creatophora 640
 crecca, Anas 43
 Crex 138
 crex, Crex 138
 Crinifer 244
 Criniger 452
 cristata, Alcedo 303
 cristata, Corythaeola 242
 cristata, Fulica 25
 croceus, Macronyx 603
 cruentus, Rhodophoneus 619
 crumeniferus, Leptoptilus 30
 Cryptospiza 742
 cucullata, Lonchura 770
 cuculoides, Aviceda p.21
 Cuculus 256

cuprea, Nectarinia 664
cupreus, Chrysococcyx 250
curruca, Sylvia 563
cursor, Cursorius 200
Cursorius 200
curvirostris, Andr padus 442
cuvieri, Falco 113
cyanea, Platysteira 591
cyanocampter, Cossypha 469
cyanocephalus, Uraeginthus 766
cyanogaster, Coracias 326
Cypsiurus 296
darnaudii, Trachyphonus 361
daurica, Hirundo 398
decepiens, Streptopelia 226
delegorguei, Columba 221
delegorguei, Coturnix 122
Delichon 395
Dendrocygna 38
Dendropicos 378
denhami, Neotis 152
dentata, Petronia 726
deserti, Oenanthe 481
diademata, Alethe 461
Dicrurus 410
Dinemellia 717
dinemelli, Dinemellia 717
domesticus, Passer 721
donaldsoni, Caprimulgus 279
dorsalis, Lanius 627
dorsostriatus, Serinus 784
Drymocichla 537
Dryocichloides 474
dubius, Charadrius 161
dubius, Hieraaetus p.21
dybowskii, Clytospiza 740

ecaudatus, Terathopius 73
edoloides, Melaenornis 575
edouardi, Guttera 132
egregia, Crex
Egretta 16
Elanus 105
elegans, Sarothrura 146
eleonora, Falco p.21
Emberiza 774
emini, Ploceus 696
Eminia 538
eminibey, Passer 722
Empidonis 572
Ephippiorhynchus 29
episcopus, Ciconia 27
epops. Upupa 330
Erannornis 593
Eremomela 539
Eremopterix 384

erythrocephalus, Trachyphonus 362
erythroceria, Nectarinia 665
erythronotos, Estrilda 744
erthrophthalma, Netta 51
erythroops, Cisticola 527
erythroops, Quelea 713
erythroptera, Heliolais 542
erythropygia, Mirafra 391
erythrorhynchus, Anas 44
erythrorhynchus, Tockus 340
erythrothoraxm Stiphornis 492
Estrilda 743
Euplectes 683
Eupodotis 148
Eurocephalus 636
europaeus, Caprimulgus 280
Eurystomus 329
excubitor, Lanius 628
excubitorius, Lanius 629
exilis, Indicator 363
eximia, Bleda 447
exustus, Pterocles 215

falcinellus, Limicola 192
falcinellus, Plegadis 33
Falco 108
familiaris, Cercomela 463
famosa, Nectarinia 666
fasciata, Amadina 768
fasciatus, Tockus 341
fasciinucha, Falco 114
ferina, Aythya p.14
ferruginea, Calidris 189
ferrugineus, Laniarius 612
Ficedula 573
fischeri, Turdus 496
fischeri, Vidua 733
flammeus, Asio
flammulata, Megabyas 589
flava, Campephaga 437
flava, Motacilla 607
flavicans, Macrosphenus 546
flavicollis, Chlorocichla 449
flavida, Apalis 507
flavigaster, Hyliota 588
flavigularis, Chrysosomus 251
flavirostra, Limnocorax 141
flavirostris, Rhynchops 214
flavirostris, Tockus 342
flaviventris, Emberiza 775
forbesi, Charadrius 162
forbesi, Emberiza 776
fraenatus, Caprimulgus 281
fransiscanus, Euplectes 687
Francolinus 123
fraseri, Stizorhina 494

fringilloides, *Lonchura* 771
frontalis, *Sporopipes* 728
fulgidus, *Onychognathus* 648
Fulica p.15
fuliginosa, *Artomyias* 569
fuligula, *Aythya* 49
fuligula, *Hirundo* 399
fulvescens, *Trichastoma* 431
funebri, *Laniarius* 613
funereus, *Parus* 422
fuscescens, *Dendropicos* 377
fuscus, *Larus*

gabar, *Melierax* 50
galactotes, *Cercotrichas* 464
galactotes, *Cisticola* 529
galbula, *Ploceus* 697
Galerida 387
gallicus, *Circaetus* 72
Gallinago 183
gallinago, *Gallinago* 183
Gallinula 140
gambagae, *Muscicapa* 580
gambensis, *Dryoscopus* 610
gambensis, *Plectropterus* 53
garrulus, *Coracias* 327
garzetta, *Egretta* 18
Gelochelidon 212
gierowii, *Euplectes* 688
glandarius, *Clamator* 253
Glareola 205
glareola, *Tringa* 177
Glaucidium 271
glaucus, *Eurystomus* 329
goertae, *Mesopicos* 379
goliath, *Ardea* 10
Gorsachius 20
gracillirostris, *Acrocephalus* 501
gracillirostris, *Andropadus* 443
grillii, *Centropus* 262
griseicapilla, *Lonchura* 772
griselidis, *Acrocephalus* 502
griseopyga, *Hirundo* 400
griseus, *Passer* 723
gubernator, *Lanius* 630
guifsobalito, *Lybius* 349
guinea, *Columba* 222
gularis, *Cuculus* 258
gularis, *Serinus* 785
guttata, *Cichladusa* 466
Guttera 132
Gymnobucco 347
Gypaetus p.15
Gypohierax 58
Gyps 59

habessinica, *Nectarinia* p.85
hagedash, *Bostrychia* 32
Halcyon 306
Haliaeetus 101
haliaeetus, *Pandion* 107
hartlaubi, *Pteronetta* 54
hartlaubii, *Eupodotis* 148
Heliolais 542
hemprichii, *Tockus* 343
heuglini, *Cossypha* 470
heuglini, *Neotis* 153
heuglini, *Ploceus* 698
hiaticula, *Charadrius* 163
Hieraetus 92
Himantopus 196
himantopus, *Himantopus* 196
Hippolais 543
hirsutus, *Lybius* 350
hirundineus, *Merops* 317
Hirundo 396
hispanica, *Oenanthe* 482
hordeaceus, *Euplectes* 689
hortensis, *Sylvia* 564
hortulana, *Emberiza* p.97
horus, *Apus* p.44
hottentota, *Anas* 45
hunteri, *Nectarinia* p.85
hybridus, *Chlidonias* 210
Hylia 544
Hyliota 588
hypermetra, *Mirafr* 392
Hypochoera 729
hypocherina, *Vidua* 734
hypoleucos, *Actitis* 175
ianthinogaster, *Uraeginthus* 767
ibis, *Ardeola* 14
ibis, *Mycteria* 31
icterina, *Hippolais* p.72
icteropygialis, *Eremomela* 540
icterorhynchus, *Francolinus* 124
imberbis, *Anomalospiza* 682
incana, *Drymocichla* 537
Indicator 363
indicator, *Baepogon* 446
indicator, *Indicator* 364
inornatus, *Caprimulgus* 282
insignis, *Ploceus* 699
insignis, *Prodotiscus* 368
interjecta, *Vidua* 735
intermedia, *Egretta* 19
intermedius, *Ploceus* 700
interpres, *Arenaria* 195
isabellina, *Oenanthe* 483
Ispidina 311
Ixobrychus 8

- jacksoni, Apalis 508
 jacksoni, Ploceus 701
 jacksoni, Tockus 344
 jacobinus, Clamator 254
 jamesi, Tchagra 621
 jamesoni, Platysteira blissetti 592
 juncidis, Cisticola 530
 Jynx 369

 Kaupifalco 94
 klaas, Chrysococcyx 252
 kori, Otis 155
 krameri, Psittacula 241

 lacrymosus, Lybius 351
 lacteus, Bubo 269
 laetissima, Chlorocichla 450
 lagonosticta 750
 lamelligerus, Anastomus 24
 Lamprotornis 641
 languida, Hippolais p.72
 Laniarius 611
 Lanius 625
 Larus 208
 larvata, Aplopelia 218
 larvata, Lagonosticta 750
 larvatus, Oriolus 413
 lateralis, Cisticola 531
 lathamii, Francolinus 125
 latirostris, Andropadus 444
 lecontei, Ispidina 311
 lepida, Emina 538
 Leptoptilos 30
 leschenaultii, Charadrius p.27
 leucocephala, Halcyon
 leucocephalus, Lybius
 leucogaster, Cinnyricinclus 638
 leucogaster, Corythaixoides 243
 leucolaima, Pogoniulus 358
 leucolophus, Tauraco 246
 leucomelas, Parus 423
 leucomelos, Lybius 353
 leuconotus, Gorsachius 20
 leuconotus, Thalassornis 56
 leucophrys, Anthus 599
 leucophrys, Cercotrichas 465
 leucopogon, Prinia 554
 leucopterus, Chlidonias 211
 leucopygius, Serinus 786
 leucopygius, Turdoides 433
 leucorodia, Platalea 36
 leucoscepus, Francolinus 126
 leucotis, Eremopterix 384
 leucotis, Otus 272
 leucotis, Tauraco
 leucurus, Vanellus 171

 levaillantii, Clamator 255
 levaillantoides, Francolinus 127
 lichtensteinii, Pterocles 216
 Limicola 192
 Limnicorax 141
 Limosa 193
 limosa, Limosa 193
 Linurgus 780
 Locustella 545
 locustella, Ortygospiza p.95
 Lonchura 769
 longicausa, Erannornis 593
 longipennis, Macrodipteryx 288
 longuemarei, Anthreptes 657
 Lophæetus 95
 ludwigii, Dicrurus 411
 luehderi, Laniarius 615
 lugens, Parisoma 547
 lugens, Streptopelia 227
 Luscinia 475
 luscinia, Luscinia 475
 luscinioides, Locustella 545
 luteolus, Ploceus 702
 luteus, Passer 724
 Lybius 348
 Lymnocyptes 186

 Macheiramphus 106
 Macrodipteryx 288
 Macronyx 603
 Macrosphenus 546
 macroura, Vidua 736
 macrourus, Circus 66
 macrourus, Euplectes 690
 macrourus, Urocolius 299
 macrourus, Urotriorchis 100
 maculatus, Indicator 365
 mahali, Plocepasser 718
 malabarica, Lonchura 773
 Malaconotus 616
 malimbica, Halcyon 309
 Malimbus 691
 Mandingoa 756
 mariquensis, Nectarinia 667
 maxima, Ceryle 301
 media, Gallinago 184
 Megabyas 589
 megarhynchos, Luscinia 476
 meiffreni, Ortyxelos 134
 Melaenornis 574
 melanocephala, Ardea 11
 melanocephalus, Lybius 354
 melanocephalus, Ploceus 703
 melanogaster, Eupodotis 149
 melanogaster, Ploceus 704
 melanoleucus, Accipiter 76

melanotis, Estrilda 745
 melanotos, Sarkidiornis 55
 melba, Apus 294
 melba, Pytilia 762
 meleagris, Numida 133
 Melierax 96
 mentalis, Sphenoeacus 558
 Merops 313
 Mesopicos 379
 metabates, Melierax 97
 metallicus, Anthreptes p.84
 meyeri, Poicephalus 240
 Microparra 157
 microrhynchus, Bradornis 570
 migrans, Milvus 102
 Milvus 102
 minimus, Lymnocyptes 186
 minor, Batis 583
 minor, Indicator 366
 minor, Lanius 632
 minor, Phoeniculus 333
 minullus, Accipiter 77
 minuta, Calidris 190
 minuta, Tchagra 622
 minutus, Ixobrychus 7
 Miraфра 388
 modesta, Galerida 387
 molitor, Batis 584
 monachus, Centropus 263
 monachus, Neophron 61
 mongolus, Charadrius 164
 monogrammicus, Kaupifalco 94
 monteiri, Clytosipiza 741
 Monticola 477
 morio, Onychognathus 649
 Motacilla 604
 motitensis, Passer 725
 mozambicus, Serinus 787
 Muscicapa 576
 musculus, Remiz 424
 musicus, Bias 587
 Musophaga 245
 Mycteria 31
 Myioparus 582
 myioptilus, Schoutedenapus 297
 Myrmecocichla 480

naevia, Coracias 328
 namaquus, Thripias 382
 nana, Cisticola 532
 narina, Apaloderma 300
 nasutus, Tockus 345
 natalensis, Caprimulgus 283
 natalensis, Chloropeta 518
 natalensis, Cisticola 533
 natalensis, Cossypha 471

naumanni, Falco p.21
 nebularia, Tringa 178
 Nectarinia 661
 Neophron 61
 Neotis 152
 Nesocharis 757
 Netta 51
 Nettapus 52
 niansae, Apus p.44
 Nicator 453
 niger, Bubalornis 716
 niger, Chlidonias p.33
 nigeriae, Hypochera 731
 nigerrimus, Ploceus 705
 nigra, Ciconia 28
 nigra, Myrmecocichla 479
 nigriceps, Eremopterix 385
 nigricollis, Ploceus 706
 nigripennis, Gallinago 185
 nigripennis, Oriolus 414
 Nigrita 758
 nigromitratus, Trochocercus 595
 Nilais 618
 nilotica, Gelochelidon 212
 nipalensis, Aquila 81
 nisoria, Sylvia 565
 nitens, Malimbus 691
 nitens, Trochocercus 596
 nitidula, Mandingoa 756
 niveicapilla, Cossypha 472
 nivosa, Campethera 373
 nonnula, Estrilda 746
 nordmanni, Glareola 205
 novaeseelandiae, Anthus 600
 nubica, Campethera 374
 nubicus, Caprimulgus p.42
 nubicus, Lanius 633
 nubicus, Merops 318
 nuchalis, Glareola 206
 Numida 133
 Nycticorax 21
 nycticorax, Nycticorax 21
 nyroca, Aythya 50

obsoletus, Picoides 381
 occipitalis, Lophæetus 95
 occipitalis, Trigoniceps 64
 ochropus, Tringa 179
 ocularis, Ploceus 707
 oedinemus, Burhinus p.31
 Oena 224
 oenanthe 480
 oenanthe, Oenanthe 484
 olivacea, Nectarinia 668
 olivaceus, Linurgus 780
 onocrotalus, Pelecanus 3

Onychognathus 648
oreobates, *Merops* 319
orientalis, *Anthreptes* 658
orientalis, *Batis* 585
orientalis, *Merops* 320
orientalis, *Vidua* 737
Oriolus 412
oriolus, *Oriolus* 415
Ortygospiza 759
Ortyxelos 134
osea, *Nectarinia* 669
ostrinus, *Pyrenestes* 760
Otis 154
Otus 272
ovampensis, *Accipiter* p.21

Pachyococcyx 260
pallida, *Hippolais* 543
pallidus, *Apus* 295
pallidus, *Bradornis* 571
paludicola, *Estrilda* 747
paludicola, *Riparia* 408
Pandion 107
paradisaea, *Vidua* 738
Parisoma 547
Parus 421
parva, *Porzana* p.24
parvulus, *Remiz* 425
parvus, *Cypsiurus* 296
Passer p.91
pavonina, *Balearica* 137
pectoralis, *Caprimulgus* 284
pectoralis, *Coracina* 441
pecuarius, *Charadrius* 165
Pelecanus 3
pelegrinoides, *Falco* 115
peli, *Scotopelia* 175
pelios, *Turdus* 497
penelope, *Anas* 46
pennatus, *Hieraaetus* 92
Pentholaea 486
percnopterus, *Neophron* 62
peregrinus, *Falco* 116
perkeo, *Batis* 586
perlatum, *Glaucidium* 271
permista, *Campethera* 375
Pernis 103
persicus, *Merops* 321
Petronia 726
petrosus, *Ptilopachus* 131
Phalacrocorax 5
Philomachus 194
phoenicea, *Campephaga* 438
phoenicoptera, *Pytilia* 763
Phoenicopterus 37
Phoeniculus 331

Phoenicurus 487
phoenicurus, *Phoenicurus* 487
Phyllastrephus 454
Phyllolais 548
Phylloscopus 549
piaggiae, *Turdus* 498
Picoides 381
picta, *Ispidina* 312
placidus, *Phyllastrephus* 456
Platalea 35
platurus, *Anthreptes* 659
Platysteira 590
platyura, *Schoenicola* 557
plebejus, *Turdoides* 434
Plectopterus 53
Plegadis 33
pleschanka, *Oenanthe* 485
Plocepasser 718
Ploceus 693
plumata, *Prionops* 637
plumbeus, *Myioparus* 582
Pluvialis 167
Pluvianus 202
podobe, *Cercotrichas* p.63
poecilolaemus, *Dendropicos* 378
poecilosterna, *Mirafra* 393
Poeoptera 651
Poginiulus 357
Pogonocichla 488
Poicephalus 239
Polemaetus 98
Polihierax 119
poliocephala, *Alethe* 462
poliocephalus, *Caprimulgus* 285
poliogaster, *Zosterops* 678
polioptera, *Emberiza* 777
poliopterus, *Melierax* p.21
Polyboroides 69
pomarina, *Aquila* 82
Porphyrio 142
porphyrio, *Porphyrio* p.25
Porzana 143
porzana, *Porzana* 143
prasina, *Hylia* 544
pratincola, *Glareola* 207
preussi, *Nectarinia* 670
Prinia 554
Prionops 637
pristoptera, *Psalidoprocne* 406
Prodotiscus 368
Psalidoprocne 405
Pseudonigrita 720
Psittacula 241
Pterocles 215
Pteronetta 54
Ptilopachus 131

Ptilostomus 420
Ptyrticus 429
pugnax, *Philomachus* 194
pulchella, *Nectarinia* 671
pulchella, *Phyllolais* 548
pulchra, *Apalis* 509
pulchra, *Sarothrura* 147
pullaria, *Agapornis* 238
punctifrons, *Remiz* 426
punctuligera, *Campethera* 376
purpuratus, *Trachylaemus* 360
purpurea, *Ardea* 12
purpureus, *Lamprotornis* 645
purpureus, *Phoeniculus* 334
purpuropterus, *Lamprotornis* 646
pusilla, *Eremomela* 541
pusilla, *Porzana* p.25
pusillus, *Merops* 322
pusillus, *Pogoniulus* 359
puveli, *Trichastoma* 432
Pycnonotus 459
pygargus, *Circus* 67
Pyrenestes 760
pyrgita, *Petronia* 727
Pytilia 762

quadribrachys, *Alcedo* 304
quadricinctus, *Pterocles* 217
Quelea 712
quelea, *Quelea* 714
querquedula, *Anas* 47
quisqualina, *Campephaga* 439

radiatus, *Polyboroides* 69
ralloides, *Ardeola* 13
Rallus 144
ranivorus, *Circus* 68
rapax, *Aquila* 83
rara, *Lagonosticta* 751
rectirostris, *Anthreptes* 660
Recurvirostra 197
reichardi, *Serinus* 788
Remiz 424
rex, *Balaeniceps* 22
Rhinoptilus 203
ripidurus, *Corvus* 418
rhodopareia, *Lagonosticta* 752
Rhodophoneus 619
rhodopyga, *Estrilda* 748
Rhynchops 214
ridibundus, *Larus* 209
riocourii, *Chelictinia* 104
Riparia 407
riparia, *Riparia* 409
robusta, *Cisticola* 534
rolleti, *Lybius* 355

rossae, *Musophaga* 245
Rostratula 158
ruber, *Phoenicopterus* 37
rubescens, *Nectarinia* 672
rubetra, *Saxicola* 490
rubiginosus, *Ploceus* 708
rubiginosus, *Turdoides* 435
rubricata, *Lagonosticta* 753
rubriceps, *Anaplectes* 681
rubricollis, *Malimbus* 692
rudis, *Ceryle* 302
rueppelli, *Eurocephalus* 636
rueppelli, *Gyps* 60
rufa, *Anhinga* 6
rufescens, *Acrocephalus* 503
rufescens, *Pelecanus* 4
ruficapilla, *Spermophaga* 764
ruficeps, *Cisticola* 535
ruficollis, *Corvus* 419
ruficollis, *Jynx* 369
ruficollis, *Tachybaptus* 2
ruficrista, *Eupodotis* 150
rufifrons, *Spiloptila* 559
rufinus, *Buteo* 90
rufipennis, *Butastur* 86
rufipennis, *Trichastoma* p.60
rufiventris, *Accipiter* 78
rufocinerea, *Monticola* 477
rufocinnamomea, *Mirafraga* 394
rufogularis, *Apalis* 510
rufopicta, *Lagonosticta* 754
rufus, *Bathmocercus* 511
rupicoloides, *Falco* p.21
rustica, *Hirundo* 401

Sagittarius 57
Salpornis 427
salvadorii, *Cryptospiza* 742
Sarkidiornis 55
Sarothrura 145
saxatilis, *Monticola* 478
Saxicola 489
scandens, *Phyllastrephus* 457
schlegelii, *Francolinus* 128
Schoenicola 557
schoenobaenus, *Acrocephalus* 504
Schoutedenapus 297
schuetti, *Tauraco* 248
scirpaceus, *Acrocephalus* 505
scops, *Otus* 273
Scopus 23
scotocerca, *Cercomela* p.63
Scotopelia 275
seimundi, *Nectarinia* 673
semipartitus, *Empidonax* 572
semirufa, *Cossypha* 473

semirufa, *Hirundo* 402
 semitorquata, *Alcedo* 305
 semitorquata, *Streptopelia* 228
 semitorquatus, *Polihierax* 119
 senator, *Lanius* 634
 senegala, *Lagonosticta* 755
 senegala, *Tchagra* 623
 senegalensis, *Burhinus* 199
 senegalensis, *Centropus* 264
 senegalensis, *Ephippiorhynchus* 29
 senegalensis, *Eupodotis* 151
 senegalensis, *Halcyon* 310
 senegalensis, *Hirundo* 403
 senegalensis, *Nectarinia* 674
 senegalensis, *Otus* 274
 senegalensis, *Streptopelia* 229
 senegalensis, *Zosterops* 679
 senegallus, *Vanellus* 172
 sephaena, *Francolinus* 129
 Serinus 781
 serpentarius, *Sagittarius* 57
 sharpii, *Cynnyricinclus* 639
 shelleyi, *Spreo* 652
 Sheppardia 491
 sibilatrix, *Phylloscopus* 551
 signata, *Eremopterix* 386
 similis, *Anthus* 601
 similis, *Chloropeta* 519
 simplex, *Camaroptera* 517
 simplex, *Chlorocichla* 451
 smithii, *Hirundo* 404
 solitarius, *Cuculus* 259
 solitarius, *Monticola* p.65
 somalica, *Prinia* 555
 somalicus, *Lanius* 635
 sparsa, *Anas* 48
 Spermothaga 764
 Sphenoeacus 558
 spilogaster, *Hieraaetus* 93
 spilonota, *Salpornis* 427
 Spiloptila, 559
 spinosus, *Vanellus* 173
 splendidus, *Lamprotornis* 647
 Sporopipes 728
 Spreo 652
 squamatus, *Francolinus* 130
 squatarola, *Pluvialis* 167
 stagnatilis, *Tringa* 180
 stellaris, *Botaurus* p.8
 stellata, *Pogonocichla* 488
 stellatus, *Caprimulgus* 286
 Stephanoaetus 99
 Sterna 213
 Stiphornis 492
 Stizorhina 493
 strepera, *Anas* p.14
 strepitans, *Phyllastrephus* 458
 Streptopelia 225
 streptophorus, *Francolinus* p.23
 striata, *Muscicapa* 581
 striatus, *Butorides* 15
 striatus, *Colius* 298
 striolatus, *Serinus* 789
 striolata, *Emberiza* 778
 Struthio 1
 stuhlmanni, *Poeyoptera* 651
 sturmi, *Ixobrychus* 8
 subbuteo, *Falco* 117
 subcylindricus, *Bycanistes* 336
 subflava, *Amandava* 739
 subflava, *Prinia* 556
 sulfurepectus, *Malaconotus* 617
 superbus, *Spreo* 653
 superciliiaris, *Camaroptera* p.69
 superciliosus, *Centropus* 265
 superciliosus, *Merops* 323
 superciliosus, *Plocepasser* 719
 superciliosus, *Ploceus*
 sylvatica, *Turnix* 135
 Sylvia 560
 Sylvieta 566
 syndactyla, *Bleda* 448
 Tacazze, *Nectarinia* 675
 tachardus, *Buteo* 91
 tachiro, *Accipiter* 79
 Tachybaptus 2
 taeniopterus, *Ploceus* 710
 tahapisi, *Emberiza* 779
 Tauraco 246
 Tchagra 620
 tectus, *Vanellus* 174
 Telecanthura p.44
 temminckii, *Calidris* 191
 temminckii, *Cursorius* 201
 tenebrosus, *Turdoides* 436
 tenellus, *Tmetothylacus* 608
 tephronotus, *Glaucidium* p.41
 Terathopius 73
 Terpsiphone 594
 Thalassornis 56
 Thamnotalae 494
 Threskiornis 34
 Thripas 382
 tinnunculus, *Falco* 118
 Tmetothylacus 608
 Tockus 338
 Torgos 63
 torquata, *Saxicola* 490
 Torquilla, *Jynx* 370
 totanus, *Tringa* 181
 tousenellii, *Accipiter* 80

tracheliotus, Torgos 63
Trachylaemus 360
Trachyphonus 361
Treron 236
Trichastoma 430
tricolor, Charadrius 166
tricolor, Pycnonotus 460
Trigonoceps 64
Tringa 177
Tristigma, Caprimulgus 287
trivialis, Anthus 601
trochilus, Phylloscopus 552
Trochocercus 595
troglodytes, Cisticola 536
troglodytes, Estrilda 749
turdinus, Ptyrticus 429
Turdoides 433
Turdus 495
Turnix 135
Turtur 232
turtur, Streptopelia 230
tympanistria, Turtur 235
Tyto 266

ultramarina, Hypochera
umbretta, Scopus 23
umbrovirens, Phylloscopus 553
undulata, Anas p.14
unicincta, Columba 223
Upupa 330
Uraeginthus 765
urbica, Delichon 395
Urocolius 299
Urotriorchis 100
ussheri, Telecanthura p.44

Vanellus 168
variegatus, Indicator 367
variegatus, Merops 324
velatus, Ploceus 711
venusta, Nectarinia 676
verreauxi, Aquila 84
verticalis, Nectarinia 677
verspertinus, Falco p.21
vexillarius, Macrodipteryx 289
Vidua 733
viduata, Dendrocygna 39
vielloti, Lybius 356
vinacea, Streptopelia 231
virens, Andropadus 445
virens, Sylvietta 567
virgo, Anthropoides 136
viridis, Terpsiphone 594
vocifer, Haliaeetus 101
vulturinum, Acryllium p.23

waalia, Treron 237
wahlbergi, Aquila 85
walleri, Onychognathus 650
whytii, Sylvietta 568
wilsoni, Hypochera 730
woodfordii, Ciccaba 270

xantholophus, Mesopicos 380
Xenus 182

zonurus, Crinifer 244
Zosterops 678

INDEX OF ENGLISH NAMES

Numbers refer to species, not pages

- Akalet, Equatorial 492
- Alethe, Brown-chested 462
 - Fire-crested 461
- Apalis, Black-collared 509
 - Black-throated 508
 - Buff-throated 510
 - Grey 506
 - Yellow-breasted 507
- Avocet 197
- Babbler, African Hill 428
 - Brown 434
 - Dusky 436
 - Thrush 429
 - White-rumped 433
- Barbet, Black-billed 349
 - Black-breasted 355
 - Black-throated 354
 - d'Arnaud's 361
 - Double-toothed 348
 - Grey-throated 347
 - Hairy-breasted 350
 - Red and Yellow 362
 - Red-fronted 353
 - Spotted-flanked 351
 - Vieillot's 356
 - White-headed 352
 - Yellow-billed 360
- Bateleur 73
- Batis, Black-headed 583
 - Chinspot 584
 - Grey-headed 585
 - Pygmy 586
- Bee-eater, Black-headed 315
 - Blue-breasted 324
 - Blue-cheeked 321
 - Carmine 318
 - Cinnamon-chested 319
 - Eurasian 314
 - Little 322
 - Little Green 320
 - Madagascar 323
 - Red-throated 316
 - Swallow-tailed 312
 - White-throated 313
- Bishop, Black 688
 - Black-winged Red 689
 - Red 687
 - Yellow 686
 - Yellow-crowned 683
- Bittern p.8
 - Dwarf 8
 - Little 7
- Blackcap 560
- Bluebill, Red-headed 764
- Bluethroat p.64
- Boubou, Slate-coloured 613
 - Sooty 614
 - Tropical 611
- Bristlebill 448
 - Green-tailed 447
- Brownbul, Northern 458
- Brubru, Northern 618
- Bulbul, Common 459
 - Yellow-vented 460
- Bunting, Brown-rumped 776
 - Cabanis' 774
 - Cinnamon-breasted Rock 779
 - Cretzschmar's p.97
 - Golden-breasted 775
 - House 778
 - Somali Golden-breasted 777
- Bustard, Arabian 154
 - Black-bellied 149
 - Buff-crested 150
 - Denham's 152
 - Hartlaub's 148
 - Heuglin's 153
 - Kori 155
 - Senegal 151
- Button Quail 135
- Buzzard, Augur 87
 - Common 89
- Grasshopper 86
- Honey 103
- Lizzard 94
- Long-legged 90
- Mountain 91
- Red-necked 88
- Camaroptera, Grey-backed 515
 - Olive-green 516
 - Yellow-browed p.69
- Canary, White-bellied 784
 - Yellow-crowned 782
 - Yellow-fronted 787
- Chat, Brown-tailed Rock p. 63
 - Cliff 480
 - Red-tailed 463
 - Rufous Bush 464

Chat, sooty 481
 White-fronted Black 479
 Chatterer, Rufous 435
 Chiffchaff 550
 Cisticola, Ashy 526
 Black-backed 528
 Boran 522
 Croaking 533
 Foxy 536
 Rattling 525
 Red-faced 527
 Red-pate 535
 Rock-loving 520
 Siffling 523
 Singing 524
 Stout 534
 Tiny 532
 Whistling 531
 Winding 529
 Wing-snapping 521
 Zitting 530
 Citril, African Grey-faced 783
 Coot, Red-knobbed p.25
 Cordon-bleu, Blue-capped 766
 Red-cheeked 765
 Cormorant p.7
 Long-tailed 5
 Corncrake 138
 Coucal, Black 262
 Blue-headed 263
 Senegal 264
 White-browed 265
 Courser, Cream-coloured 200
 Heuglin's 204
 Temminck's 201
 Violet-tipped 203
 Crane, African 139
 Baillon's p.25
 Black 141
 Buff-spotted Pygmy 146
 Chestnut-tailed Pygmy 145
 Little p.24
 Spotted 143
 White-spotted Pygmy 147
 Crane, Crowned 137
 Demoiselle 136
 Creeper, spotted 427
 Crimson-wing, Abyssinian 742
 Crombec 566
 Green 567
 Red-faced 568
 Crow, Pied 416
 Cuckoo, African 258
 Black 257
 Black and White 254
 Didric 249
 Emerald 250
 Eurasian 256
 Great Spotted 253
 Klaas' 252
 Levallant's 255
 Red-chested 259
 Thick-billed 260
 Yellow-throated Green 251
 Cuckoo Shrike, Black 437
 Grey 440
 Purple-throated 439
 Red-shouldered 438
 White-breasted 441
 Curlew, Stone p.31
 Cut-throat 768
 Darter 6
 Dove, Black-billed Wood 232
 Dusky Turtle 227
 Blue-spotted Wood 233
 Emerald-spotted Wood 234
 Laughing 229
 Lemon 218
 Mourning 226
 Namaqua 224
 Red-eyed 228
 Ring-necked 225
 Tambourine 235
 Turtle 230
 Vinaceous 231
 Drongo 410
 Square-tailed 411
 Duck, Black 48
 Fulvous Whistling 38
 Hartlaub's 54
 Red-billed 44
 Tufted 49
 White-backed 56
 White-faced Whistling 39
 Yellow-billed p.14
 Dunlin 188
 Eagle, African Hawk 93
 Ayres' Hawk p.21
 Banded Snake 70
 Beaudouin's Snake 72
 Black-chested 72
 Booted 92
 Brown Snake 71
 Crowned 99
 Fish 101
 Greater Spotted p.21

Lesser Spotted 82
 Long-crested Hawk 95
 Martial 98
 Short-toed Snake 72
 Steppe 81
 Tawny 83
 Verreaux's 84
 Wahlberg's 85
 Egret, Cattle 14
 Great White 16
 Little 18
 Yellow-billed 19
 Eremomela, Brown-crowned 539
 Green-backed 541
 Yellow-bellied 540
 Falcon, Barbary 115
 Eleonora's p.21
 Lanner 110
 Peregrine 116
 Pygmy 119
 Red-footed p.21
 Red-necked 112
 Saker 111
 Sooty p.21
 Taita 114
 Finch, Locust p.95
 Oriole 780
 Quail 759
 Firefinch, African 753
 Bar-breasted 754
 Black-bellied 751
 Black-faced 750
 Jameson's 752
 Red-billed 755
 Fiscal 625
 Grey-backed 629
 Somali 635
 Taita 627
 Flamingo, Greater 37
 Lesser p.11
 Flycatcher, Ashy 578
 Black 575
 Black and White 587
 Blue 593
 Blue-headed Crested 596
 Collared 573
 Dusky 576
 Dusky Blue 579
 Dusky Crested 595
 Gambaga 580
 Grey 570
 Lead-coloured 582
 Pale 571
 Paradise 594
 Shrike 589
 Sooty 569
 Spotted 581
 Swamp 577
 White-eyed Slaty 574
 Yellow-bellied 588
 Francolin, Banded 128
 Clapperton's 123
 Crested 129
 Forest 125
 Heuglin's 124
 Ring-necked p.23
 Scaly 130
 Gadwall p.14
 Gallinule, Allen's 142
 Purple p.25
 Garganey 42
 Go-away Bird, White-bellied 243
 Godwit, Black-tailed 193
 Gonolek, Black-headed 612
 Goose, Egyptian 40
 Knob-billed 55
 Pygmy 52
 Spur-winged 53
 Goshawk, African 79
 Dark Chanting 97
 Gabar 96
 Pale Chanting p.21
 Red-chested 80
 Grebe, Little 2
 Greenbul, Cameroon Sombre 442
 Honeyguide 446
 Joyful 450
 Little 445
 Olive Mountain 456
 Red-tailed 452
 Simple 451
 Slender-billed 443
 Toro Olive 455
 White-throated 454
 Yellow-whiskered 444
 Greenshank 178
 Grenadier, Purple 767
 Greywing, Archer's 127
 Smith's 127
 Guineafowl, Crested 132
 Tufted 133
 Vulturine p.23
 Gull, Black-headed 209
 Grey-headed p.33
 Lesser Black-backed 208
 Hadada 32
 Hamerkop 23
 Harrier, African Marsh 68

Marsh 65
 Montagu's 62
 Pallid 66
 Hawk, Bat 106
 Cuckoo p.21
 Harrier 69
 Long-tailed 100
 Heron, Black 17
 Black-headed 11
 Goliath 10
 Green-backed 15
 Grey 9
 Night 21
 Purple 12
 Squacco 13
 White-backed Night 20
 Hobby 117
 African 113
 Honeybird, Cassin's 368
 Honeyguide, Black-throated 364
 Least 363
 Lesser 366
 Scaly-throated 367
 Spotted 365
 Hoopoe 330
 Abyssinian Wood 333
 Black Wood 331
 Green Wood 334
 White-headed Wood 332
 Hornbill, Abyssinian Ground 346
 Black and White Casqued 336
 Black-wattled 337
 Crowned 338
 Grey 345
 Hemprich's 343
 Jackson's 344
 Pied 341
 Red-billed 340
 Red-billed Dwarf 339
 Silvery-cheeked 335
 Yellow-billed 342
 Hylia, Green 544
 Ibis, Glossy 33
 Sacred 34
 Illadopsis, Brown 431
 Pale-breasted p.60
 Puvel's 432
 Scaly-breasted 430
 Indigobird, Bar-breasted Firefinch 732
 "Camerunensis" 729
 "Nigeriae" 731
 Red-billed Firefinch 730
 Jacana 156
 Lesser 157
 Kestrel 118
 Fox 108
 Grey 109
 Lesser p.21
 White-eyed p.21
 Kingfisher, Blue-breasted 309
 Chestnut-bellied 308
 Chocolate-backed 306
 Dwarf 311
 Giant 301
 Half-collared 305
 Malachite 303
 Pied 302
 Pygmy 312
 Shining-blue 304
 Striped 307
 Woodland 310
 Kite, Black 102
 Black-shouldered 105
 Swallow-tailed 104
 Lammergeyer p.15
 Lapwing, Crowned 169
 Long-toed 170
 Lark, Chestnut-backed Sparrow 384
 Chestnut-headed Sparrow 386
 Crested p.54
 Flappet 116
 Kordofan Bush 390
 Northern White-tailed Bush 388
 Pink-breasted 393
 Red-tailed Bush 391
 Red-winged Bush 392
 Short-toed 383
 Singing Bush 389
 Sun 387
 White-fronted Sparrow 385
 Leaflove 457
 Yellow-throated 449
 Longbill, Yellow 546
 Longclaw, Yellow-throated 603
 Lovebird, Red-headed 238
 Malimbe, Blue-billed 691
 Red-headed 692
 Mannikin, Black and White 769
 Bronze 770
 Magpie 771
 Marabou 30
 Martin, African Rock 399
 African Sand 408
 Banded Sand 407
 House 395
 Sand 409

Moorhen p.25
 Lesser 140
 Mousebird, Blue-naped 299
 Speckled 298
 Negrofinch, Grey-headed 758
 Nicator 453
 Nightingale 476
 Nightjar, Montane 285
 Donaldson-Smith's 279
 Dusky 281
 Egyptian 276
 Eurasian 280
 Fiery-necked 284
 Freckled 287
 Long-tailed 278
 Nubian p.42
 Pennant-winged 289
 Plain 282
 Rufous-cheeked p.42
 Slender-tailed 277
 Standard-winged 288
 Star-spotted 286
 White-tailed 283
 Olive-back, Grey-headed 758
 Oriole, African Golden 412
 Black-headed 413
 Black-winged 414
 Golden 415
 Ortolan p.97
 Osprey 107
 Ostrich 1
 Owl, African Marsh 267
 African Scops 274
 African Wood 270
 Barn 266
 Pel's Fishing 275
 Long-eared p.41
 Scops 273
 Short-eared P.41
 Spotted Eagle 268
 Verreaux's Eagle 269
 White-faced Scops 272
 Owlet, Pearl-spotted 271
 Red-chested p.41
 Oxpecker, Red-billed 655
 Yellow-billed 654
 Parisoma, Brown 547
 Parakeet, Rose-ringed 241
 Parrot, Brown 240
 Niam Niam 239
 Partridge, Stone 131
 Pelican, Pink-backed 4
 White 3
 Petronia, Bush 726
 Yellow-spotted 727
 Piapiac 420
 Pigeon, Afep 223
 Bronze-naped 221
 Bruce's Green 237
 Green 236
 Olive 220
 Speckled 222
 White-naped 219
 Pintail 41
 Pipit, Golden 608
 Long-billed 601
 Plain-backed 599
 Red-throated 598
 Richard's 600
 Tawny 597
 Tree 602
 Plantain Eater, Eastern Grey 244
 Plover, Black-headed 174
 Caspian 160
 Egyptian 202
 Forbe's 162
 Grey 167
 Kentish 159
 Little Ringed 161
 Quail 134
 Ringed 163
 Spur-winged 173
 Three-banded 166
 Wattled 172
 White-headed 168
 White-tailed 171
 Pochard, European p.14
 Southern 51
 White-eyed 50
 Pratincole, Common 207
 Black-winged 205
 White-collared 206
 Prinia, Pale 555
 Tawny-flanked 556
 White-chinned 554
 Puffback 609
 Pink-footed 610
 Pytilia, Green-winged 762
 Orange-winged 761
 Red-winged 763
 Quail 121
 Blue 120
 Harlequin 122
 Quelea, Cardinal 712
 Red-billed 714
 Red-headed 713

- Rail, African Water 144
 Raven, Brown-necked 419
 Fan-tailed 418
 Redshank 181
 Spotted 176
 Redstart 488
 Robin, Black Bush p.63
 Forest 493
 Grey-winged Ground 474
 White-browed Scrub 465
 White-starred Forest 489
 Robinchat 468
 Blue-shouldered 469
 Red-capped 471
 Rüppell's 473
 Snowy-headed 472
 White-browed 470
 White-crowned 467
 Roller, Abyssinian 325
 Blue-bellied 326
 Broad-billed 329
 Eurasian 327
 Lilac-breasted p.47
 Rufous-crowned 328
 Rook, Cape 417
 Roughwing, Black 406
 White-headed 405
 Ruff 194

 Sanderling 187
 Sandgrouse, Chestnut-bellied 215
 Four-banded 217
 Lichtenstein's 216
 Sandpiper, Broad-billed 192
 Common 175
 Curlew 189
 Green 179
 Marsh 180
 Terek 182
 Wood 177
 Sandplover, Great p.27
 Kittlitz's 165
 Mongolian 164
 Secretary Bird 57
 Seedcracker, Large-billed 760
 Seed-eater, Streaky 789
 Streaky-headed 785
 Stripe-breasted 788
 White-rumped 786
 Yellow-rumped 781
 Shikra 74
 Shoebill 22
 Shoveler 42

 Shrike, Emin's 630
 Great Grey 628
 Grey-headed Bush 616
 Helmet 637
 Lesser Grey 632
 Lühder's 615
 Nubian 633
 Red-backed 626
 Red-tailed 631
 Rosy-patched 619
 Sulphur-breasted Bush 617
 White-crowned 636
 Woodchat 634
 Yellow-billed 624
 Silverbill 773
 Grey-headed 772
 Silverbird 572
 Skimmer 214
 Snipe, African 185
 Common 183
 Great 184
 Jack 186
 Painted 158
 Sparrow, Chestnut 722
 Grey-headed 723
 House 721
 Parrot-billed 723
 Rufous 725
 Somali p.91
 Sudan Golden 724
 Sparrowhawk, Great 76
 Levant 75
 Little 77
 Ovampo p.21
 Rufous 78
 Spinetail, Mottle-throated p.44
 Spoonbill 36
 African 35
 Sprosser 475
 Spurfowl, Yellow-necked 126
 Starling, Blue-eared Glossy 643
 Brown-tailed Glossy 642
 Chestnut-winged 648
 Lesser Blue-eared Glossy 644
 Long-tailed Glossy 641
 Purple Glossy 645
 Red-winged 649
 Rüppell's Long-tailed Glossy 646
 Sharpe's 639
 Shelley's 652
 Splendid Glossy 647
 Stuhlmann's 651

Superb 653
 Violet-backed 638
 Waller's Chestnut-winged 650
 Wattled 640
 Stilt, Black-winged 196
 Stint, Little 190
 Temminck's 191
 Stonechat 491
 Stork, Abdim's 25
 Black 28
 Open-billed 24
 Saddle-billed 29
 White 26
 Wooly-necked 27
 Yellow-billed 31
 Sunbird, Amethyst 661
 Beautiful 671
 Collared 656
 Copper 664
 Eastern Pygmy p.84
 Eastern Violet-backed 658
 Green 660
 Green-headed 677
 Green-throated 672
 Hunter's p.85
 Little Green 673
 Malachite 666
 Mariqua 667
 Northern Double-collared 670
 Northern Orange-tufted 669
 Olive 668
 Olive-bellied 662
 Pygmy 659
 Red-chested 665
 Scarlet-chested 674
 Shining p.85
 Splendid 663
 Tacazze 675
 Variable 676
 Violet-backed 657
 Swallow 401
 Ethiopian 397
 Grey-rumped 400
 Mosque 403
 Red-rumped 398
 Rufous-chested 402
 Striped 396
 Wire-tailed 404
 Swift, African Black p.44
 Alpine 294
 Common 292
 Horus p.44
 Little 291
 Mottled 290
 Nyanza p.44
 Pallid 295
 Palm 296
 Scarce 297
 White-rumped 293
 Tchagra, Black-capped 622
 Black-headed 623
 Brown-headed 620
 Three-streaked 621
 Teal 43
 Hottentot 45
 Tern, Black p.33
 Caspian 212
 Gull-billed 213
 Whiskered 210
 White-winged Black 211
 Thicknee, Senegal 199
 Spotted 198
 Thrush, Abyssinian Ground 498
 African 497
 Blue Rock p.65
 Little Rock 477
 Olive 495
 Rock 478
 Rufous 494
 Spotted Ground 496
 Spotted Morning 466
 Tinkerbird, Lemon-rumped 358
 Red-fronted 359
 Yellow-fronted 357
 Tit, Black 423
 Dusky 422
 Mouse-coloured Penduline 424
 Sennar Penduline 426
 White-bellied 421
 Yellow Penduline 425
 Trogon, Narina's 300
 Turaco, Black-billed 248
 Great Blue 242
 Ross's 245
 White-cheeked 247
 White-crested 246
 Turnstone 195
 Twinspot, Brown 741
 Dybowski's 740
 Green-backed 756
 Vulture, Egyptian 62
 Hooded 61
 Lappet-faced 63
 Palm-nut 58
 Rüppell's 60
 White-backed 59
 White-headed 64

Wagtail, African Pied 604
 Grey 606
 Mountain p.78
 White 605
 Yellow 607
 Warbler, African Reed 500
 Bamboo 512
 Barred 565
 Basra Reed 502
 Black-faced Rufous 511
 Bonelli's 549
 Brown Woodland 553
 Buff-bellied 548
 Cinnamon Bracken 514
 Fan-tailed 557
 Garden 561
 Great Reed 499
 Greater Swamp 503
 Grey-capped 538
 Grey Wren 517
 Icterine p.72
 Lesser Swamp 501
 Little Rush 513
 Marsh p.67
 Mountain Yellow 519
 Moustached 558
 Olivaceous 543
 Orphean 564
 Reed 505
 Red-faced 560
 Red-fronted 559
 Red-winged 542
 Red-winged Grey 537
 Savi's 545
 Sedge 504
 Upcher's p.72
 Willow 552
 Wood 551
 Yellow 518
 Wattle-eye 591
 Chestnut 590
 Jameson's 592
 Waxbill 743
 Black-cheeked 744
 Black-crowned 746
 Black-rumped 749
 Crimson-rumped 748
 Fawn-breasted 747
 Yellow-bellied 745
 Zebra 739
 Weaver, Baglafaecht 694
 Black 705
 Black-billed 704
 Black-necked 706
 Brown-capped 699
 Chestnut 708
 Chestnut-crowned Sparrow 719
 Cinnamon 693
 Compact 709
 Dark-backed 695
 Emin's 696
 Golden-backed 701
 Grey-headed Social 720
 Grosbeak 680
 Heuglin's Masked 698
 Little 702
 Masked 700
 Northern Masked 710
 Parasitic 682
 Red-billed Buffalo 716
 Red-headed 681
 Rüppell's 697
 Speckle-fronted 728
 Spectacled 707
 Vitelline Masked 711
 White-billed Buffalo 715
 White-browed Sparrow 718
 White-headed Buffalo 717
 Yellow-backed 703
 Wheatear, Black-eared 484
 Desert 483
 Isabelline 485
 Northern 486
 Pied 487
 Red-breasted 482
 Whinchat 490
 White-eye, Montane 678
 Yellow 679
 Whitethroat 562
 Lesser 563
 Whydah, Northern Broad-tailed
 Paradise 735
 Northern Paradise 737
 Paradise 738
 Pin-tailed 736
 Steel-blue 734
 Straw-tailed 733
 Widowbird, Fan-tailed 685
 Red-collared 684
 Red-naped 684
 White-winged p.87
 Yellow-mantled 690
 Wigeon 46
 Cape p.14
 Woodpecker, Bearded 382
 Brown-backed 381
 Brown-eared 372
 Buff-spotted 373

Cardinal 377
Fine-spotted 376
Golden-tailed 371
Green-backed 375
Grey 379
Nubian 374

Uganda Spotted 378
Yellow-crested 380
Wryneck 370
Red-throated 369
Yellowbill 261





